



BIOACTIVE COMPOUNDS AND BIOLOGICAL ACTIVITIES
OF SOME MEDICINAL PLANTS FROM CENTRAL NEPAL

M.Sc. Thesis

2016

Submitted to

CENTRAL DEPARTMENT OF BIOTECHNOLOGY

Tribhuvan University

Kirtipur, Kathmandu, Nepal

By

Gyandra Prasad Ghimire

Exam Roll No.: BT105/069

Registration No.: 5-2-37-540-2007



BIOACTIVE COMPOUNDS AND BIOLOGICAL ACTIVITIES OF SOME MEDICINAL PLANTS FROM CENTRAL NEPAL

M.Sc. Thesis

2016

Submitted to

Central Department of Biotechnology

Tribhuvan University

Kirtipur, Kathmandu, Nepal

By

Gyandra Prasad Ghimire

Registration No.: 5-2-37-540-2007

Supervisors

Prof. Dr. Krishna Das Manandhar

Prof. Dr. Krishna Kumar Shrestha

Acknowledgements

I am deeply indebted to my research supervisor, Prof. Dr. Krishna Das Manandhar, for his kind encouragement, patient supervision, insightful advice, enduring support, and creative suggestions. He has taught me to think critically, select problems, solve them, and present their solutions. I would also like to express my gratitude to my another supervisor, Prof. Dr. Krishna Kumar Shrestha, Former Head, Central Department of Botany, for offering me an opportunity to work in a research project entitled “Bioprospecting of underutilised plants of Dhading district” which was conducted in the Central Department of Biotechnology, in collaboration with Ethnobotanical Society of Nepal (ESON), Kathmandu, and Korea Research Institute of Bioscience and Biotechnology (KRIBB), South Korea. My sincere gratitude goes to Prof. Dr. Rajani Malla, Head, Central Department of Biotechnology for her persistent love and guidance. She rendered her valuable time and effort and helped arrange this work for me.

I would like to thank Tribhuvan University Teaching Hospital (TUTH), Maharajgunj, Kathmandu for providing bacterial cultures of *Staphylococcus aureus*, methicillin resistant *Staphylococcus aureus*, *Klebsiella pneumoniae*, and *Salmonella* Typhi. I would like to appreciate Central Department of Chemistry, Tribhuvan University for providing *Artemia salina* eggs. Special recognition must be given to National Centre for Cell Science (NCCS), Pune for providing different cell lines including HeLa and MDCK. The author’s gratitude goes to Dr. Rabindra M. Sapkota, Mr. Bikash Pandey, Ms. Priyam Shah, and Mr. Rajendra Ghimire from Everest Biotech Pvt. Ltd., Khumaltar, Lalitpur for their technical assistance and first-class facilities on cell culture and MTT assay. I cannot go without mentioning National Forensic Science Laboratory (NAFOL), Khumaltar, Lalitpur, and Advanced Instrumentation Research Facility (AIRF), Jawaharlal Nehru University, New Delhi for offering the training, and the technical support on GC-MS analysis.

I am grateful to my respected teachers, Prof. Dr. Tribikram Bhattra, Prof. Dr. Mohan Kharel, Prof. Dr. Ganga Prasad Kharel, Prof. Dr. Tilak R. Shrestha, Dr. Giri Raj Tripathi, Ms. Jarina Joshi, Mr. Bal Hari Poudel, and Dr. Smita Shrestha, for their endless encouragement and support throughout the whole period. I would like to acknowledge Mohan Bahadur Shrestha and Ujjwal Pradhan for their assistance in the lab and in the library, respectively. Appreciation is given to Shrawan Kumar Mishra, Birendra Prasad Gupta, Bimlesh Kumar Jha and Anurag Adhikari for their support and scholarly interactions at various points of my research. My sincere thanks to Sujindra Subedi, Namita Bhattarai, Suyog Lamichhane, Yadu Nath Poudel, and Sneha Joshi. Thanks are due to my friends Rajindra Napit, Nirmal Aryal, Pradip Paudel, Ishwar katawal, Mitesh Shrestha, Roshan Nepal, Baidhyanath Kumar Jha, Santosh Dahal,

Uddhav Karki, Mukesh Thapa, Bimala Dhakal, Sandipty Kayastha, Aarati Poudel, and Suvash Chandra Awal. Here I would like to express my infinite gratitude to my mother for her unconditional love and support.

Gyandra Ghimire

List of Abbreviations

µg	Microgram
µl	Microliter
amu	atomic mass unit
ATCC	American Type Culture Collection
EC ₅₀	Effective concentration 50
CDBT	Central Department of Biotechnology
CDB	Central Department of Botany
CDC	Central Department of Chemistry
DPPH	2,2-diphenyl-1-picrylhydrazyl
EB	Everest Biotech
ESON	Ethnobotanical Society of Nepal
EMEM	Eagle's Minimum Essential Medium
eV	electron volt
GAE	Gallic Acid Equivalent
GC-MS	Gas Chromatography - Mass Spectroscopy
HeLa	Human Cervical Carcinoma Cell Line
IC ₅₀	Inhibitory Concentration 50
IUCN	International Union for Conservation of Nature and Natural Resources
KRIBB	Korea Research Institute of Bioscience and Biotechnology
MDCK	Madin-Darby Canine Kidney Epithelial Cells
MDR	Multi-Drug Resistance
MHA	Mueller Hinton Agar
MRSA	methicillin resistant <i>Staphylococcus aureus</i>
MTT	3-(4,5-dimethyl thiazol-2-yl)-2,5-diphenyl tetrazolium bromide

NA	Nutrient Agar
NCCS	National Centre for Cell Science
NPHL	National Public Health Laboratory
NAFOL	National Forensic Science Laboratory
QE	Quercetin Equivalent
RSA	Radical Scavenging Activity
RT	Room Temperature
TFC	Total Flavonoid Content
TPC	Total Phenolic Content
UV	Ultraviolet
WHO	World Health Organization

Table of Contents

Acknowledgements.....	i
List of Abbreviations.....	iii
List of figures.....	ix
List of Tables.....	xi
List of Plates.....	xiii
Abstract.....	xiv
1 Introduction.....	1
1.1 Background.....	1
1.2 Medicinal Plants.....	1
1.2.1 Medicinal Plants in Nepal.....	2
1.3 Bioactive Compounds.....	2
1.3.1 Phenolic.....	3
1.3.2 Flavonoid.....	3
1.4 Biological Activities.....	4
1.4.1 Antioxidant Activities.....	4
1.4.2 Antimicrobial Effects.....	5
1.4.3 Anticancer Properties.....	6
1.4.4 Brine Shrimp Cytotoxicity.....	6
1.5 Hypothesis.....	8
1.6 Objectives.....	8
1.6.1 General Objective.....	8
1.6.2 Specific Objectives.....	8
1.7 Rationale.....	8
2 Literature Review.....	10
2.1 <i>Anemone rivularis</i> Buchanan-Hamilton ex De Candolle.....	10
2.2 <i>Chrysopogon aciculatus</i> (Retzius) Trinius.....	11

2.3	<i>Geranium wallichianum</i> D. Don ex Sweet.....	13
2.4	<i>Hydrocotyle javanica</i> Thunb.....	14
2.5	<i>Loxogramme involuta</i> (D. Don) Persl	16
2.6	<i>Phyllanthus emblica</i> Linnaeus	16
2.7	<i>Ranunculus scleratus</i> Linnaeus.....	17
2.8	<i>Rhododendron lepidotum</i> Wallich ex G. Don	18
2.9	<i>Scurrula elata</i> (Edgeworth) Danser	20
2.10	<i>Symplocos lucida</i> (Thunberg) Siebold & Zuccarini	20
2.11	<i>Terminalia bellirica</i> (Gaertner) Roxburgh	22
2.12	<i>Tsuga dumosa</i> (D. Don) Eichler	23
3	Materials and Methods	24
3.1	Collection of Plant Materials.....	24
3.2	Methanol Extraction of Plant Samples.....	26
3.3	Total Phenolic Content.....	26
3.4	Total Flavonoid Content.....	26
3.5	Antioxidant Activity.....	27
3.6	GC-MS Analysis.....	28
3.7	Antimicrobial activity	28
3.7.1	Nutrient agar (NA).....	28
3.7.2	Mueller Hinton Agar (MHA).....	29
3.7.3	Standard bacterial culture Inoculums.....	29
3.7.4	Antibacterial test.....	29
3.7.5	Potato Dextrose Agar (PDA) and Potato Dextrose Broth	30
3.7.6	Preparation of the standard fungal cultures	30
3.7.7	Antifungal test.....	30
3.8	Brine Shrimp Cytotoxicity.....	30
3.9	MTT Cell Proliferation	31
3.10	Statistical Analysis	32

4	Results.....	33
4.1	Methanol Extraction of Plant Materials.....	33
4.2	Determination of Total Phenolic Content.....	34
4.3	Determination of Total Flavonoid Content.....	35
4.4	Estimation of Antioxidant Activity.....	36
4.5	Correlation between Total Phenolic Content and Antioxidant Activity.....	38
4.6	Correlation between Total Flavonoid Content and Antioxidant activity.....	39
4.7	GC-MS Analysis.....	40
4.7.1	GC-MS Analysis of <i>Loxogramme involuta</i>	40
4.7.2	GC-MS Analysis of <i>Hydrocotyle javanica</i>	42
4.7.3	GC-MS Analysis of <i>Phyllanthus emblica</i>	45
4.7.4	GC-MS Analysis of <i>Chrysopogan aciculatus</i>	47
4.7.5	GC-MS Analysis of <i>Rhododendron lepidotum</i>	50
4.7.6	GC-MS Analysis of <i>Geranium wallichianum</i>	52
4.7.7	GC-MS Analysis of <i>Tsuga dumosa</i>	55
4.7.8	GC-MS Analysis of <i>Anemone rivularis</i>	57
4.7.9	GC-MS Analysis of <i>Scurrula elata</i>	60
4.7.10	GC-MS Analysis of <i>Terminalia bellerica</i>	63
4.7.11	GC-MS Analysis of <i>Ranunculus scleratus</i>	66
4.7.12	GC-MS Analysis of <i>Symplocos lucida</i>	69
4.8	Antimicrobial Effects.....	71
4.9	Brine Shrimp Cytotoxicity.....	75
4.10	MTT Cell Proliferation Assay.....	77
4.10.1	HeLa Cell Survival.....	78
4.10.2	MDCK Cell Survival.....	80
5	Discussion.....	83
5.1	Methanol Extraction.....	83
5.2	Total Phenolic Content.....	84

5.3	Total Flavonoid Content	86
5.4	Antioxidant Activity	87
5.4.1	Correlation of TPC and TFC with Antioxidant Activity	89
5.5	Antimicrobial Activity	89
5.6	Brine Shrimp Cytotoxicity	93
5.7	Anticancer Activities	94
5.8	GC-MS analysis	95
6	Summary and Conclusion	104
7	Recommendation	106
	References	107
	Appendices	142
	Photographs	144

List of figures

Figure 4.1 Standard graph of gallic acid.....	34
Figure 4.2 Total phenolic content in methanolic extracts of 12 medicinal plants.	35
Figure 4.3 Standard graph of quercetin.....	36
Figure 4.4 Total flavonoid content in methanolic extracts of 12 medicinal plants.....	36
Figure 4.5 DPPH free radical scavenging activity of ascorbic acid.....	37
Figure 4.6 DPPH free radical scavenging activity of methanolic extracts of 12 medicinal plants.	37
Figure 4.7 IC ₅₀ of DPPH free RSA of methanolic extracts of 12 medicinal plants.....	38
Figure 4.8 Correlation between total phenolic content and IC ₅₀ of DPPH free RSA of methanolic extracts of 12 medicinal plants.....	39
Figure 4.9 Correlation between total flavonoid content and IC ₅₀ of DPPH free RSA of methanolic extracts of 12 medicinal plants.....	39
Figure 4.10 GC-MS Chromatogram of methanolic extract of <i>Loxogramme involuta</i> whole plant.	40
Figure 4.11 GC-MS Chromatogram of methanolic extract of <i>Hydrocotyle javanica</i> whole plant.	43
Figure 4.12 GC-MS Chromatogram of methanolic extract of <i>Phyllanthus emblica</i> bark.	46
Figure 4.13 GC-MS Chromatogram of methanolic extract of <i>Chrysopogan aciculatus</i> whole plant.	48
Figure 4.14 GC-MS Chromatogram of methanolic extract of <i>Rhododendron lepidotum</i> leaf and stem.	50
Figure 4.15 GC-MS Chromatogram of methanolic extract of <i>Geranium wallichianum</i> root.	53
Figure 4.16 GC-MS Chromatogram of methanolic extract of <i>Tsuga dumosa</i> bark.	55
Figure 4.17 GC-MS Chromatogram of methanolic extract of <i>Anemone rivularis</i> whole plant.	57
Figure 4.18 GC-MS Chromatogram of methanolic extract of <i>Scurrula elata</i> branches.....	61
Figure 4.19 GC-MS Chromatogram of methanolic extract of <i>Terminalia bellerica</i> fruit.	64
Figure 4.20 GC-MS Chromatogram of methanolic extract of <i>Ranunculus scleratus</i> whole plant.	66
Figure 4.21 GC-MS Chromatogram of methanolic extract of <i>Symplocos lucida</i> bark.	69
Figure 4.22 Antimicrobial activity of <i>T. dumosa</i> by agar well diffusion method as shown by the zone of inhibition against methicillin-resistant <i>Staphylococcus aureus</i> and <i>Salmonella</i> Typhi, respectively.	72

Figure 4.23 Percentage mortality of *Artemia salina* (brine shrimp) larvae on treatment of methanolic extracts of 12 medicinal plants..... 76

Figure 4.24 LC₅₀ of methanolic extracts of 12 medicinal plants on *Artemia salina* (brine shrimp) larvae..... 77

Figure 4.25 Effects of *H. javanica* methanol extract on HeLa cell line with its decreasing concentration..... 78

Figure 4.26 Effects of methanolic extracts of 12 medicinal plants on HeLa cell survival..... 78

Figure 4.27 Effects of methanolic extracts of 12 medicinal plants on HeLa cell survival..... 79

Figure 4.28 Effects of methanolic extracts of 12 medicinal plants on MDCK cell survival..... 80

Figure 4.29 Effects of methanolic extracts of 12 medicinal plants on MDCK cell survival..... 81

List of Tables

Table 3.1 Name, collection site, parts used and ethnobotanical information of 12 medicinal plants.....	24
Table 4.1 Percentage yield and physical characteristics of the crude methanol extracts.	33
Table 4.2 Phytocomponents identified in the methanolic extract of <i>Loxogramme involuta</i> whole plant by GC-MS.	41
Table 4.3 Phytocomponents identified in the methanolic extract of <i>Hydrocotyle javanica</i> whole plant by GC-MS.	43
Table 4.4 Phytocomponents identified in the methanolic extract of <i>Phyllanthus emblica</i> bark by GC-MS.....	46
Table 4.5 Phytocomponents identified in the methanolic extract of <i>Chrysopogan aciculatus</i> whole plant by GC-MS.	48
Table 4.6 Phytocomponents identified in the methanolic extract of <i>Rhododendron lepidotum</i> leaf and stem by GC-MS.....	50
Table 4.7 Phytocomponents identified in the methanolic extract of <i>Geranium wallichianum</i> root by GC-MS.....	53
Table 4.8 Phytocomponents identified in the methanolic extract of <i>Tsuga dumosa</i> bark by GC-MS.	56
Table 4.9 Phytocomponents identified in the methanolic extract of <i>Anemone rivularis</i> whole plant by GC-MS.	58
Table 4.10 Phytocomponents identified in the methanolic extract of <i>Scurrula elata</i> branches by GC-MS.....	61
Table 4.11 Phytocomponents identified in the methanolic extract of <i>Terminalia bellerica</i> fruit by GC-MS.....	64
Table 4.12 Phytocomponents identified in the methanolic extract of <i>Ranunculus scleratus</i> whole plant by GC-MS.	66
Table 4.13 Phytocomponents identified in the methanolic extract of <i>Symplocos lucida</i> bark by GC-MS.	70
Table 4.14 Antibacterial activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against <i>Staphylococcus aureus</i> using agar well diffusion method....	72
Table 4.15 Antibacterial activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against methicillin-resistant <i>Staphylococcus aureus</i> using agar well diffusion method.....	73
Table 4.16 Antibacterial activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against <i>Klebsiella pneumoniae</i> using agar well diffusion method...	73

Table 4.17 Antibacterial activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against *Salmonella* Typhi using agar well diffusion method. 74

Table 4.18 Antifungal activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against *Saccharomyces cerevisiae* using agar well diffusion method. 74

Table 4.19 Antifungal activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against *Pichia sp.* using agar well diffusion method. 75

Table 4.20 EC₅₀ of methanolic extracts of 12 medicinal plants on HeLa cell line..... 80

Table 4.21 EC₅₀ of methanolic extracts of 12 medicinal plants on MDCK cell line..... 82

List of Plates

Plate 1 Some medicinal plants (a) <i>Loxogramme involuta</i> , (b) <i>Hydrocotyle javanica</i> Thunb, (c) <i>Phyllanthus emblica</i> Linnaeus, and (d) <i>Chrysopogon aciculatus</i> (Retzius) Trinius	144
Plate 2 Some medicinal plants (a) <i>Rhododendron lepidotum</i> Wallich ex G. Don, (b) <i>Geranium wallichianum</i> D. Don ex Sweet, (c) <i>Tsuga dumosa</i> (D. Don) Eichler, and (d) <i>Anemone rivularis</i> Buchanan-Hamilton ex De Candolle	145
Plate 3 Some medicinal plants (a) <i>Scurrula elata</i> (Edgeworth) Danser, (b) <i>Terminalia bellirica</i> (Gaertner) Roxburgh, (c) <i>Ranunculus scleratus</i> Linnaeus, and (d) <i>Symplocos lucida</i> (Thunberg) Siebold & Zuccarini	146

Abstract

Considerable attempts have been made globally to validate the traditional use of plants in phytomedicine. Phytomedicine can counter the challenge posed by drug resistance and toxicity by continuous search for novel drugs. This study focuses on the estimation of total phenolic content, total flavonoid content, examination of antioxidant activities, GC-MS identification of bioactive compounds, observation of antimicrobial, cytotoxic, and anticancer effects. Twelve medicinal plants with some history of use as traditional medicine have been selected for the purpose. They include *Loxogramme involuta*, *Hydrocotyle javanica*, *Phyllanthus emblica*, *Chrysopogan aciculatus*, *Rhododendron lepidotum*, *Geranium wallichianum*, *Tsuga dumosa*, *Anemone rivularis*, *Scurrula elata*, *Terminalia bellerica*, *Ranunculus scleratus*, and *Symplocos lucida*.

Total phenolic content was determined by the Folin-Ciocalteu method, total flavonoid content by aluminium chloride colorimetric method. Antioxidant activity was tested by their capacity to scavenge DPPH (2,2-diphenyl-1-picrylhydrazyl) free radicals. GC-MS analysis was performed to identify the bioactive compounds. The antimicrobial efficacy was tested against *Staphylococcus aureus*, methicillin-resistant *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Salmonella Typhi*, *Saccharomyces cerevisiae* and *Pichia sp.* by agar well diffusion method. Cytotoxicity was determined by brine shrimp bioassay. Anticancer activity was determined by MTT assay on HeLa and MDCK cell lines.

P. emblica (55.66 ± 0.84 mgGAE/g) possessed the highest amount of total phenolic content. Higher amount of TPC was also observed in *G. wallichianum* (54.99 ± 0.74 mgGAE/g), *S. elata* (54.71 ± 0.68 mgGAE/g), *T. bellerica* (54.69 ± 0.77 mgGAE/g), and *T. dumosa* (54.19 ± 0.93 mgGAE/g). The lowest amount of TPC was observed in *R. scleratus* (6.74 ± 0.62 mgGAE/g). The highest amount of total flavonoid content was observed in *P. emblica* (25.65 ± 0.77 mgQE/g). The lowest amount of TFC was observed in *T. bellerica* (7.51 ± 0.54 mgQE/g). TFC for other plant extracts locate within the short range between these two values. *T. dumosa* (4.54 ± 0.18 μ g/ml) showed the highest antioxidant activity with the lowest IC₅₀ value – lower than that observed in ascorbic acid (4.57 ± 00 μ g/ml). Higher antioxidant activity was also observed in *P. emblica* (6.64 ± 0.71 μ g/ml), *G. wallichianum* (7.24 ± 0.48 μ g/ml), *T. bellerica* (12.08 ± 0.76 μ g/ml), and *S. elata* (14.20 ± 0.94 μ g/ml). *R. scleratus* (195.55 ± 2.45 μ g/ml) showed the lowest antioxidant activity. A significant Pearson's correlation between TPC and IC₅₀ of DPPH RSA ($r = -0.9684$) attributed phenolic compounds to be the main contributor of antioxidant activity in plants. GC-MS analysis showed the presence of a wide range of bioactive compounds such as fatty acids, esters, sugars, steroidal, phenolic, and heterocyclic

compounds. Maximum inhibition of *S. aureus* and MRSA both were shown by *T. bellerica*. Highest inhibition of *S. Typhi* was caused by *T. dumosa*. Minimum number of plants inhibited *K. pneumoniae* on which *L. involuta* exerted the highest effect. Ten of the twelve plants inhibited at least three of the six organisms. Eight of them inhibited at least four organisms. *G. wallichianum* inhibited five of the six organisms. *S. elata* and *T. bellerica* both inhibited all of them. Plants showed higher effects on gram positive bacteria. Nine of the twelve plants showed significant toxicity to brine shrimps ($LC_{50} < 500 \mu\text{g/ml}$) of which three were found to be highly toxic ($LC_{50} < 100 \mu\text{g/ml}$) which includes *A. rivularis*, *L. involuta* and *T. bellerica*. Maximum number of plants showed higher inhibition of HeLa cell line than of MDCK. The highest inhibition of HeLa cell was shown by *S. lucida* ($52.22 \mu\text{g/ml}$) which precedes other ten with EC_{50} value less than $1000 \mu\text{g/ml}$. *T. bellerica* ($1048 \mu\text{g/ml}$) showed the lowest inhibition. Highest inhibition of MDCK cell was shown by *L. involuta* ($240 \mu\text{g/ml}$) followed by *T. dumosa* ($305 \mu\text{g/ml}$), *A. rivularis* ($407.7 \mu\text{g/ml}$), *S. elata* ($751.1 \mu\text{g/ml}$), and *P. emblica* ($925.3 \mu\text{g/ml}$). Other showed less inhibition with EC_{50} value more than $1000 \mu\text{g/ml}$.

The results provide justification for traditional use of medicinal plants in treatment of various diseases. The biological activities of medicinal plants may have been attributed due to the presence of bioactive compounds identified. These plants can be the candidates as the potential source of high valued phytopharmaceuticals. Further meticulous analysis of plants might lead to isolation, identification and characterization of biologically active substances.

Keywords: medicinal plants, phenolic, flavonoid, antioxidant, bioactive compounds, antimicrobial, cytotoxicity, anticancer

1 Introduction

1.1 Background

Recorded history of the use of medicinal plants is much shorter than the actual one which in fact began with the beginning of the human civilization. The vast majority of modern medicines have been derived originally from ancient herbal traditions. Plants are a great source of therapeutic molecules. It has been more than three billion years since nature started its own combinatorial chemistry (Holland, 1997). The total number of natural products produced by plants has been estimated at over 500,000 (Mendelson & Balick, 1995). Medicinal plants rich in phytochemicals have been used for centuries in the treatment and prevention of diseases. Phytochemicals may affect health, but are not essential nutrients. Most of the phytochemicals used in the modern pharmaceutical industry are secondary metabolites. Emerging factors like drug-resistance, cost-effectiveness and side-effects of synthetic drugs have led to the resurgence of phytomedicine.

It is estimated that the worldwide market potential for herbal drugs is around US\$40 billion (Patwardhan et al., 2004). About one-fourth of the drugs approved by FDA during the period 1981–2002 was either natural products or based on natural products (Newmann et al., 2003). About two dozen of such drugs introduced to the market during the period 2000–2005 include drugs for cancer, neurological, cardiovascular, metabolic and immunological diseases, and genetic disorders (Chin et al., 2006).

Total number of higher plant species on earth has been estimated to be more than 250,000 (Ayensu & DeFilipps, 1978). Of that only around 6% are screened for biological activity, and 15% are phytochemically characterized (Verpoorte, 2000). Fifty million screening tests have been conducted so far making a variation of molecules, concentrations and/or bioassays (Drews, 2000).

1.2 Medicinal Plants

WHO has estimated that almost 80% of the world population rely on traditional medicine for their primary health protection (Diallo et al., 1999; Kim et al., 2005). Some countries have incorporated traditional medicine extensively in their public health system (WHO, 2000). Plants continue to be the major source of medicine in rural regions of developing countries

(Chitme et al., 2004). More than 3.3 billion people have been reported to utilize medicinal plants on a regular basis (Davidson-Hunt, 2000).

The number of species of plants estimated to exist on earth has been on the order of 250,000 to 500,000 of which only 1 to 10% are used as food by humans and other animals (Borris, 1996; Cowan, 1999). Nearly 3000 species of plants have been documented to be used as food during the course of human civilization, of which only about 150 species have been cultivated (Council, 1982), and less than 10 plant species are meeting more than 90% of the world food demand (Wilkes, 1981). About 20,000 to 70,000 species of plants are used medicinally (Alves & Rosa, 2007; Farnsworth & Soejarto, 2009; Schippmann et al., 2006).

Medicinal plants have been used in treatment of different diseases for thousands of years (Palombo, 2011). Plants have been used in the treatment of cancer for a long time (Richardson, 2001). More than one thousand plants have been found to possess significant anticancer properties (Mukherjee et al., 2001). About 50% of breast cancer patients and 37% of prostate cancer patients are estimated to use herbal products (Sini et al., 2012). Phytochemicals have also been suggested to prevent colorectal cancer and other cancers (Birt et al., 2001; Yuvaraj et al., 2010).

1.2.1 Medicinal Plants in Nepal

Nepal contains about 1,950 species of medicinal plants (about 28% of the local flora) (Baral & Kurmi, 2006; Ghimire, 2008; Manandhar, 2002; Shrestha & Shrestha, 1999; Shrestha et al., 2002; Tiwari, 1999), including 1,614 native medicinal species (Ghimire et al., 2008; Shrestha et al., 2002). The contribution of medicinal plants in Nepal is important (Kunwar, 2006). More than 80% of the rural population of Nepal are reported to use herbal remedies (Dani, 1986). Herbal medicine has been used since ancient time in many parts of the country where access to formal and modern healthcare is limited (Kunwar et al., 2010).

1.3 Bioactive Compounds

The medicinal values of plants have been claimed to lie in their phytochemical components including alkaloids, tannins, flavonoids and other phenolic compounds (Anyasor et al., 2011). With the discovery of profound physiological effect exerted by active principles on mammalian systems, efforts are being made to know the exact chemical nature of compounds and to obtain by chemical synthesis (Ramawat, 2007). Natural products are a major source of inspiration for drug development (Newmann & Cragg, 2007), with only 30% of all novel molecules introduced into the market in the period 1981–2006 being pure synthetic and all others being natural products or natural product related.

1.3.1 Phenolic

Phenolic compounds and polyphenols are the most abundant structures in plants. Polyphenols are a structural class of natural and synthetic or semisynthetic, organic chemicals characterized by the presence of large multiples of phenol structural units. The number and characteristics of these phenol structures underlie the unique physical, chemical and biological (metabolic, toxic, therapeutic, etc.) properties of particular members of the class. Examples include tannic acid and ellagitannin (Chuang et al., 2013; Quideau et al., 2011). More than 8000 different polyphenols of diverse structures have been reported (Strack & Wray, 1992).

Polyphenols are antioxidants with redox properties, which allow them to act as reducing agents, hydrogen donors, and singlet oxygen quenchers. Some show metal chelation properties (Kähkönen et al., 1999; Pietta, 2000). Some have antimicrobial activity (Rauha et al., 2000). A great number of aromatic plants have been reported as having anti-inflammatory, antiallergic, antimutagenic, antiviral, antithrombotic, and vasodilatory actions (Hollman et al., 1996). Polyphenols can have favorable effects on the incidence of cancers and chronic diseases, including cardiovascular disease, type II diabetes, and impaired cognitive function (Rio et al., 2013).

Polyphenols historically have been considered as anti-nutrients by nutritionists, because some, eg tannins, have such adverse effects as decreasing the activities of digestive enzymes, energy, protein and amino acid availabilities, mineral uptake and having other toxic effects (Salunkhe et al., 1982). Recognition of the antioxidant activities of many polyphenols has realigned thinking toward the health benefits provided by many of these compounds (Bravo, 1998). The most important dietary phenolics are the phenolic acids (including hydroxybenzoic and hydroxycinnamic acids), polyphenols (hydrolyzable and condensed tannins) and flavonoids (King & Young, 1999).

1.3.2 Flavonoid

Flavonoids are secondary metabolites widely distributed in the plant kingdom. They are the most abundant polyphenols in the human diet. There are six subclasses of flavonoids including anthocyanins, flavonols, flavanols, flavanones, flavones and isoflavones. More than 6000 varieties of flavonoids have been identified in plants (Khatiwora et al., 2010). The best-described property of almost every group of flavonoids is their capacity to act as antioxidants. Among the biological activities of flavonoids are actions against free radicals, free radical mediated cellular signaling, inflammation, allergies, platelet aggregation, microbes, ulcers,

viruses and tumors and hepatotoxins (Kinsella et al., 1993). Flavonoid intake has been associated with reduced risk from death from coronary heart disease (Hertog et al., 1993).

1.4 Biological Activities

1.4.1 Antioxidant Activities

Free radicals are atoms or group of atoms with an unpaired electron. These highly reactive free radicals can start a chain reaction. Free radicals such as the reactive oxygen species (ROS) and reactive nitrogen species (RNS) are generated through endogenous processes such as metabolism, respiration and phagocytosis. They are also generated by exogenous systems such as chemical agents, pesticides, pollutants, toxins, organic solvents, radiation, cigarette smoke, deep fried foods and herbicides (Davies, 1995; Thatoi et al., 2014). Chemical compounds that can generate free radicals are called Prooxidants. In a normal cell there is a proper balance between pro-oxidants and antioxidants but when the level of pro-oxidants increases as compared to antioxidants, they lead to oxidative stress (Collins, 2005). Free radicals react with important cellular components such as DNA, and the cell membrane, and can lead to lipid peroxidation, tissue injury and protein degradation (Gilbert, 1981; Phang et al., 2011). Oxidative stress has been implicated in the etiology of a number of diseases including cancer (Kinnula & Crapo, 2004), cardiovascular disease (Singh & Jialal, 2006), neural disorders (Sas et al., 2007), Alzheimer's disease (Smith et al., 2000), mild cognitive impairment (Guidi et al., 2006), Parkinson's disease (Bolton et al., 2000), alcohol induced liver disease (Arteel, 2003), ulcerative colitis (Ramakrishna et al., 1997), aging (Hyun et al., 2006), arthritis (Ramamoorthy & Awang, 2007), atherosclerosis (Upston et al., 2003), nephritis, diabetes mellitus, rheumatism, and ischemia (Amel et al., 2013).

Antioxidants are compounds that interact with free radicals and terminate the chain reaction before the vital molecules such as lipids, proteins and DNA are damaged (Becker et al., 2004; Halliwell & Gutteridge, 1989; Kumar & Chattopadhyay, 2007; Ozsoy et al., 2008; Ratnam et al., 2006; Temitope et al., 2010). Generation of these free radicals is normally balanced by an equivalent production of antioxidants through our natural antioxidant defense mechanism, which are the enzymatic antioxidants (superoxide dismutase, glutathione peroxidase, quinone reductase and catalases) and the non-enzymatic antioxidant (ascorbic acid, α -tocopherol, melatonin, β -carotene) obtained from the diet (Halliwell, 1996; Davies, 2000; Phang et al., 2011). All aerobic organisms have antioxidant defence systems to protect from harmful effects caused by free radicals. In the case of failure of the antioxidant defence system, antioxidants need to be supplemented from outside sources which can be either

synthetic and natural. Synthetic antioxidants such as butylated hydroxyanisole, butylated hydroxytoluene, and propyl gallate have toxic effects due to which restrictions have been imposed on their use. Natural antioxidants are safe with nutritional and therapeutical effects (Lopez-Velez et al., 2003). Therefore, researchers have focused their studies on plant-derived natural antioxidants (Kulisic et al., 2004). Carotenoids, cinnamic acids, benzoic acids, folic acid, ascorbic acid, tocopherols, tocotrienols, polyphenols such as phenolics acids, flavonoids, proanthocyanidins, among others, are some of the antioxidants produced by plants for their survival (Jimoh et al., 2008; Ramamoorthy & Awang, 2007). Higher level of natural antioxidants (vitamins, carotenoids, phenolics) in human everyday diet can protect against cardiovascular diseases, cataract, cancer and aging- related disorders (Steffen et al., 2003).

1.4.2 Antimicrobial Effects

Infectious diseases represent an important cause of morbidity and mortality among the general population, particularly in developing countries. Drug resistance to pathogenic bacteria has been commonly reported from all over the world (Mulligan et al., 1993; Piddock & Wise, 1989; Singh et al., 1992). Antibiotics are sometimes associated with adverse effects on host which include hypersensitivity, depletion of beneficial gut and mucosal microorganism, immunosuppression and allergic reactions (Lopez et al., 2001). This has created immense clinical problem in the treatment of infectious diseases (Davies, 1994). Therefore, there is a need to develop alternative antimicrobial drugs for the treatment of infectious diseases. One approach is to screen local medicinal plants for possible antimicrobial properties. Plant materials remain an important resource to combat serious diseases in the world.

Major groups of antimicrobial compounds from plants include simple phenols and phenolic acids, quinones, flavones, flavonoids and flavonols, tannins, coumarins, alkaloids, terpenoids and essential oils, lectins and polypeptides (Cowan, 1999).

The action mechanisms of natural compounds are related to disintegration of cytoplasmic membrane, destabilization of the proton motive force, electron flow, active transport and coagulation of the cell content. Not all action mechanisms work on specific targets, and some sites may be affected due to other mechanisms (Burt, 2004). Important characteristics responsible for the antimicrobial action of essential oils include hydrophobic components that allow the participation of lipids from the bacterial cell membrane, which disturbs cell structures and make them more permeable (Sikkema et al., 1994). Chemical compounds from essential oils also act on cytoplasmic membrane proteins (Knobloch et al., 1989). Cyclic hydrocarbons act on ATPases, enzymes known to be located at the cytoplasmic membrane

and surrounded by lipid molecules. In addition, lipid hydrocarbons may distort the lipid-protein interaction, and the direct interaction of lipophilic compounds with hydrophobic parts of the protein is also possible (Sikkema et al., 1995). Some essential oils stimulate the growth of pseudo-mycelia, evidencing that they may act on enzymes involved in the synthesis of bacterium structural components (Conner & Beuchat, 1984).

1.4.3 Anticancer Properties

Cancer is considered to be the third leading cause of death worldwide (12.4%), the first being cardiovascular disease (30%) and the second being infectious diseases, including HIV/AIDS (18.8%) (Mathers et al., 2001). An estimated 12.7 million new cases are registered each year with 7.6 million deaths and 24.6 million persons living with cancer worldwide (Jemal et al., 2011). It is estimated that the number of deaths due to cancer may double in next 50 years and new cases of cancer may rise to 15 million by 2020 (Ferlay et al., 2015).

Cancer cells occur as a result of unique multiple genetic disorders that may arise from exposure to environmental and occupational carcinogenic agents or dietary habits and infectious agents (Sugimura, 1992). Drug resistance, particularly multidrug resistance (MDR), can make many of the clinically established anticancer drugs ineffective (Borowski et al., 2005).

Chemotherapy remains the treatment of choice in many malignant diseases (Carter & Livingston, 1982; Lage et al., 2010). Natural product could serve as chemopreventive and chemotherapeutic agent with minor side effect on normal cells and high tumor selectivity (Mehta et al., 2010). More than 60% of the anticancer drugs are of natural origin, such as plants (i.e., vincristine, irinotecan, camptothecines) and microorganisms (i.e., doxorubicin, dactinomycines, mitomycin and bleomycin) (Grever, 2001; Newman & Cragg, 2012; Newmann & Cragg, 2007; Stevigny et al., 2005). Over 50% of drugs used in clinical trials for anticancer activity were isolated from natural sources or are related to them (Newmann & Cragg, 2007). The plant-derived drugs such as taxol from *Taxus* species, vinblastine and vincristine from *Catharanthus roseus*, topotecan and irinotecan from *Camptotheca accuminata*, and etoposide and teniposide from *Podophyllum peltatum* are currently used in treatment of various types of cancers (Patwardhan et al., 2004).

1.4.4 Brine Shrimp Cytotoxicity

Although many plants have valuable properties, some of them are known to carry toxicological properties as well. Recent studies indicate that although numerous plants are used as food sources, some of them may have mutagenic or genotoxic potential (Çelik &

Aslantürk, 2007). Study on pharmacology and toxicity of medicinal plants is of high importance in order to achieve a safe treatment with plant products (Parra et al., 2001).

Various assays are used for the research of potential toxicity of herbal extracts based on different biological models, such as *in vivo* assays on laboratory animals. However, recent studies employed efforts for alternative biological assays that include species of *Artemia salina*, *Artemia franciscana*, *Artemia urmiana* and *Thamnocephalus platyurus*. These toxicity tests are considered a useful tool for preliminary assessment of toxicity (Carballo et al., 2002; Mayorga et al., 2010; Veni & Pushpanathan, 2014).

During the past 30 years, the Brine Shrimp Assay has been widely used to test the toxicity of a great variety of plant products. Brine shrimp (*Artemia salina*) is most extensively studied of the *Artemia* species, estimated to represent over 90% of the studies in which *Artemia* is used as an experimental test organism (Campbell et al., 1994). The Brine Shrimp Toxicity Assay was proposed and developed by Michael et al. (1956) and later adapted by Vanhaecke et al. (1981), Meyer et al. (1982), and Sleet & Brendel (1983).

Brine Shrimp Lethality Assay has been applied as an alternative bioassay technique to screen the toxicity of plant extracts (Gadir, 2012; McLaughlin et al., 1998; Meyer et al., 1982; Moshi et al., 2010; Ogugu et al., 2012; Sharma et al., 2013; Solanki & Selvanayagam, 2013).

Since its introduction, this *in vivo* test has been successively employed for bioassay-guided fractionation of active cytotoxic and antitumor agents (Ahmed et al., 2010; Ramachandran et al., 2011). Additionally, several studies demonstrated that there is a good correlation between the results for the lethal concentration that kills 50% of the exposed population (LC₅₀) obtained with the Brine Shrimp Lethality Assay using *A. salina* and the results of the acute oral toxicity assay in mice (Arslanyolu & Erdemgil, 2006; Parra et al., 2001).

Brine Shrimp Lethality Assay is a convenient system for monitoring biological activities of various plant species. Although this method does not provide any adequate information regarding the mechanism of toxic action, it is a very useful method for the assessment of the toxic potential of various plant extracts (Gadir, 2012; Naidu et al., 2014). This method provides preliminary screening data that can be backed up by more specific bioassays once the active compounds have been isolated (Pisutthanan et al., 2004).

Due to the ethical issues in toxicological tests, substituting animals with alternative models is very important. The effectiveness of the *Artemia salina* bioassay for predicting the toxicity of plant extracts was evaluated by comparing the LC₅₀ results for the brine shrimps with the LD₅₀

results for acute toxicity in rats and mice (Naidu et al., 2014; Parra et al., 2001; Sharma et al., 2013).

1.5 Hypothesis

People have been using plants in medicine since ancient past. They contain bioactive compounds such as phenolics and flavonoids. They possess antioxidant, antimicrobial, anticancer properties due to which the medicinal plants are being used for a number of ailments.

1.6 Objectives

1.6.1 General Objective

Study of twelve medicinal plants for the presence of bioactive compounds and biological activities.

1.6.2 Specific Objectives

1. Methanol extraction of twelve medicinal plants
2. Estimation of total phenolic and flavonoid content
3. Determination of antioxidant activity by DPPH free radical scavenging assay
4. Antibacterial property evaluation of the extracts on *S. aureus* ATCC – 25923, MRSA, *K. pneumoniae*, and *S. Typhi*
5. Antifungal property evaluation on *S. cerevisiae*, and *Pichia sp.*
6. Analysis of cytotoxicity of the extracts by brine shrimp bioassay
7. Observation of anticancer effects on HeLa and MDCK cell lines
8. Identification of bioactive compounds by GC-MS analysis

1.7 Rationale

Medicinal plants are known to contain bioactive compounds responsible for antioxidant, antimicrobial and anticancer activities. Antioxidants such as flavonoids and phenolics are associated with reduced risk of chronic diseases like heart disease, cancer, diabetes, neurodegeneration and other age related degenerative diseases. Growing problem of resistance and toxicity of presently available commercial drugs has necessitated the search for novel phytopharmaceuticals. Phytomedicine can counter the challenge posed by drug resistance and toxicity by continuous search for novel drugs. Nepal preserves highly valued medicinal plants which find use in varieties of ailments such as in cough, fever, cuts, wounds,

malaria, nephritis, hepatitis, cancer, diabetes, tooth aches, and joint pains, and so on. People are intimately connected to phytomedicine when faced with frequent diseases. Research inspired from tribal medicine contributes to drug discovery and socioeconomic development.

2 Literature Review

2.1 *Anemone rivularis* Buchanan-Hamilton ex De Candolle

Anemone rivularis (Ranunculaceae) (Plate 2) is a perennial herbaceous plant distributed in Nepal, northern India, Bhutan, Sri Lanka, southern Tibet, southwestern China and Myanmar (Manandhar, 2002). The genus *Anemone* (Ranunculaceae) consists of about 150 species with a near global distribution, of which about 50 species were found in the northwest and southwest of China (Wang, 1980). More than 10 species of this genus have been used as Chinese folk medicines for a long time. The rhizome of *A. raddeana* is recorded in the Chinese Pharmacopoeia for the treatment of rheumatism and neuralgia (National Pharmacopoeia Committee, 2010). The entire plant and rhizome of *Anemone rivularis* have been used as folk medicines for treatment of hepatitis, muscle and joint pain, stranguria, emission, edema, etc. (Wang, 1980) (Flora of China Editorial Committee of Chinese Academy of Sciences, 1980). In Nepal, paste of the plant is given to treat cough and fever, a decoction of the root is applied to cuts and wounds and juice of the leaf is inhaled to relieve sinusitis (Manandhar, 2002).

A new saponin, rivularinin, shown to be [α -L-arabinofuranosyl(1 \rightarrow 2)- α -L-rhamnopyranosyl(1 \rightarrow 4)- β -D-glucopyranosyl(1 \rightarrow 4)- β -D-glucuronopyranosyl(1 \rightarrow 3)]-3 β -hydroxy-olean-12-en-28-oic acid was isolated from the ethanolic extract of *Anemone rivularis* (Tiwari & Singh, 1978). Six saponins were first reported from *Anemone rivularis* var. *flore-minore* one of which was a new compound with gypsogenin as aglycone first reported in genus *Anemone* (Ding et al., 2011). A new triterpene ester olean-9(11),12-dien-3-O-palmitate and six known triterpenoids lupeol, betulin, betulic acid, oleanolic acid, ursolic acid and β -amyrin were isolated from *Anemone rivularis* (Zhao et al., 2012). Phytochemical investigation on the whole plant of *Anemone rivularis* var. *flore-minore* led to the isolation of a new labdane-type diterpene glycoside, β -D-glucopyranosyl (13S)-13-hydroxy-7-oxo-labda-8,14-diene-18-oate and a new trihydroxyfuranoid lignanoid glycoside (7S,7'R,8R,8'S)-7'-butoxy-7,9'-epoxy-4,4',9-trihydroxy-3,3'-dimethoxylignane 9-O- β -D-glucopyranoside, together with three known triterpene and triterpenoid glycosides 3-O- α -L-arabinopyranosyl-hederagenin, 3-O- β -D-glucopyranosyl-(1 \rightarrow 2)- α -L-arabinopyranosyl-oleanolic acid, hederagenin (Ding et al., 2012). A new cerebroside, rivulacerebroside, was isolated from *Anemone rivularis* (Shao et al., 2013). Phytochemical investigation of the *n*-BuOH extract of the rhizomes of *Anemone rivularis* var. *flore-minore* led to the isolation of five new oleanane-type triterpenoid

saponins 3-O- β -D-xylopyranosyl-(1 \rightarrow 3)- α -L-rhamnopyranosyl-(1 \rightarrow 2)-[β -D-glucopyranosyl-(1 \rightarrow 4)]- α -L-arabinopyranosyl gypsogenin 28-O- β -D-glucopyranosyl ester, 3-O- β -D-xylopyranosyl-(1 \rightarrow 3)- α -L-rhamnopyranosyl-(1 \rightarrow 2)-[β -D-glucopyranosyl-(1 \rightarrow 4)]- α -L-arabinopyranosyl gypsogenin 28-O- α -L-rhamnopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranosyl ester, 3-O- β -D-xylopyranosyl-(1 \rightarrow 3)- α -L-rhamnopyranosyl-(1 \rightarrow 2)-[β -D-glucopyranosyl-(1 \rightarrow 4)]- β -D-glucopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-(1 \rightarrow 4)]- α -L-arabinopyranosyl oleanolic acid 28-O- β -D-glucopyranosyl ester, 3-O- α -L-rhamnopyranosyl-(1 \rightarrow 2)-[β -D-glucopyranosyl-(1 \rightarrow 4)]- α -L-arabinopyranosyl 21 α -hydroxyoleanolic acid 28-O- α -L-rhamnopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranosyl ester, 3-O- β -D-glucopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-(1 \rightarrow 4)-[α -L-rhamno-pyranosyl-(1 \rightarrow 2)]- α -L-arabinopyranosyl 21 α -hydroxyoleanolic acid 28-O- α -L-rhamnopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranosyl ester, together with five known saponins sapindoside B, pulsatilla saponin D, 3-O- β -D-xylopyranosyl-(1 \rightarrow 3)- α -L-rhamnopyranosyl-(1 \rightarrow 2)-[β -D-glucopyranosyl-(1 \rightarrow 4)]- α -L-arabinopyranosyl oleanolic acid, sieboldianoside B, 3-O- α -L-arabinopyranosyl gypsogenin 28-O- α -L-rhamnopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranosyl ester (Wang et al., 2014).

The genus *Anemone* has been found to be the rich source of glycosides, especially triterpenoid glycosides with a range of biological and pharmacological activities including antitumor, antiperoxidation, antibacterial, insect deterrence (Lu et al., 2009; Mimaki et al., 2009; Sun et al., 2011; Wang et al., 2012; Wang et al., 1998; Wang et al., 1997; Ye et al., 2001; Zhang et al., 2008). The triterpene ester, olean-9(11),12-dien-3-O-palmitate, from *Anemone rivularis* was found to be effective against *B. subtilis*, *S. aureus*, *E. coli*, *S. typhimurium*, and *P. aeruginosa* (Zhao et al., 2012). The monodesmosidic saponins from *Anemone rivularis* var. *flore-minore* exhibited cytotoxic activity against HL-60 (promyelocytic leukemia), HepG2 (hepatocellular liver carcinoma), A549 (lung carcinoma) and HeLa (cervical carcinoma) human cancer cell lines (Wang et al., 2014).

2.2 *Chrysopogon aciculatus* (Retzius) Trinius

C. aciculatus (Retz.) Trin. (Plate 1) is a perennial grass with a creeping rhizome (Paria & Chattopadhyay, 2005) widely distributed in tropics of Asia in hills and plains common along the banks of rivers, streams and canals usually forming dense carpets (Dileep & Nair, 2015). Its culm is divided into creeping base and erect portion. The creeping base is covered with imbricate scale like old sheaths. Sheaths are long, striate, sometimes purple-tinged and imbricate (Singh et al., 2001). Leaf-blades are flat. Panicles are reddish purple, narrowly elliptic and long. Chorant is usually found in sunny, dry, exposed areas such as roadsides,

lawns, pasture, bank of rivers, water courses, etc. (Noltie, 2000). Cattle eat this species in default of anything else. It can tolerate grazing, mowing and trampling by animals (Kabir & Nair, 2009). It is very difficult to eradicate if it becomes established. It is a very good soil binder which prevents soil erosion (Dileep & Nair, 2015). The spikelets are easily disarticulating, acicular and the callus is extremely sharp, often gets attached to the skin and hairs of animals and clothes of human beings. Hence it is popularly called 'Love Grass' (Sreekumar & Nair, 1991).

Whole plants of *Chrysopogon aciculatus* is reported to be applied topically as decoction in treating swollen bodies, legs, arms in Sepik Province, Papua New Guinea (Koch et al., 2015). Leaves of *Saraca asoca*, *Chrysopogon aciculatus* and *Vitex negundo* are warmed and applied topically to affected areas 2-3 times daily in cases of pain and paralysis (Shahnaj et al., 2015). Rhizome of *Chrysopogon aciculatus* powdered with *Piper nigrum* and made into a paste is taken early in the morning on empty stomach for stomach ache and gastric disorders by tribal people in India (Saha et al., 2014; Dileep & Nair, 2015). The plant is used in Ayurveda as a diuretic (Johnson T, 1998). In Malaysia, the plant is burnt with lalang and the ashes swallowed as a remedy for rheumatism. The plant has been used in traditional Chinese medicine to treat common cold and fever. In China and Taiwan, a decoction of the roots is used to treat poisonous snake bites. In Indochina, the "seeds" are used to expel intestinal roundworms. In Nepal, a paste of the root is applied to boils (Manandhar, 2002) and smoke of the plant is used for healing the wound of toes caused due to mud (Rajbhandari, 2001).

Four new apigenin derivatives, 7-de-O-methylaciculatin, 8-C- β -D-boivinopyranosylapigenin, aciculatinone, and 4'-O-glucosylaciculatin, along with eight known compounds, apigenin-8-carbaldehyde, kaempferol, tricetin, taxifolin, 6,7,4'-trihydroxyflavone, *trans*-oxyresveratrol, aciculatin, and luteolin-7-sulfate, were isolated from an ethanolic extract of *Chrysopogon aciculatus* (Shen et al., 2012). Aciculatin (8-((2R,4S,5S,6R)-tetrahydro-4,5-dihydroxy-6-methyl-2H-pyran-2-yl)-5-hydroxy-2-(4-hydroxyphenyl)-7-methoxy-4H-chromen-4-one), a major bioactive component of *Chrysopogon aciculatus*, was first reported in 1991 showing DNA-binding activity and cytotoxicity *in vitro* (Carte' et al., 1991; Krause & Eggleston, 1991).

Methanol extract of *Chrysopogon aciculatus* exhibited α -glucosidase inhibitory activity (Nguyen et al., 2012). Compounds isolated from *Chrysopogon aciculatus* have been evaluated for cytotoxic activities on human cancer cell lines including MCF-7, H460, HT-29, and CEM (Shen et al., 2012). It has been found that aciculatin exerts dual inhibitory effects on inducible nitric oxide synthase (iNOS) and cyclooxygenase-2 (COX-2) due to regulation of the NF-kappaB and c-Jun N-terminal kinase (JNK)/p38 pathways (Hsieh et al., 2011). Aciculatin shows

a potent anti-inflammatory effect by inhibiting the expression of lipopolysaccharide-mediated inducible nitric oxide synthase and cyclooxygenase-2 (Shen et al., 2012). Aciculatin inhibits IL-1 β -stimulated G-CSF expression and the subsequent neutrophil differentiation, suggesting that it might have therapeutic potential for inflammatory arthritis (Shih et al., 2012). Aciculatin induces cell cycle arrest and apoptosis *via* inhibition of MDM2 expression, thereby inducing p53 accumulation without significant DNA damage and genome toxicity (Lai et al., 2012). Water extracts of *Chrysopogon aciculatus* exhibited allelopathic effects on *Borreria latifolia* (Li et al., 2012).

2.3 *Geranium wallichianum* D. Don ex Sweet

Geranium wallichianum D. Don ex Sweet (Plate 2) belongs to family Geraniaceae and is commonly known as “Shephred’s needles” (Nadkarni, 1976). It is a tall much branched procumbent or erect perennial herb found in the Himalayas, from Kashmir to Nepal at the altitude of 7000-11000 feet (Vakil, 1956; CSIR, 1968). The herb possesses the astringent properties of the genus to a marked degree. The root stock is used in eye troubles as a substitute for that of *Cotis teeta* Wall. The herb is also in the treatment of toothache (Watt, 1972). The rhizomes of the plant are used in the treatment of mouth ulceration, dysentery, diarrhea, passive hemorrhage, and leucorrhoea (Shinwari et al., 2003). Juice of the plant is applied to stop bleeding from fresh cuts and a paste is applied to relieve joint pains while root of *Geranium wallichianum* mixed with bark of *Prunus cerasoides* is used for peptic ulcer (Manandhar, 2002). *G. thunbergii* has been used in intestinal disorders in Japan (Ito et al., 1999).

The GC and GC-MS analysis of essential oil of *G. wallichianum* showed the presence of bioactive compounds such as germacrene D, β -caryophyllene, selinene < 7-*epi*- α >, δ -cadinene, α -cadinol, linalool, β -bourbonene, γ -amorphene, α -humulene, caryophyllene oxide, isophytol, β -(*E*)-farnesene, (*E,E*) geranyl linalool, terpineol-4-ol, sabinene and cubenol (Tewari et al., 2015). Tannins, Geranin, Geranic acid B and C, Phyllanthusine F, Dehydroallagittalin, Dehydrogeranin, Furosine and Furosine were isolated from *G. wallichianum* (Okuda et al., 1982). Six compounds ursolic acid, β -sitosterol, stigmasterol, β -sitosterol galactoside, herniarin, and 2,4,6-trihydroxyethylbenzoate were isolated and characterized for the first time from *Geranium wallichianum* (Ismail Mohammad et al., 2009).

The antioxidant capacity of *Geranium wallichianum* has been investigated by DPPH radical scavenging assay and EtOAc extract was found to be most potent with IC₅₀ 19.05 ug/mL (Ismail et al., 2009). Antibacterial, antifungal (Atta-ur-Rahman et al., 1991), phytotoxic

(McLaughlin et al., 1991), cytotoxic (Meyer et al., 1982) and insecticidal (Naqvi & Parveen, 1991) activities of *G. wallichianum* has been reported. The crude extracts and pure compounds isolated from rhizomes and leaves of *Geranium wallichianum* showed varied degree of antimicrobial activities (Ismail et al., 2012). The crude extract of *Geranium wallichianum* when tested for antifungal, antibacterial, insecticidal, phytotoxic and cytotoxic activities, displayed moderate activity against human, animal and plant pathogens in antifungal tests and similar activity against *Staphylococcus aureus* in antibacterial tests but not any remarkable activity in rest of the bioassays (Ahmad et al., 2003). Ursolic acid and benzoic ester from *Geranium wallichianum* were found to be capable of inhibiting urease enzyme but β -sitosterol, stigmasterol and β -sitosterol galactoside were not (Ismail et al., 2012). Lipoxygenase enzyme was more significantly inhibited as compared to urease (Ismail et al., 2012).

2.4 *Hydrocotyle javanica* Thunb

Hydrocotyle javanica Thunb (Umbelliferae) (Plate 1) is a prostrate herb, occurring from tropical to temperate zones, distributed in Nepal, northern India, Bhutan, Tibet, and Myanmar (Manandhar, 2002). Juice of the plant is used to treat fever, cough and boils (Manandhar, 2002). The genus *Hydrocotyle* contains some 130 species worldwide (Du and Ren, 2010). Most of them inhabit marshy, moist shady and understory environments in tropical and subtropical forests. The whole plants of the *Hydrocotyle* species are often used in Taiwan folk medicine for treating common cold, tonsillitis, cephalitis, enteritis, dysentery, zoster, eczema, period pain, hepatitis and jaundice (Chang et al., 2003). Traditionally, the fresh plant parts of *H. javanica* are ingested orally to cure sores of throats and lungs. Leaf juice is often used as eye drops to cure eye infection and leaf paste was used in dressing of wounds to reduce swelling and juice of shoots can treat gastritis and constipation. The leaves of *H. javanica* Thunb. are used in traditional medicine in China and Thailand as liniment on rashes (Inta et al., 2008). The poultices of *H. bonariensis* Lam. are applied externally to heal wounds, including freckles. Internally, its decoction is diuretic, antiphlogistic, hepatic, against dropsy and pectoral. The juice is emetic, diuretic, and useful against illnesses of lungs, liver, spleen and bladder (Eduardo et al., 2006). The leaf juice of *Hydrocotyle mannii* Kook.f. along with *Cynoglossum amplifolium* is used locally in cataract (Giday et al., 2009). *Hydrocotyle rotundifolia* Roxb. leaf juice is taken to cure fever and sometimes is used as brain tonic (Barukial & Sarmah, 2011). *H. sibthorpioides* is used to treat fevers, wounds, boils, abscesses, boils, cirrhosis, cold, cough, hepatitis, itching, jaundice, sinusitis and sore throat in Madhya Pradesh, India (Quamar & Bera, 2014).

Methanol extract of *Hydrocotyle javanica* showed the presence of alkaloids, flavonoids, phenols, tannins, leucoanthocyanidins and cardiac glycosides while water extract showed the presence of alkaloids, saponins and triterpenoids (Mandal et al., 2016). Five new oleanane-type triterpene saponins including steganogenin 3-O- β -D-glucopyranoside, steganogenin 3-O- α -L-rhamnopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside, steganogenin 3-O- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside, chichipegenin 3-O- α -L-rhamnopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside and chichipegenin 3-O- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside, along with four known oleanane-type triterpenes and triterpene saponins were isolated from ethanolic extract of the whole plant of *Hydrocotyle nepalensis* (Ma et al., 2016). Seven new oleanane-type triterpenoid saponins, hydrocotylosides I–VII, and one known saponin, udosaponin B, were isolated from the methanol extract of the whole plants of *Hydrocotyle sibthorpioides* (Matsushita et al., 2004).

Several studies have confirmed the pharmacological activity and therapeutic potential of the genus *Hydrocotyle* (Srinivasan et al., 2001). Water extracts of four *Hydrocotyle* species *H. nepalensis*, *H. setulosa*, *H. batrachium* and *H. sibthorpioides* showed antioxidant activities in descending order consistent with the antiproliferative activities against human hepatoma Hep3B cells suggesting *Hydrocotyle nepalensis* to be a good source of natural antioxidant among the *Hydrocotyle* genus (Huang et al., 2008). The compounds isolated from *H. nepalensis* exhibited cytotoxic activities against five human cancer cell lines (HL-60, SMMC-7721, A-549, MCF-7 and SW480) (Ma et al., 2016). *H. javanica* exhibited larvicidal activity against *Culex quinquefasciatus* (Tennyson et al., 2012), venom neutralization effect and antihemolytic activity against *Naja nigricollis* venom (Kumarappan et al., 2011), bactericidal activity against food poisoning, human pathogenic gastrointestinal and topical bacteria (Mandal et al., 2016) and a high antioxidant and genotoxic properties (Wan-Ibrahim et al., 2010). *H. sibthorpioides* inhibited the growth of transplanted tumors in mice, such as Hep (hepatic carcinoma), S180 (sarcoma) and U14 (uterine cervical carcinoma) (Yu et al., 2007), demonstrated *in vitro* and *in vivo* antiviral activities against the replication of hepatitis B virus and dengue virus (Huang et al., 2013b; Husin et al., 2015) and showed effectiveness in curing chronic alcohol-induced hepatic injury and fibrosis in rats (Huang et al., 2013a). *H. leucocephala* and *H. sibthorpioides* exhibited immunomodulatory effects (Ramos et al., 2006; Yu et al., 2007). *Hydrocotyle umbellata* exhibited analgesic, anti-inflammatory, anxiolytic and sedative effects in mice (Florentino et al., 2013; Rocha et al., 2011). *H. bonariensis* showed antioxidant activity and chemopreventive effect against oxidative stress and galactose-induced cataract (Ajani et al., 2009).

2.5 *Loxogramme involuta* (D. Don) Persl

Loxogramme involuta (Polypodiaceae) (Plate 1) is an epiphytic fern found in Nepal, India, Sri Lanka and Indo-China (Manandhar, 2002). Juice of the rhizome is used to treat cuts and wounds (Manandhar, 2002).

2.6 *Phyllanthus emblica* Linnaeus

Phyllanthus emblica Linnaeus (Euphorbiaceae) (Plate 1) is a deciduous tree indigenous to tropical South-East Asia and occurs mainly in Nepal, India, Sri Lanka, Malaysia, Myanmar (Khan, 2009; Zhang et al., 2003). Traditionally, the fruit of *P. emblica* is used to treat various ailments including diarrhoea, jaundice, inflammation, cerebral and intestinal disorders, diabetes mellitus, coronary heart disease, cancer, rheumatic pain, diseases of the eye and genitalia, gonorrhoea, constipation, asthma (Aslokar et al., 1992; Baliga & Dsouza, 2011; Perry, 1980; Rao & Siddiqui, 1964; Scartezzini & Speroni, 2000).

P. emblica contains hydrolyzable tannins Emblicanin A, Emblicanin B, punigluconin and pedunculagin (Ghosal et al., 1996). The seed contains various fatty acids like linolenic acid, linoleic acid, oleic acid, palmitic acid, stearic acid and myristic acid (Thakur et al., 1989), flavonoids such as Kaempferol-3-O- α -L-(6''-methyl) rhamnopyranoside, Kaempferol-3-O- α -L-(6''-ethyl) rhamnopyranoside (Rahman, 2007), alkaloids such as Phyllantidine and phyllantine (Khanna & Bansal, 1975). The fruit contains quercetin, Gallic acid, ellagic acid, 1-O-galloyl- β -D-glucose, 3,6-di-O-galloyl-D-glucose, chebulinic acid, chebulagic acid, corilagin and isostrictinnin (Zhang et al., 2003). A new acylated glucoside, apigenin7-O-(6''-butyryl-beta)-glucopyranoside was isolated from the methanolic extract of the leaves of *P. emblica* (El-Desouky et al., 2008). Phyllaemblic acid, a novel highly oxygenated norbisabolane were isolated from the roots of *P. emblica* (Zhang et al., 2000). Roots of *P. emblica* are also a rich source of Ellagic acid and lupeol (Kapoor, 1990; Rastogi & Mehrotra, 1993). Four new compounds 3,4,8,9,10-pentahydroxydibenzo[b,d]pyran-6-one, 3,4,3'-tri-O-methylellagic acid, lup-20,29-en-3 β ,30-diol and betulin were isolated first time from the *P. emblica* (Zhang et al., 2013).

Methanol extract of *P. emblica* exhibited the significant scavenging activity against DPPH, O₂^{•-}, OH[•] and NO radicals and also significantly inhibited the oxidation of low density lipoprotein (Nampoothiri et al., 2011). The ethyl acetate fraction of a methanolic extract of *P. emblica* fruits showed strong NO scavenging activity *in vitro* (Kumaran & Karunakaran, 2006). Extracts of *P. emblica* also exhibited significant protection to DNA against oxidative damage (Kumar et al., 2006). *P. emblica* inhibited the growth of Chinese hamster ovary cells

(CHO) (Sumantran et al., 2007), human hepatocellular carcinoma (HepG2) and lung cancer cells (A549) (Pinmai et al., 2008). *P. emblica* fruits inhibited hepatocarcinogenesis (Jeena et al., 1999), protected from sublethal gamma radiation (Singh et al., 2005), and reduced the effects of radiation (Jagetia et al., 2002). *P. emblica* inhibited cyclophosphamide-induced suppression of humoral immunity (Haque et al., 2001), chromium induced free radical production, apoptosis and DNA fragmentation (Sai Ram et al., 2003) and restored the antioxidant status. *P. emblica* showed potent antibacterial activity against *E. coli*, *K. pneumoniae*, *K. ozaenae*, *Proteus mirabilis*, *P. aeruginosa*, *S. Typhi*, *S. paratyphi* A & B, and *Serratia marcescens* (Saeed & Tariq, 2007) and also inhibited *Bacillus subtilis*, *S. Typhi*, *Bacillus cereus*, *P. aeruginosa*, *Shigella boydii*, *Shigella dysenteriae*, *S. aureus*, *Sternbergia lutea*, *E. coli*, *S. paratyphi*, *Vibrio parahaemolyticus* and *V. mimicus* (Rahman et al., 2009). *P. emblica* reduced bacterial colonisation in the lung of *Klebsiella pneumoniae*-induced mice (Saini et al., 2008). *P. emblica* inhibited the mRNA expressions of tyrosinase and related proteins (TRP-1 and TRP-2) in B16 murine melanoma cells and also suppressed the LPS-induced pro-inflammatory genes (COX-2, iNOS, TNF- α , IL-16 and IL-6) expression in RAW 264.7 murine macrophage cell (Sripanidkulchai & Junlatat, 2014).

2.7 *Ranunculus sceleratus* Linnaeus

Ranunculus sceleratus Linn. (marsh-crowfoot) (Plate 3) belonging to the family Ranunculaceae is an erect annual herb which is often found on damp terrain, riverbanks and small water bodies (Kanjilal, 1997). This species originated in the northern hemisphere and is widely distributed in Nepal, northern India, Bhutan, central Asia including Pakistan, China, Mongolia, Siberia, Japan, Europe, and North America (Manandhar, 2002). It is widely used in traditional Chinese medicine (Wu et al., 1999) due to its excellent therapeutic effects (Mei et al., 2012). The fresh or dry plant can be used to treat cancer of the esophagus and the breast (Li, 1999). *Ranunculus sceleratus* has unique toxicological and pharmacological activities. The leaves of this plant cause dermatitis, raise blisters on the skin and were formerly used by professional beggars to produce or maintain open sores and blisters to gain sympathy (Chopra et al., 1958). The plant is very acrid and poisonous, producing violent irritant effects if taken internally (Manandhar, 2002). However, the toxic properties of the plant get destroyed when consumed after boiling. The plant is considered stimulant and diuretic. Leaf juice is used in sciatica, rheumatism, dysuria, asthma and pneumonia. It is used in cutaneous disorders. Seeds are prescribed to cure kidney troubles (Ambasta et al., 1992; Gangwar & Joshi, 2008).

Ranunculuss sceleratus was investigated for pharmacologically active substances, namely histamine, acetylcholine, and 5-hydroxytryptamine, which are found in other plants that cause irritation (Bhargava et al., 1965). The various tryptamine derivatives were detected in *Ranunculus sceleratus* which are involved in the synthesis and metabolism of 5-hydroxytryptamine (Bhargava et al., 1965). Six compounds stigmasta-4-ene-3,6-dione, isoscopoletin, scoparone, protocatechuic aldehyde, protocatechuic acid and stigmasterol were isolated from *Ranunculus sceleratus* (Gao et al., 2005).

Leaf extract of *Ranunculus sceleratus* was thermostable up to 100°C and retained its antifungal activity even after autoclaving. It possessed a quick fungicidal action and even its volatile vapours were fungitoxic (Misra & Dixit, 1978). *Ranunculus sceleratus* exhibited anti-dermatophytic activity against five strains of dermatophytes *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Trichophyton tonsurans*, *Microsporum gypseum* and *Microsporum fulvum* (Sharma et al., 2012). Chloroform, ethyl acetate, *n*-butanol and aqueous fractions of *R. sceleratus* have exhibited enzyme inhibition, antimicrobial and antioxidant effects, which identifies it as a potential drug candidate for the treatment of Alzheimer's and other diseases (Shahid et al., 2015). In addition to its medicinal value, *R. sceleratus* has other potential applications. Recent studies suggest that it is capable of purifying organic sewage and the industrial wastewater containing abundance of heavy metals. *R. sceleratus* has also been considered as a potential bio-indicator of eutrophication in aquatic habitats (Xu et al., 2004).

2.8 *Rhododendron lepidotum* Wallich ex G. Don

The genus *Rhododendron* comprises more than 900 species, the majority of which are indigenous to Asia (Tiwari & Chauhan, 2005; Zhao et al., 2006). *Rhododendron lepidotum* (Ericaceae) (Plate 2) is distributed in Nepal, northern India, southeastern Tibet, southwestern China and northern Myanmar (Manandhar, 2002). It is mainly found in the Western Himalayas along with *R. campanulatum*, *R. arboretum*, and *R. anthopogon*. The genus *Rhododendron* has a long history of use in inflammation, symptoms of cold, gastrointestinal disorders, skin diseases and as pain killers (Popescu & Kopp, 2013). *Rhododendron lepidotum* has been used in headache (Khan et al., 2008; Li et al., 2000), fever, cough, cold, tonsillitis (Rajbhandari et al., 2009). *Rhododendron adamsii* Rehder is used as a stimulant and tonic (Aleksandrova, 1975; Blinova et al., 1961). Dried leaves of *R. dauricum* are used as an expectorant and in the treatment of acute and chronic bronchitis (Tang & Eisenbrand, 1992).

Rhododendron lepidotum showed the presence of flavonoids, terpenoids, steroids, glycosides, coumarins and saponins (Bhattarai et al., 2010). *Rhododendron lepidotum* afforded the isolation of 3 β -*O*-acetylup-20(29)-ene, lupeol, β -sitosterol and 3 β -hydroxy-12-ursen-28-oic acid, 3 β -olean-12-en-28-olic acid and β -*D*-fructofuranosyl- α -*D*-glucopyranoside (Lamichhane et al., 2014), rhodonin and rhodonetin (Khan et al., 2008), 7-*O*- β -*D*-glucopyranosyl-8-methoxybenzopyranone and 7-hydroxy-8-*O*- β -glycosylbenzopyranone, 7,8-dihydroxy coumarin (daphnetin) (Ahmad et al., 2010), daphnin, daphnetin, daphnetin glucoside, rhodonetin, rhodonin, and umbelliferone (Shakeel-U-Rehman et al., 2010), 8-[2',6'-dimethoxy-4'-(1'',2'',3''-trihydroxy-propyl)-phenyl]-7-hydroxy benzopyranone, 3-*O*- β -*D*-glycopyranosyl betulinic amide and 8-hydroxy-7,7'-oxydicoumarin (Shakeel-U-Rehman et al., 2013), 2,4,6-trihydroxyacetophenone-3,5-di-*C*- β -*D*-glucoside, hopenol B, lupeol, ursolic acid, avicularin, quercetin, myricetin, hyperoside, myricetin-3'-*O*- β -*D*-xyloside, (+)-taxifolin-3-*O*- α -*L*-arabinopyranoside, (+)-taxifolin-3-*O*- β -*D*-glucopyranoside, lyoniside, confluentin, 2-(4-hydroxyphenyl)-ethyltriacontanoate, 2,6-dimethoxy-4-hydroxyphenyl-1-*O*- β -*D*-glucopyranoside, (-)-2-hydroxy-5-(2-hydroxyethyl)phenyl-*O*- β -*D*-glucopyranoside, (-)-isolariciresinol, and isofraxoside (Zhou et al., 2012). Chemical constituents such as triterpene and flavanone glycosides (Takahashi et al., 2001), anti-HIV principles rhodoaurichromanin acid A and rhodoaurichromanin acid B (Kashiwada et al., 2001), diterpenoids rhodomolleins XV-XVIII, kalmanol and rhodojaponins III and VI (Li et al., 2000), dihydroflavonol glycoside (Chosson et al., 1998a), phloroacetophenone glycoside (Chosson et al., 1998b), terpenoids (Kloocke et al., 1986) and iridiod glycosides (Fan et al., 2001) were isolated from different species of *Rhododendron* genus.

Six coumarin compounds isolated from *Rhododendron lepidotum* showed better antibacterial activity than their acyl derivatives against *Staphylococcus aureus*, methicillin resistant *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* in which Daphnetin displayed the best antibacterial activity followed by rhodonetin against *S. aureus* and MRSA (Shakeel-U-Rehman et al., 2010). *Rhododendron lepidotum* showed high anti-proliferative activity against HeLa cells (Bhattarai et al., 2010), very mild to no toxicity against Vero and MDCK cells (Rajbhandari et al., 2009) and a weak activity against Influenza virus A (Rajbhandari et al., 2009). *Rhododendron lepidotum* exhibited cytotoxicity against brine shrimp larva and antimicrobial activity against *Staphylococcus aureus* (Bhattarai et al., 2010). The leaves and twigs of *R. dauricum* display significant anti-HIV activity (Kashiwada et al., 2001).

2.9 *Scurrula elata* (Edgeworth) Danser

Scurrula elata (Edgeworth) Danser (Plate 3) is a hemiparasitic plant of family Loranthaceae distributed in Nepal, India, Bhutan, Tibet, and western China. The family Loranthaceae contains about 75 genera most of which are known as mistletoes (Barlow, 1991; Calvin & Wilson, 2006). The leaves of *S. elata* are ground into a paste to relieve joint pain (Acharya & Acharya, 2009) and entire plants to hasten fracture recovery (O'Neill & Rana, 2016). The fruits of *S. elata* are edible and used for trapping birds (Acharya & Acharya, 2009; Joshi & Joshi, 2007; Kunwar & Adhikari, 2005; Limbu & Rai, 2013). Leaves are also used as fodder (Duwadee & Kunwar, 2001; Manandhar, 2002; Nepal, 1999; Shrestha, 1988, 1989).

Chemical and pharmacological studies of plants from the genus *Scurrula* led to the isolation of several bioactive compounds. Quercetin, quercitrin and glycoside 4"-O-acetylquercitrin were isolated from *Scurrula ferruginea* (Lohézic-Le et al., 2002). (Z)-9-octadecenoic acid, (Z,Z)-octadeca-9,12-dienoic acid, (Z,Z,Z)-octadeca-9,12,15-trienoic acid, octadeca-8,10-dienoic acid, (Z)-octadec-12-ene-8,10-dienoic acid, octadeca-8,10,12-trienoic acid, theobromine, caffeine, quercitrin, rutin, icaraside B, aviculin, (+)-catechin, (-)-epicatechin, (-)-epicatechin-3-O-gallate, (-)-epigallocatechin-3-O-gallate, (+)-galocatechin, and (-)-epigallocatechin were isolated from *Scurrula atropurpurea* (Ohashi et al., 2003).

Fatty acids, monoterpene glycoside, lignan glycoside and flavanes from *Scurrula atropurpurea* inhibited cancer cell invasion (Ohashi et al., 2003). *Scurrula atropurpurea* also modulated the total plasma nitrate/nitrite levels and reduced endothelial damage by increasing endothelial progenitor cells in DOCA-salt hypertensive rats (Athiroh et al., 2014). *Scurrula oortiana* showed significant cytotoxicity against WEHI-164 cells and increased tumour cell sensitivity to TNF α mediated lysis (Murwani, 2003). Quercetin from *Scurrula ferruginea* potentially inhibited U251 (human glioblastoma cells) (Lohézic-Le et al., 2002). Polysaccharides from *Scurrula parasitica* down-regulated the expression of Ki-67, CyclinD1 and Bcl2 proteins and up-regulated the expression of Bax protein which inhibited the cancer cell proliferation and enhanced apoptosis (Xiao et al., 2010).

2.10 *Symplocos lucida* (Thunberg) Siebold & Zuccarini

Symplocos lucida (Thunberg) (Plate 3) belongs to a unigeneric family Symplocaceae consisting of only one genus named *Symplocos* (Cronquist, 1981; Takhtajan, 1997). The genus *Symplocos* consists of about 300–500 species. These plants are distributed in tropical and subtropical regions of Asia, Australia and America (Cronquist, 1981; Hegnauer, 1973; Takhtajan, 1997). Traditionally, plants of the genus *Symplocos* have been widely used in the treatment of

various diseases. *S. racemosa* is used in Ayurvedic medicine for the treatment of diarrhea, dysentery, eye diseases, gum bleeding, menorrhagia and uterine disorders (Watt, 1972). *S. chinensis* is a potent remedy for malaria, tumefaction, enteritis, nephritis, and snake bite (Li et al., 2003). *S. cochinchinensis* is used in the treatment of leprosy, tumors, diarrhea, dysentery, menorrhagia, inflammation, and uterine disorders (CCTHD, 1992).

The phytochemical investigation of *Symplocos* species had resulted in isolation of confusoside, trilobatin, β -sitosterol, β -amyirin, symplocoside, salireposide, benzoylsalireposide, oleanolic acid and stigmasterol (Abbasi, 2004). Compounds such as (-)-pinoresinol- β -D-glucoside and (-)-pinoresinol monomethyl ether- β -D-glucoside were isolated from *Symplocos lucida* (Inouye et al., 1973). *Symplocos racemosa* afforded the isolation of flavans, symposide and its glycoside, (-)-epiafzelechin (Dhaon et al., 1989), *n*-hexacosanol, *n*-octacosanol, nonaeicosanol, *n*-hentriacontanol, methyl triacontanoate, tricontyl palmitate (Gupta & Sharma, 2015), 28-hydroxy-20 α -urs-12,18(19)-dien-3 β -yl acetate, 3-oxo-urs-20 α -12,18(19)-dien-28-oic acid, 24-hydroxyolean-12-en-3-one, betulin and oleanolic acid (Ali et al., 1990). Symplocosides A-F were isolated from *Symplocos chinensis* (Tang et al., 2004). Octacos-1-ene, stigmasterol and lupeol were isolated from *Symplocos paniculata* (Kumar & Jangwan, 2012).

Recently much attention has been paid to the genus *Symplocos* due to their diverse biological activity in treating various disorders. *Symplocos racemosa* exhibited antioxidant, and antitumor properties (Vijayabaskaran et al., 2010), antifibrinolytic activity (Dhaon et al., 1989), anti-inflammatory, and analgesic effects (Mehjabeen et al., 2014), and antibacterial activities against *Propionibacterium acnes* and *Staphylococcus epidermidis* (Devmurari, 2010; Kumar et al., 2007). *S. racemosa* exerted anticancer effects on HL 60, and HeLa cancer cell lines (Bhuvan et al., 2009). Symconoside A and symconoside B from *Symplocos racemosa* inhibited phosphodiesterase-I (Ahmad et al., 2005). Locoracemosides A, B and C from *Symplocos racemosa* inhibited α -chymotrypsin (Rashid et al., 2008). *S. chinensis* root extract and an ursane triterpenoid, 2 β ,3 β ,19 α ,24-tetrahydroxy-23-norurs-12-en-28-oic acid, from it inhibited B16 and BGC-823 cells (Li et al., 2003). Triterpenoid saponins from *Symplocos chinensis* were found cytotoxic against different cells such as symplocoside A against KB, HCT-8, A549, and HELF cells, symplocoside C against HCT-8 and BGC-823 cells, and symplocoside F against HCT-8 cells (Tang et al., 2004). Matairesinol and harman from *Symplocos setchuensis* inhibited HIV replication in H9 lymphocyte cells (Ishida et al., 2001). *S. cochinchinensis* showed antidiabetic and antilipidemic effects in streptozotocin induced diabetic rats (Sunil et al., 2012).

2.11 *Terminalia bellirica* (Gaertner) Roxburgh

Terminalia bellirica Roxb (Combretaceae) (Plate 3) is a large deciduous tree distributed in Nepal, India, Sri Lanka, Southeast Asia (Manandhar, 2002), and equally in Africa and America (Sastri, 1952). It is widely used in Ayurveda, Siddha and Chinese systems of medicine (Saraswathi et al., 2012) for the treatment of anaemia, asthma, cancer, constipation, diarrhoea, dysuria, headache, hypertension, inflammation and rheumatism (Duke et al., 2002). *T. bellirica* is effective in diarrhoea, leucoderma, impotence, bronchitis, cold, cholera, respiratory tract infections, anaemia, haemorrhoids, eye infections, as a brain tonic and as a diuretic to remove kidney stones (Chopra et al., 1956; Sharma et al., 2001; Singh et al., 2002).

Terminalia bellirica contains alkaloids, flavonoids, tannins, saponin, glycosides, terpenoids and polyphenols (Arulmozhi & Surya, 2016). *Terminalia bellirica* afforded the isolation of termilignan, thannilignan, 7-hydroxy-3',4'-(methylenedioxy) flavone, anolignan B, gallic acid, ellagic acid, β -sitosterol, arjungenin, belleric acid, bellericoside and cannogenol 3-O- β -D-galactopyranosyl-(1 \rightarrow 4)-O- α -L-rhamnopyranoside (Khan & Gilani, 2008; Nandy et al., 1989). It also contains numerous phytoconstituents such as gallo-tannic acid, bellericanin, ellagic acid, gallic acid, termilignan, flavone and anolignan B, tannins, ellagic acid, galloyl glucose and chebulaginic acid, phenyllembin, mannitol, glucose, fructose and rhamnose, flavonoids (Khan & Gilani, 2008).

Terminalia bellirica showed antidiabetic and hypolipidemic activity (Arulmozhi & Surya, 2016). It was effective against obesity, insulin resistance, and hyperlipidemia (Makihara et al., 2012). *Terminalia bellirica* suppressed absorption of triacylglycerol and inhibited pancreatic lipase activity (Makihara et al., 2012). *Terminalia bellirica* stimulates the secretion of insulin and enhance its action and inhibits starch digestion (Kasabri et al., 2010). *Terminalia bellirica* showed better glucose uptake potential in *Vero*, L-6 and 3T3 cell lines (Das & Devi, 2015). Ethnolic extract of *Terminalia bellirica* triggers both non-specific and specific cellular immunity (Manjunatha et al., 2011). Ethyl acetate and methanolic extract of *Terminalia bellirica* fruits displayed antiviral activity at very low concentrations (Nagendraswamy et al., 2013). *Terminalia bellirica* exhibits anti-asthmatic, antitussive, anti-spasmodic, anti-mutagenic and anti-HIV properties (Wadekar & Patil, 2008). *T. bellirica* exhibited antioxidant, antimicrobial, and antiinflammatory activities (Manohar et al., 2012; Upadhyay et al., 2015). *Terminalia bellirica* Roxb. fruits extracts have shown nephroprotective effect in ethylene glycol induced urolithiatic model (Upadhyay et al., 2015). *Terminalia bellirica* are therapeutically effective for the treatment of CaOx stones (Upadhyay et al., 2015). Alcoholic extract of *Terminalia bellirica* showed anti-colitic activity in dextran sulfate sodium induced

colitis (Desai et al., 2014). *Terminalia bellirica* leaf showed a strong angiogenic activity (Prabhu & Gopal, 2011; Vinoth et al., 2012). Ethanolic and aqueous extracts of *Terminalia bellirica* possessed significant analgesic and antipyretic in mice and rats (Sharma et al., 2010).

2.12 *Tsuga dumosa* (D. Don) Eichler

Tsuga dumosa D. Don (Pinaceae) (Plate 2) is an economically as well as medicinally important conifer (Fang et al., 1985) found in Nepal, India and Myanmar (Manandhar, 2002). In Nepal, it is found in the northern belt at an altitude of 2000–3600 m in association with other conifers like *Taxus*, *Abies* and *Picea* (Bhattacharyya et al., 1992). It is commonly known as “Hemlock Spruce” and locally called as “Chune Salla”, “Gobre Salla” or “Thingre Salla” (Manandhar, 2002). The plant has been extensively used for timbering and lumber products because of its resistance to decay. The bark of this plant is a rich source of tannin, hence can be used for dyeing (Zhao et al., 2005). A solid self bioadhesive herbal compound isolated from *T. dumosa* was used for the treatment of the oral mucosal disorder and for patients exhibiting herpetic stomatitis lesions, aphthous ulcers, mucosal inflammation, toothache and lesions on the lips and gingival (Domb & Wolnerman, 2002).

Phytochemical screening of *T. dumosa* showed the presence of volatile oil, fatty acids, triterpenes, sterols, carotenoids, polyphenols, reducing compounds, anthracenocoids, coumarins, flavone aglycones, tannins and saponins (Dhakal et al., 2011). *Tsuga dumosa* afforded the isolation of nonacosan-10-ol, stigmasterol, α -sitosterol, α -sitosterol glucoside (Dhakal et al., 2011), dumosaol elucidated as 3-(4-hydroxy-3-methoxy-benzyl)-5-[2-(4-hydroxy-3-methoxy-phenyl)-3-hydroxymethyl-7-methoxy-2,3-dihydrobenzofuran-5-yl]-4-hydroxymethyl-dihydro-furan-2-one (Zhao et al., 2005), 4,8-dihydroxyepipinoresinol (Zhao et al., 2004b), (+)-parryin-15,16-o-isopropylidene, (-)-isopimara-8 (14)-en-15,16-O-isopropylidene, isopimara-8-en-15,16-O-isopropylidene and isopimara-8-en-15,16-diol (Zhao et al., 2004a).

3 Materials and Methods

3.1 Collection of Plant Materials

Plants were selected based on their ethnomedicinal use in the study area. During sample collection, habitat features of plant species like GPS location, associated species, habitat type and forest type were recorded. Ethnobotanical information was documented on the basis of field based primary data collection through semi-structured and unstructured questionnaires and secondary information was collected from published and unpublished literatures. Information of plant specimens like local names, vernacular name, plant parts used, purpose of use, mode of preparation and administration for medicinal and other purposes were collected. The plant specimens were collected from the study area during the period of June – October 2014. The specimens were identified in the field with the help of the local people for the vernacular name. The voucher specimens were collected and dried which were then authenticated by Ethnobotanical society of Nepal (ESON) with the help of taxonomic literatures (Grierson & Long, 1984-1999; Polunin & Stainton, 1987; Press et al., 2000) along with consultation of taxonomists and thus prepared herbarium specimens were deposited at Tribhuvan University Central Herbarium (TUCH).

Table 3.1 Name, collection site, parts used and ethnobotanical information of 12 medicinal plants.

Scientific name	Family	Vernacular name	Collection site	GPS coordinates	Elevation (m)	Plant parts	Ethnobotanical information
<i>Loxogramme involuta</i>	Polypodiaceae		Phulkharka-3, Dhading	N 28° 04.510', E 84° 55.104'	1750	Whole plant	Cuts and wounds
<i>Hydrocotyle javanica</i>	Apiacea	Ghortapre	Baseri-5, Dhading	28° 06' 14.7", E 84° 51' 42.7"	1378	Whole plant	Fever
<i>Phyllanthus emblica</i>	Euphorbiaceae	Amala	Budhathum-9, Dhading	N 28° 04' 39.9", E 84° 50' 22.4"	850	Bark	Stomach uneasiness

<i>Chrysopogon aciculatus</i>	Poaceae	Kuro ghans	Baseri-5, Dhading	N 28° 06' 14.8", E 84° 51' 42.9"	1380	Whole plant	Wound, cut delivery problems
<i>Rhododendron lepidotum</i>	Ericaceae	Sundhup	Phulkharka-2, Dhading	N 28° 06.286', E 84° 54.436'	2890	Leaf, Stem	Rheumatic pain and Syphilis
<i>Geranium wallichianum</i>	Geraniaceae	Urmen	Phulkharka-3, Dhading	N 28° 05.456', E 84° 55.379'	2350	Root	Gastritis and female problems
<i>Tsuga dumosa</i>	Pinaceae	Gobresalla	Phulkharka-3, Dhading	N 28° 06.290', E 84° 54.512'	2870	Bark	NA
<i>Anemone rivularis</i>	Ranunculaceae	Hattipaile	Phulkharka-3, Dhading	N 28° 05.459', E 84° 55.455'	2366	Whole plant	Cough and fever
<i>Scurrula elata</i>	Loranthaceae	Liso	Phulkharka-3, Dhading	N 28° 05.292', E 84° 55.285'	2275	Branches	NA
<i>Terminalia bellerica</i>	Combretaceae	Barro	Amalachour-8, Baglung	N 28° 12.475', E 83° 38.778'	952	Fruit	Fever, Anthihelminthic
<i>Ranunculus scleratus</i>	Ranunculaceae	Mudulejhar	Amalachour-8, Baglung	N 28° 12.475', E 83° 38.778'	952	Whole plant	Blisters, Gastric
<i>Symplocos lucida</i>	Symplocaceae	Tebba	Phulkharka-4, Dhading	N 28° 06.180', E 84° 54.916'	2805	Bark	NA

3.2 Methanol Extraction of Plant Samples

Air-dried plant samples were grinded to fine powder using an electric grinder. 100g of plant powder was weighed and soaked in 750 ml of HPLC grade methanol and then subjected to intermittent sonication at 40 kHz and 45°C for 75 minutes each time for three consecutive days to complete the extraction. The resulting solution was filtered through Whatmann No.1 filter paper. The filtrate was then subjected to rotoevaporation with water bath at 45°C, and round bottom flask at 60-80 rpm for 20-45 minutes. Extract was thus concentrated to 50-60ml and transferred to clean and weighed petriplates and kept in an incubator at 37°C for about 3 days to remove excess methanol. The dried extracts in petriplates were weighed to estimate the percentage yield using following formula,

$$\text{Percentage Yield} = \frac{\text{Dry Weight of Extract}}{\text{Dry Weight of Plant Material}} \times 100$$

The dried extracts were collected in cryovials and stored at -20°C until use.

3.3 Total Phenolic Content

Total phenolic content (TPC) of methanolic extracts were determined by Folin-Ciocalteu method (Singleton & Rossi, 1965). A hundred microliter of extract (2.5mg/ml) was separately mixed with 1 ml Folin Ciocalteu phenol reagent (Merck Ltd, India) (1:10 dilution with the distilled water) and 0.8ml of aqueous 1M Na₂CO₃ solution. The reaction mixture was allowed to stand for about 15 minutes at room temperature. After completion of incubation the absorbance of the reaction mixture (Blue/violet solution) was measured at 765nm using the UV-Visible Spectrophotometer (Thermo fisher scientific, Genesystem- 10-5). For blank, 0.1 ml of absolute methanol was added instead of extract solution in above reaction mixture. A callibration curve was obtained using gallic acid (Moly chem, Mumbai, india) as standard in methanol and water mixture (50:50 v/v) with the concentration ranging from 5-350µg/ml. Total phenolic content of plant extract were then quantified by using standard calibration curve generated. The result was expressed in terms of milligram of gallic acid equivalent per gm extract (mgGAE/g). For each extract, triplicates were performed to get more accurate result.

3.4 Total Flavonoid Content

Total flavonoid content (TFC) of methanol extracts were determined using spectrophotometric assay based on aluminium-flavonoid complex formation in presence of

potassium acetate as suggested by Pękal & Pyrzynska (2014) with slight modification. Two hundred and fifty microliters of each extract (10mg/ml) was separately mixed with the 0.75 ml of methanol, 0.05ml of 10% aluminium chloride, 0.05 ml of 1M potassium acetate (CH₃COOK) and 1.4 ml of distilled water. The reaction mixtures were shaken and incubated for 30 minutes at room temperature. The absorbance of the mixture (yellow coloured complex) was measured at 415nm using the UV-Visible spectrophotometer (Thermo Fisher Scientific, Genesystem-10-5). Blank was prepared adding all the reagents except the plant sample.

Callibration curve was obtained with the help of quercetin (Sigma) as standard solution in methanol with the concentration ranging from 10-100µg/ml. The total flavonoid content was then expressed in terms of milligram of quercetin equivalent per gram extract (mgQE/g). All the tests were carried in triplicates to get more accurate result.

3.5 Antioxidant Activity

The antioxidant activity of plant extracts and standard ascorbic acid in methanol were determined based on their ability to scavenge 1, 1-diphenyl-2-picrylhyrazyl (DPPH) free radical. The method was carried out as described by Xu & Chang (2007) and Chen & Ho (1995) with slight modification. Different concentration of plant extract (4-300µg/ml) and ascorbic acid (1-25 µg/ml) were prepared in methanol on clean test tubes to which equal volume (0.5 ml) of the 0.2mM DPPH solution was added. Tubes were shaken uniformly for proper mixing and incubated in dark for half an hour. The control was prepared as above but without the plant extract or ascorbic acid. Methanol was taken as blank. Absorbance was taken at 517nm using UV-Visible spectrophotometer (Thermo Fisher Scientific, genesystem- 10-5).

Free radical scavenging activity of the plant extracts and ascorbic acid were calculated by using the given formula:

$$\% \text{ Radical Scavenging Activity (RSA)} = \frac{\text{Abs.control} - \text{Abs.sample}}{\text{Abs.control}} \times 100$$

IC₅₀ value of plant extracts and ascorbic acid was calculated by using the given formula:

$$IC_{50} = EXP \left(LN(\text{conc.} > 50\%) - \left(\frac{pi > 50\% - 50}{pi > 50\% - pi < 50\%} \times LN \left(\frac{\text{conc.} > 50\%}{\text{conc.} < 50\%} \right) \right) \right)$$

EXP: Exponential function; LN: Natural log function, both used in Microsoft Excel 2016; pi>50%: RSA value just above 50%; pi<50%: RSA value just below 50% conc.>50%: Concentration of Signal>50%; conc.<50%: Concentration of Signal <50%.

3.6 GC-MS Analysis

GC-MS analyses of methanol extracts of 12 medicinal plants were performed in Nepal Forensic Science Laboratory, Khumaltar, Lalitpur and Advanced Instrumentation Research Facility, JNU, New Delhi using GCMS-QP2010 Plus and GCMS-QP2010 Ultra Shimadzu systems each comprising an automatic operation controller (AOC-20i, 20s) and Gas chromatograph interfaced to a mass spectrometer (GC-MS) equipped with a Rtx-5MS (5% diphenyl/95% dimethyl polysiloxane) fused silica column (30 m × 0.25 mm ID × 0.25 μm df). For GCMS detection, an electron ionization system was operated in electron impact mode with an ionization energy of 70 eV. Helium gas (99.999%) was used as a carrier gas at a constant flow rate of 1.35 ml/min, and an injection volume of 1 μl was employed (a split ratio of 2:1). The injection temperature was maintained at 280 °C, the ion-source temperature was 200 °C, the oven temperature was programmed from 50 °C (isothermal for 0 min), with an increase of 10 °C/min to 250°C (isothermal for 2 min), and 15 °C/min to 280°C (isothermal for 16 min). Mass spectra were taken at 70 eV; a scan interval of 0.5 s and fragments from 45 to 450 Da. The solvent delay was 0 to 2 min, and the total GC/MS running time was 60 min. The relative percentage amount of each component was calculated by comparing its average peak area to the total areas.

Interpretation on mass spectrum GCMS was conducted using the software database of NIST08, NIST08s, NIST11 and WILEY8 libraries. The spectrum of the unknown components was compared with the spectrum of known components stored in the library. The name, molecular weight, and structure of the components of the plant extracts were ascertained.

3.7 Antimicrobial activity

3.7.1 Nutrient agar (NA)

NA plates were prepared for bacterial culture in antibacterial tests. About 28 gram of NA powder (Hi Media Laboratories Pvt. Ltd, Mumbai, India) was carefully weighed and poured in distilled water. The contents were dissolved on the water completely and the final volume was maintained to 1000 ml followed by boiling for uniform mixing. This media was sterilized on an autoclave at 15lbs pressure at 121°C for 15 minutes. The autoclave tape was used as an indicator for the completeness of sterilization. After cooled to about 45-50°C, the media was poured on sterilized and properly labeled petridishes. About 25 ml of the media was poured on each petridish of 9 cm diameter in sterile conditions under a laminar flow hood. All of the plates were left for the solidification of media. For the preparation of the NB broth, screw tight bottles were filled with NB media and autoclaved. It was then cooled and used.

3.7.2 Mueller Hinton Agar (MHA)

The Mueller Hinton Agar (MHA) media was required for antimicrobial tests. About 38 gram of MHA powder (Hi Media Laboratories Pvt. Ltd, Mumbai, India) was weighed and transferred on a conical flask. The content was dissolved in distilled water and final volume was maintained to 1000 ml. The media was sterilized by autoclaving at 15 lbs pressure and 121°C for 15 minutes. Autoclave tape was used for the indication of the completeness of the sterilization. The media was cooled in laminar airflow and dispensed on sterile and dry petridishes.

3.7.3 Standard bacterial culture Inoculums

The individual pure culture of *Staphylococcus aureus* (ATCC 25923), methicillin-resistant *Staphylococcus aureus* (MRSA), *Klebsiella pneumoniae* (MDR), and *Salmonella* Typhi (MDR) were obtained from Tribhuvan University Teaching Hospital, Kathmandu and were streaked on NB broth with the help of the sterilized inoculating loop. The inoculated culture bottles were kept on the shaking incubator at 37°C and 120 rpm for overnight. The turbidity of the sub-cultured bacterial suspension was adjusted at the 0.5 McFarland standards freshly prepared on the other day for antibacterial tests. These bacterial inoculums were used for the swabbing on the MHA plates to test the antimicrobial effects of the plant extracts.

3.7.4 Antibacterial test

The antimicrobial activities of medicinal plant extracts were tested by using agar well diffusion method. The sterile and dry MHA plates were properly labeled with name of bacteria, name of the plant species and the concentration of the plant extract to be added. The plates were inoculated with the appropriate bacterial culture by a sterile cotton swab aseptically. These plates were then allowed to dry for about 15 to 20 minutes.

Five wells were prepared on the solid MHA media with the help of the sterile cork borer (4 mm diameter). Four different concentrations (100 mg/ml, 50 mg /ml, 25 mg /ml, and 12.5 mg /ml) of the plant sample were prepared on DMSO. With the help of the sterile pipette the 30 µl of each individual plant extract were poured in the above prepared wells. The DMSO was taken as negative control while the antibiotic disc – Piperacillin (PI100), Methicillin (MET5), or Piperacillin/Tazobactam (PIT100/10) – was taken as the positive control. The plates were incubated on the microbial incubator overnight at 37°C and the zone of inhibition was observed against each type of bacteria at different concentrations of each plant extract.

3.7.5 Potato Dextrose Agar (PDA) and Potato Dextrose Broth

About 25 gram of PDA powder (Hi Media Laboratories Pvt. Ltd, Mumbai, India) was carefully weighed, dissolved in distilled water and final volume was maintained to 1000 ml. This media was autoclaved at 15 lbs pressure and 121°C for 15 minutes, was cooled in laminar airflow and dispensed in sterile petriplates. Likewise, PD broth was prepared in culture tubes excluding agar.

3.7.6 Preparation of the standard fungal cultures

The individual pure and characterized cultured of *Saccharomyces cerevesiae*, and *Pichia sp.* was obtained from CDBT, TU and were sub-cultured in PD broth with the help of the sterilized inoculating loop and kept on the shaking incubator at 37°C and 120 rpm for overnight. The turbidity of the sub-cultured fungal suspension was adjusted to 0.5 McFarland standards. These bacterial inoculums were used for the swabbing on the MHA plates to test the antimicrobial effects of the plant extracts.

3.7.7 Antifungal test

The antifungal activities of medicinal plant extracts were tested by using agar well diffusion method. The sterile and dry PDA plates were properly labeled with name of fungi, name of the plant species and the concentration of the plant extract to be added. The plates were inoculated with the appropriate fungal culture by a sterile cotton swab aseptically. These plates were then allowed to dry for about 15 to 20 minutes.

Six wells were prepared in solid agar in PDA plates with the help of the sterile cork borer (4 mm diameter). Four different concentrations (100 mg/ml, 50 mg /ml, 25 mg /ml, and 12.5 mg/ml) of the plant extracts were prepared on DMSO. With the help of the sterile pipette, 30 µl of different concentrations of each plant extract were dispensed in their respective wells. The DMSO was taken as negative control. Fluconazole (2.5 mg/ml) was taken as the positive control. The plates were incubated on the microbial incubator overnight at 28°C and the zone of inhibition was observed against different type of fungi at different concentrations of each plant extract.

3.8 Brine Shrimp Cytotoxicity

Brine shrimp lethality bioassay was carried out as previously described by Solis et al. (1993) with some modification. Artificial sea water was prepared by dissolving 38 gm of pure NaCl in one litre of distilled water supplemented with 6 mg of dried yeast. *Artemia salina* (brine shrimp) eggs, obtained from Central Department of Chemistry, were left to hatch in artificial

sea water for 48 hours at room temperature (25-29°C). Brine shrimp larvae (nauplii) were attracted to one side of the vessel exposing them to a light source and separated from eggs by pipetting them 2-3 times in small beakers containing saline water. Plant extracts were dissolved in artificial sea water and made up to 2 mg/ml. Water insoluble compounds were dissolved in 50 µl DMSO prior to adding sea water. Plant extracts were pipetted into the wells of 96-well microplates (Nunc, Denmark) and serial dilutions were made so that the final concentration of each extract after adding 100 µl of suspension containing 10-12 nauplii to form the final volume of 200 µl would be 1000 µg/ml, 100 µg/ml, and 10 µg/ml in triplicate. Control wells with Potassium dichromate, DMSO and artificial sea water were included in each experiment. The plates were covered and incubated at room temperature (25-29°C) for 24 hours. Plates were then examined under the binocular stereomicroscope and the numbers of dead (non-motile) nauplii in each well were counted. One hundred microliter of methanol were added to each well to immobilize the nauplii and after 15 minutes the total numbers of brine shrimp in each well were counted. Percentage mortality was calculated as given by the formula:

$$\%Mortality = \frac{Observed\ Proportion - Proportion\ at\ Zero\ Concentration}{1 - Proportion\ at\ Zero\ Concentration} * 100$$

Here the proportion is meant for the number of dead to the total number of brine shrimps.

Probit analysis was performed to determine the lethal concentration to half of the test organisms (LC_{50}) on Excel 2016 as described by Currell (2015). Plant extracts were classified for their toxicity based on the benchmark created by Clarkson according to which extracts with $LC_{50} > 1000$ µg/ml are non-toxic, 500 - 1000 µg/ml are low toxic, 100 - 500 µg/ml are medium toxic, and < 100 µg/ml are highly toxic (Clarkson et al., 2004).

3.9 MTT Cell Proliferation

Cytotoxicity of plant extracts on Hela (human cervical adenocarcinoma) and MDCK (normal canine kidney) cell lines obtained from NCCS, Pune was determined by a rapid colorimetric assay, using 3-(4,5-dimethyl thiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT), which is based on the metabolic reduction of soluble MTT by mitochondrial enzyme activity of viable cells into an insoluble colored formazan product. Five hundred milliliter of Eagle's Minimum Essential Medium (EMEM) was supplemented with 10% of fetal calf serum, 5 ml of penicillin/streptomycin (50 IU/ml and 50 µg/ml respectively), 5 ml of sodium pyruvate (1 mM), $NaHCO_3$ (1 g) and 5 ml of L-glutamine (2 mM). The final medium was then sterilized using 0.22µ microfilters and stored at 4°C before use. Methanolic extracts of twelve medicinal

plants were dissolved in DMSO and further diluted with cell culture medium so that the final DMSO concentration was $\leq 0.3\%$ of total volume of the medium in all treatments. The monolayer cell culture in T175 flask was trypsinized and the cell count was adjusted to 1×10^4 cells per ml using medium containing 10% FBS. 100 μ l of cell suspension (1×10^3 cells per well) were seeded into 96-well microplate and incubated for 24 hours at 37°C in 5% CO₂ incubator. The medium was flicked off and 100 μ l of each plant extract was added to the respective well. Cells with no treatment and 5-fluorouracil treatment were examined as negative and positive controls, respectively. Microplates containing cells and extracts were incubated for another 24 hours. The cells were checked for granularity, shrinkage, swelling. 20 μ L of MTT reagent (Merck, Germany) (5mg/ml in PBS) was added to each well and incubated for 3 hours. The medium was removed and 100 μ l of DMSO (culture grade) was added to each well and gently shaken for 15 minutes to dissolve any formazan crystals formed. The formazan salts were then quantified by measuring the absorbance at 595nm and 655nm on a microplate reader. Cell viability was calculated as a percentage of untreated cells.

$$\% \text{ Cell Survival} = \frac{At - Ab}{Ac - Ab} \times 100$$

At: Absorbance of tested concentration, Ab: Absorbance of blank, Ac: Absorbance of negative control

The cytotoxicity of each plant extract was presented as EC₅₀ using GraphPad Prism 6 Software.

3.10 Statistical Analysis

The data were expressed as mean \pm standard deviation of three independent experiments. The data were analyzed using one-way analysis of variance (ANOVA). Differences were considered statistically significant at $P < 0.05$. Statistical analysis was performed using the Microsoft Excel 2016 and the GraphPad Prism 6.

4 Results

4.1 Methanol Extraction of Plant Materials

Among the twelve medicinal plants selected, the extract yield in methanol varied from 4.39% to 32.48%. The highest yield was obtained from *Terminalia bellerica* and lowest yield was found in *Chrysopogan aciculatus*. Extracts varied in color from yellowish green to reddish brown which were either greasy (sticky) or powdery in consistency. The percentage yield and characteristics of the extracts are shown in Table 4.1.

Table 4.1 Percentage yield and physical characteristics of the crude methanol extracts.

Plant Sample	Dry Weight (gm)	Weight of extract (gm)	Percentage Yield	Characteristics of Extract	
				Color	Consistency
<i>Loxogramme involuta</i>	100	10.3	10.3	Yellowish green	Sticky
<i>Hydrocotyle javanica</i>	100	9.85	9.85	Yellowish green	Sticky
<i>Phyllanthus emblica</i>	100	20.76	20.76	Greenish brown	Powdery
<i>Chrysopogan aciculatus</i>	100	4.39	4.39	Yellowish green	Sticky
<i>Rhododendron lepidotum</i>	100	7.12	7.12	Yellowish green	Sticky
<i>Geranium wallichianum</i>	100	19.34	19.34	Reddish brown	Powdery
<i>Tsuga dumosa</i>	100	9.18	9.18	Reddish brown	Powdery
<i>Anemone rivularis</i>	100	5.75	5.75	Greenish brown	Sticky
<i>Scurrula elata</i>	100	10.19	10.19	Greenish brown	Powdery
<i>Terminalia bellerica</i>	100	32.48	32.48	Greenish yellow	Powdery
<i>Ranunculus scleratus</i>	100	7.41	7.41	Brownish green	Sticky
<i>Symplocos lucida</i>	100	30.17	30.17	Yellowish red	Sticky

4.2 Determination of Total Phenolic Content

Standard curve of gallic acid offered an equation of $y = 0.0047x + 0.1819$, $R^2 = 0.9884$ (Figure 4.1). It helped determine the total phenolic content of methanolic extracts of each plant whose values were located at or somewhere within the range in between 6.74 mgGAE/g and 55.66 mgGAE/g (Figure 4.2). Highest amount of total phenolic content was found in *P. emblica* (55.66 ± 0.84 mgGAE/g) followed by *G. wallichianum* (54.99 ± 0.74 mgGAE/g), *S. elata* (54.71 ± 0.68 mgGAE/g), *T. bellerica* (54.69 ± 0.77 mgGAE/g), and *T. dumosa* (54.19 ± 0.93 mgGAE/g). The lowest amount of total phenolic content was found in *R. scleratus* (6.74 ± 0.62 mgGAE/g) (Figure 4.2).




Figure 4.1 Standard graph of gallic acid.




Figure 4.2 Total phenolic content in methanolic extracts of 12 medicinal plants.

4.3 Determination of Total Flavonoid Content

Standard curve of quercetin presented itself with an equation of $y = 0.0198x - 0.1337$, $R^2 = 0.9740$ (Figure 4.3). It helped determine the total flavonoid content of each extract whose value was positioned at or somewhere within the narrow range between 7.51 mgQE/g and 25.65 mgQE/g (Figure 4.4). The highest amount of total flavonoid content was found in *P. emblica* (25.65 ± 0.77 mgQE/g) followed by *S. elata* (22.26 ± 0.69 mgQE/g), *C. aciculatus* (19.83 ± 0.94 mgQE/g), *G. wallichianum* (18.53 ± 0.67 mgQE/g), *R. lepidotum* (18.35 ± 0.76 mgQE/g), and *H. javanica* (18.24 ± 0.83 mgQE/g). The lowest amount of total flavonoid content was found in *T. bellerica* (7.51 ± 0.54 mgQE/g) (Figure 4.4).




Figure 4.3 Standard graph of quercetin.




Figure 4.4 Total flavonoid content in methanolic extracts of 12 medicinal plants.

4.4 Estimation of Antioxidant Activity

Plant extracts exerted a dose dependent DPPH free radical scavenging activity with their IC_{50} values ranging from 4.54 $\mu\text{g/ml}$ to 195.55 $\mu\text{g/ml}$ (Figure 4.6 and Figure 4.7). *T. dumosa* ($4.54 \pm$

0.18 $\mu\text{g/ml}$) showed the highest antioxidant activity with the lowest IC_{50} value – lower than that observed in ascorbic acid ($4.57 \pm 00 \mu\text{g/ml}$) – followed by *P. emblica* ($6.64 \pm 0.71 \mu\text{g/ml}$), *G. wallichianum* ($7.24 \pm 0.48 \mu\text{g/ml}$), *T. bellerica* ($12.08 \pm 0.76 \mu\text{g/ml}$), and *S. elata* ($14.20 \pm 0.94 \mu\text{g/ml}$). *R. scleratus* ($195.55 \pm 2.45 \mu\text{g/ml}$) showed the highest IC_{50} value with the lowest antioxidant activity (Figure 4.7).




Figure 4.5 DPPH free radical scavenging activity of ascorbic acid.




Figure 4.6 DPPH free radical scavenging activity of methanolic extracts of 12 medicinal plants.




Figure 4.7 IC₅₀ of DPPH free RSA of methanolic extracts of 12 medicinal plants.

4.5 Correlation between Total Phenolic Content and Antioxidant Activity

A significant correlation was observed between the antioxidant activity and the total phenolic content while comparing the TPC with the IC₅₀ values of DPPH free radical scavenging activity of twelve medicinal plants ($r = -0.9684$, $p < 0.0001$).




Figure 4.8 Correlation between total phenolic content and IC₅₀ of DPPH free RSA of methanolic extracts of 12 medicinal plants.

4.6 Correlation between Total Flavonoid Content and Antioxidant activity

No significant correlation could be established between antioxidant activity and total flavonoid content while comparing the TFC with the IC₅₀ values of DPPH free radical scavenging activity of twelve medicinal plants ($r = -0.09653$, $p = 0.7654$).




Figure 4.9 Correlation between total flavonoid content and IC₅₀ of DPPH free RSA of methanolic extracts of 12 medicinal plants.

4.7 GC-MS Analysis

GC-MS analyses presented an excellent separation of phyto-compounds as shown by the chromatograms in Figure 4.10 to Figure 4.21. It permitted the inspection of each peak for their identification by comparison of the observed spectra of each constituent with that from NIST and WILEY software database libraries. The active principals with their retention time, molecular formula, molecular weight, and concentration percentage in the methanol extract are presented in

Table 4.2 to Table 4.13.

4.7.1 GC-MS Analysis of *Loxogramme involuta*

Loxogramme involuta afforded the identification of compounds such as 2,4-Bis[4-chloro-*trans*-styryl]-6-[(3-pyrrolidinomethyl-4-hydroxyphenyl)amino]pyrimidine (20.19%), 2-Furancarboxaldehyde, 5-(hydroxymethyl)- (9%), Cyclopentane, 1-acetyl-1,2-epoxy- (7.13%), Benzaldehyde, 2-hydroxy-6-methyl- (4%), 1,2-Benzenediol (3.87%), 9-Octadecynoic acid (3.81%), Octadecane, 2,2,4,15,17,17-hexamethyl-7,12-bis(3,5,5-trimethylhexyl)- (3.75%), Crypto dioxmine diacetate (3.66%), 2-Deoxy-D-galactose (3.41%), and 2-(Heptyl-methyl-carbamoyl)-hexanedioic acid, bis-(heptyl-methyl-amide) (2.96%).




Figure 4.10 GC-MS Chromatogram of methanolic extract of *Loxogramme involuta* whole plant.

Table 4.2 Phytocomponents identified in the methanolic extract of *Loxogramme involuta* whole plant by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	2.533	2.13	Silane, trimethyl[5-methyl-2-(1-methylethyl)phenoxy]-		
2	2.6	2.96	2-(Heptyl-methyl-carbamoyl)-hexanedioic acid, bis-(heptyl-methyl-amide)	C ₃₁ H ₆₁ N ₃ O ₃	523
3	2.899	3.66	Crypto dioxmine diacetate	C ₃₁ H ₄₈ N ₂ O ₆	544
4	4.139	20.19	2,4-Bis[4-chloro- <i>trans</i> -styryl]-6-[(3-pyrrolidinomethyl-4-hydroxyphenyl)amino]pyrimidine	C ₃₁ H ₂₈ Cl ₂ N ₄ O	542
5	4.531	1.32	7-Chloro-1,3,4,10-tetrahydro-10-hydroxy-1-[[3-[1-piperidinyl]propyl]imino]-3-[3,4,5(trimethoxy)phenyl]-9(2H)-acridinone	C ₃₀ H ₃₆ ClN ₃ O ₅	553
6	5.517	3.75	Octadecane, 2,2,4,15,17,17-hexamethyl-7,12-bis(3,5,5-trimethylhexyl)-	C ₄₂ H ₈₆	590
7	6.256	2.27	Tetratriacontyl trifluoroacetate	C ₃₆ H ₆₉ F ₃ O ₂	590
8	6.499	0.71	Decanoic acid, 2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9-hexadecafluorononyl ester	C ₁₉ H ₂₂ F ₁₆ O ₂	586
9	6.71	0.25	2,4,7-Trinitro-9-fluorenone dinitrophenylhydrazone	2,4-C ₁₉ H ₉ N ₇ O ₁₀	495
10	6.852	7.13	Cyclopentane, 1-acetyl-1,2-epoxy-	C ₇ H ₁₀ O ₂	126
11	7.04	2.86	Maytansine	C ₃₄ H ₄₆ ClN ₃ O ₁₀	691
12	7.255	0.37	N-(4,6-Dipiperidino-1,3,5-triazin-2-yl)-N-{4,6-bis[2,2,2-trifluoro-1,1-bis(trifluoromethyl)ethyl]-1,3,5-triazin-2-yl}amine	C ₂₄ H ₂₁ F ₁₈ N ₉	777
13	7.517	2	2,6,10,14,18,22-Tetracosahexaene, 2,6,10,15,19,23-hexamethyl-, (all-E)-		
14	7.636	0.63	2-[4-Chloro- <i>trans</i> -styryl]-6-chloro-5-[4-chlorophenyl]-4-[3,5-bis[pyrrolidinomethyl]-4-hydroxyanilino]pyrimidine	C ₃₄ H ₃₄ Cl ₃ N ₅ O	633
15	7.759	1	α -d-Glucofuranose, 3-O-(2,3,4-tri-O-acetyl- α -d-ribofuranosyl)-1,2-5,6-di-O-isopropylidene-	C ₂₃ H ₃₄ O ₁₃	518

16	7.863	1.59	Acetamide, N-phenyl-		
17	8.559	3.87	1,2-Benzenediol	C ₆ H ₆ O ₂	110
18	8.795	1.07	Rhodium, bis[5,6-bis(.eta.2-ethenyl)cyclooctene]di-.mu.-chlorodi-, stereoisomer	C ₂₄ H ₃₆ Cl ₂ Rh ₂	600
19	8.974	9	2-Furancarboxaldehyde, 5-(hydroxymethyl)-	C ₆ H ₆ O ₃	126
20	9.947	0.35	Card-20(22)-enolide, 1,5,11,14,19-pentahydroxy-3-[(3,4,5-trihydroxy-6-methyltetrahydropyran-2-yl)oxy]-	C ₂₉ H ₄₄ O ₁₂	584
21	11.874	4	Benzaldehyde, 2-hydroxy-6-methyl-	C ₈ H ₈ O ₂	136
22	12.964	0.33	Pseudoatomatidin-5,20-dien diacetate	C ₃₁ H ₄₇ NO ₄	497
23	13.222	0.35	Pseudosolasodine diacetate	C ₃₁ H ₄₉ NO ₄	499
24	13.731	1.11	Diethyl Phthalate	C ₁₂ H ₁₄ O ₄	222
25	14.16	3.41	2-Deoxy-D-galactose	C ₆ H ₁₂ O ₅	164
26	16.3	2.68	Phytol	C ₂₀ H ₄₀ O	296
27	16.559	0.49	Triacotane, 1,30-dibromo-	C ₃₀ H ₆₀ Br ₂	578
28	16.747	0.93	6-Octadecenoic acid, methyl ester, (Z)-		
29	17.561	2.78	<i>n</i> -Hexadecanoic acid	C ₁₆ H ₃₂ O ₂	256
30	18.848	0.34	9,12-Octadecadienoic acid, methyl ester, (E,E)-	C ₁₉ H ₃₄ O ₂	294
31	19.016	0.92	1-Hexadecene		
32	19.234	3.81	9-Octadecynoic acid	C ₁₈ H ₃₂ O ₂	280
33	19.3	1.22	Acetamide, N-(2-methoxyphenyl)-		
34	19.44	0.82	Beryllium, hexakis[.mu.-(2,2-dimethylpropanoato-O:O')]-.mu.4-oxotetra-	C ₃₀ H ₅₄ Be ₄ O ₁₃	658
35	23.017	1.66	Hexadecanoic acid, 2,3-dihydroxypropyl ester	C ₁₉ H ₃₈ O ₄	330
36	24.659	2.17	9,12-Octadecadienoic acid (Z,Z)-, 2-hydroxy-1-(hydroxymethyl)ethyl ester	C ₂₁ H ₃₈ O ₄	354
37	24.745	0.71	4-Hexadecen-6-yne, (E)-	C ₁₆ H ₂₈	220

4.7.2 GC-MS Analysis of *Hydrocotyle javanica*

Hydrocotyle javanica afforded the identification of compounds such as 9,12-Octadecadienoic acid (Z,Z)- (29.21%), *n*-Hexadecanoic acid (15.53%), Acetic acid, fluoro-, ethyl ester (7.65%), Ethane, 1,1,1-trimethoxy- (5.03%), 3,7,11,15-Tetramethyl-2-hexadecen-1-ol (3.96%), Phytol (2.75%), 2-[[4-Chlorophenyl]thio]-4-[[7-chloro-4-quinolinyl]amino]-6-[[diethylamino]methyl]phenol (2.31%), 9,12-Octadecadienoic acid (Z,Z)-, 2-hydroxy-1-

(hydroxymethyl)ethyl ester (2.26%), Diacetamate (2.04%), and 2(3H)-Furanone, 3,4-bis(1,3-benzodioxol-5-ylmethyl)dihydro-, (3R-*trans*)- (2%).




Figure 4.11 GC-MS Chromatogram of methanolic extract of *Hydrocotyle javanica* whole plant.

Table 4.3 Phytochemicals identified in the methanolic extract of *Hydrocotyle javanica* whole plant by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	2.566	5.03	Ethane, 1,1,1-trimethoxy-	C ₅ H ₁₂ O ₃	120
2	2.705	2.31	2-[[4-Chlorophenyl]thio]-4-[[7-chloro-4-quinolinyl]amino]-6-[[diethylamino]methyl]phenol	C ₂₆ H ₂₅ Cl ₂ N ₃ OS	497
3	2.852	7.65	Acetic acid, fluoro-, ethyl ester	C ₄ H ₇ FO ₂	106
4	2.977	0.53	Tetratriacontyl trifluoroacetate	C ₃₆ H ₆₉ F ₃ O ₂	590
5	3.04	0.64	Glycerin	C ₃ H ₈ O ₃	92
6	3.178	2.04	Diacetamate		0
7	4.119	0.22	2-Cyclopentene-1,4-dione	C ₅ H ₄ O ₂	96
8	4.657	0.33	Cyclohexanone	C ₆ H ₁₀ O	98
9	5.943	0.35	4-Heptanol, 4-methyl-	C ₈ H ₁₈ O	130
10	7.847	0.49	4H-Pyran-4-one, 2,3-dihydro-3,5-dihydroxy-6-methyl-	C ₆ H ₈ O ₄	144
11	10.193	0.33	2-Methoxy-4-vinylphenol	C ₉ H ₁₀ O ₂	150

12	13.67	0.22	1H-Cycloprop[e]azulen-7-ol, decahydro- 1,1,7-trimethyl-4-methylene-, [1ar- (1 α ,4 α ,7 β ,7 α)]-	C ₁₅ H ₂₄ O	220
13	14.555	0.24	Bicyclo[4.3.0]nonan-1-ol, 7,9- bis(methylene)-2,2,6-trimethyl-	C ₁₄ H ₂₂ O	206
14	15.517	0.16	Octadecanoic acid	C ₁₈ H ₃₆ O ₂	284
15	15.893	0.28	7-Oxabicyclo[4.1.0]heptane, 1-methyl-4-(2- methyloxiranyl)-	C ₁₀ H ₁₆ O ₂	168
16	16.175	0.2	8-Octadecynoic acid, methyl ester	C ₁₉ H ₃₄ O ₂	294
17	16.315	3.96	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C ₂₀ H ₄₀ O	296
18	16.389	0.27	2-Undecanone, 6,10-dimethyl-	C ₁₃ H ₂₆ O	198
19	16.561	0.3	Oleic Acid	C ₁₈ H ₃₄ O ₂	282
20	17.045	0.18	Z,Z,Z-1,4,6,9-Nonadecatetraene	C ₁₉ H ₃₂	260
21	17.195	0.72	Hexadecanoic acid, methyl ester	C ₁₇ H ₃₄ O ₂	270
22	17.43	1.13	9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)-	C ₁₉ H ₃₂ O ₂	292
23	17.734	15.53	<i>n</i> -Hexadecanoic acid	C ₁₆ H ₃₂ O ₂	256
24	17.937	0.18	Furan, 2-[(2-ethoxy-3,4-dimethyl-2- cyclohexen-1-ylidene)methyl]-	C ₁₅ H ₂₀ O ₂	232
25	18.337	0.24	Falcarinol	C ₁₇ H ₂₄ O	244
26	18.574	0.36	2-Aziridinone, 1-(1-adamantyl)-3-(1- methylcyclopentyl)-	C ₁₈ H ₂₇ NO	273
27	18.861	1.35	9,12-Octadecadienoic acid, methyl ester	C ₁₉ H ₃₄ O ₂	294
28	19.037	2.75	Phytol	C ₂₀ H ₄₀ O	296
29	19.112	0.16	Octadecanoic acid, methyl ester	C ₁₉ H ₃₈ O ₂	298
30	19.404	29.21	9,12-Octadecadienoic acid (Z,Z)-	C ₁₈ H ₃₂ O ₂	280
31	19.703	1.4	Cyclopropane, 1-methoxy-2,2-dimethyl-3- (3,3-dimethyl-1-propynyl)-	C ₁₂ H ₂₀ O	180
32	19.783	0.77	3,9-Dodecadiyne	C ₁₂ H ₁₈	162
33	20.023	1.32	Methyl <i>trans</i> -2-octadecenoate	C ₁₉ H ₃₆ O ₂	296
34	20.396	1.23	Octadecane, 1-(ethenyloxy)-	C ₂₀ H ₄₀ O	296
35	20.573	0.93	Cyclopentolate	C ₁₇ H ₂₅ NO ₃	291
36	20.703	0.72	Acetamide, N-(2-methoxyphenyl)-		0
37	20.971	0.98	<i>cis</i> -9-Hexadecenal	C ₁₆ H ₃₀ O	238
38	21.293	1.06	Benzene, [(butylsulfonyl)ethynyl]-	C ₁₂ H ₁₄ O ₂ S	222
39	21.383	0.43	Eicosanoic acid	C ₂₀ H ₄₀ O ₂	312
40	21.637	0.72	6-Phenylhexanoic acid	C ₁₂ H ₁₆ O ₂	192
41	22.882	0.27	<i>E,E</i> -2,13-Octadecadien-1-ol	C ₁₈ H ₃₄ O	266

42	23.045	1.49	Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester	C ₁₉ H ₃₈ O ₄	330
43	23.974	0.24	6-Monoacetylmorphine	C ₁₉ H ₂₁ NO ₄	327
44	24.683	2.26	9,12-Octadecadienoic acid (Z,Z)-, 2-hydroxy-1-(hydroxymethyl)ethyl ester	C ₂₁ H ₃₈ O ₄	354
45	24.93	0.22	Tetracosanoic acid, methyl ester	C ₂₅ H ₅₀ O ₂	382
46	25.663	0.24	4-Pentenoic acid, 2,2-diethyl-3-oxo-5-phenyl-, ethyl ester	C ₁₇ H ₂₂ O ₃	274
47	25.992	1.77	2,6,10,14,18,22-Tetracosahexaene, 2,6,10,15,19,23-hexamethyl-, (all-E)-	C ₃₀ H ₅₀	410
48	26.378	0.25	1-Pyrrolidinebutanoic acid, 2-[(1,1-dimethylethoxy)carbonyl]- α -nitro-, 2,6-bis(1,1-dimethylethyl)-4-methoxyphenyl ester, [S-(R*,R*)]-	C ₂₈ H ₄₄ N ₂ O ₇	520
49	26.637	0.26	1-Triacontanol	C ₃₀ H ₆₂ O	438
50	26.907	1	1,3-Octanediol	C ₈ H ₁₈ O ₂	146
51	27.273	0.41	Oxirane, 2,2-dimethyl-3-(3,7,12,16,20-pentamethyl-3,7,11,15,19-heneicosapentaenyl)-, (all-E)-	C ₃₀ H ₅₀ O	426
52	28.136	0.42	β -Eudesmol, trimethylsilyl ether	C ₁₈ H ₃₄ OSi	294
53	28.343	0.19	Hexadeca-2,6,10,14-tetraen-1-ol, 3,7,11,16-tetramethyl-, (E,E,E)-	C ₂₀ H ₃₄ O	290
54	29.463	0.29	Cyclononasiloxane, octadecamethyl-	C ₁₈ H ₅₄ O ₉ Si ₉	666
55	29.784	2	2(3H)-Furanone, 3,4-bis(1,3-benzodioxol-5-ylmethyl)dihydro-, (3R-trans)-	C ₂₀ H ₁₈ O ₆	354

4.7.3 GC-MS Analysis of *Phyllanthus emblica*

Phyllanthus emblica afforded the identification of compounds such as 1,2,3-Benzenetriol (57.39%), Phosphoric acid, bis(trimethylsilyl)monomethyl ester (6.79%), β -D-Glucopyranose, 1,6-anhydro- (5.49%), 2-Butenal, 2-methyl-, (E)- (5.47%), 4-Dimethylsilyloxytridecane (2.59%), 5H-1,4-Dioxepin, 2,3-dihydro-2,5-dimethyl- (2.47%), 2H-Pyran-2,6(3H)-dione (2.21%), Levoglucosenone (1.95%), Benzene-1,3-diamine, 5-[3,4,4,4-tetrafluoro-1,3-di(trifluoromethyl)-2-heptafluoroisopropyl-1-butenyloxy- (1.82%), and β -Allyloxypropionic acid (0.97%).




Figure 4.12 GC-MS Chromatogram of methanolic extract of *Phyllanthus emblica* bark.

Table 4.4 Phytocomponents identified in the methanolic extract of *Phyllanthus emblica* bark by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	2.553	5.47	2-Butenal, 2-methyl-, (E)-	C ₅ H ₈ O	84
2	2.674	1.82	Benzene-1,3-diamine, 5-[3,4,4,4-tetrafluoro-1,3-di(trifluoromethyl)-2-heptafluoroisopropyl-1-butenyloxy-	C ₁₅ H ₇ F ₁₇ N ₂ O	554
3	2.898	0.5	N,N-Dicyclohexyl-2-[2-(2-{2-[(dicyclohexylcarbamoyl)-methoxy]-ethoxy}-ethoxy)-ethoxy]-acetamide	C ₃₄ H ₆₀ N ₂ O ₆	592
4	3.407	0.22	L-Phenylalanine, N-[N-[N2,N5-bis(trifluoroacetyl)-L-ornithyl]-O-(trifluoroacetyl)-D-allothreonyl]-, methylester	C ₂₅ H ₂₇ F ₉ N ₄ O ₈	682
5	4.54	0.3	9-Amidinohydrazono-3,6-dichloro-2,7-bis-[2-piperidinoethoxy]fluorine	C ₂₈ H ₃₆ Cl ₂ N ₆ O ₂	558
6	5.099	0.53	Diacetamate		0
7	5.269	0.16	Hexadecanoic acid, methyl ester		0
8	5.522	2.21	2H-Pyran-2,6(3H)-dione	C ₅ H ₄ O ₃	112
9	6.207	2.47	5H-1,4-Dioxepin, 2,3-dihydro-2,5-dimethyl-	C ₇ H ₁₂ O ₂	128
10	6.626	0.97	β-Allyloxypropionic acid	C ₆ H ₁₀ O ₃	130

11	6.766	0.17	2,4-Bis[4-chloro- <i>trans</i> -styryl]-6-[(3-pyrrolidinomethyl-4-hydroxyphenyl)amino]pyrimidine	C ₃₁ H ₂₈ Cl ₂ N ₄ O	542
12	6.878	0.58	3-Carboxy-6- <i>n</i> -butyl-7-octadecylmercapto-4-quinolone	C ₃₄ H ₅₅ NO ₃ S	557
13	7.118	0.93	1-Hexadecene		0
14	7.304	1.95	Levoglucosenone	C ₆ H ₆ O ₃	126
15	7.751	0.29	Cholestano[3,2- <i>c</i>]isoquinolin-1'(2'H)-one, 3',4'-dihydro-6',7'-dimethoxy-	C ₃₆ H ₅₅ NO ₃	549
16	8.588	0.16	Cyclohexane, (1-hexadecylheptadecyl)-	C ₃₉ H ₇₈	546
17	8.801	0.67	Benzofuran, 2,3-dihydro-	C ₈ H ₈ O	120
18	8.996	0.83	Triacetyl pentafluoropropionate	C ₃₃ H ₆₁ F ₅ O ₂	584
19	9.193	0.75	Hexatriacontyl trifluoroacetate	C ₃₈ H ₇₃ F ₃ O ₂	618
20	10.242	0.71	iso-Valeraldehyde propyleneglycol acetal 2	C ₈ H ₁₆ O ₂	144
21	11.115	57.39	1,2,3-Benzenetriol	C ₆ H ₆ O ₃	126
22	11.47	6.79	Phosphoric acid, bis(trimethylsilyl)monomethyl ester	C ₇ H ₂₁ O ₄ PSi ₂	256
23	12.348	0.21	Cycloheptasiloxane, tetradecamethyl-	C ₁₄ H ₄₂ O ₇ Si ₇	518
24	12.656	5.49	β -D-Glucopyranose, 1,6-anhydro-	C ₆ H ₁₀ O ₅	162
25	13.745	0.75	Diethyl Phthalate	C ₁₂ H ₁₄ O ₄	222
26	13.897	0.15	Triacetyl trifluoroacetate	C ₃₂ H ₆₁ F ₃ O ₂	534
27	16.304	0.75	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C ₂₀ H ₄₀ O	296
28	16.751	0.27	6-Octadecenoic acid, methyl ester, (<i>Z</i>)-		0
29	17.192	0.21	Pentadecanoic acid, 14-methyl-, methyl ester	C ₁₇ H ₃₄ O ₂	270
30	17.554	0.65	<i>n</i> -Hexadecanoic acid	C ₁₆ H ₃₂ O ₂	256
31	18.925	0.17	11,14,17-Eicosatrienoic acid, methyl ester	C ₂₁ H ₃₆ O ₂	320
32	19.285	0.26	9,12,15-Octadecatrien-1-ol, (<i>Z,Z,Z</i>)-	C ₁₈ H ₃₂ O	264
33	20.516	2.59	4-Dimethylsilyloxytridecane	C ₁₅ H ₃₄ OSi	258

4.7.4 GC-MS Analysis of *Chrysopogon aciculatus*

Chrysopogon aciculatus afforded the identification of compounds such as 2-Propanone, 1,3-dihydroxy- (29.07%), 2-Furancarboxaldehyde, 5-(hydroxymethyl)- (8.65%), β -D-Glucopyranose, 1,6-anhydro- (4.53%), L-Proline, N-(3-fluoro-5-trifluoromethylbenzoyl)-, octadecyl ester (4.29%), Cyclopentane, 1-acetyl-1,2-epoxy- (4.01%), Sucrose (3.97%), 2-Methoxy-4-vinylphenol (3.74%), *n*-Hexadecanoic acid (2.97%), 2,2'-Bis[N-[5-

diethylaminopent-2-yl]carbamyldiphenyl disulfide (2.65%), and Palladium(II), [(S)-O,N-bis(dicyclohexylphosphino)-2-pyrrolidinemethanol](dichloro)- (2.59%).




Figure 4.13 GC-MS Chromatogram of methanolic extract of *Chrysopogan aciculatus* whole plant.

Table 4.5 Phytochemicals identified in the methanolic extract of *Chrysopogan aciculatus* whole plant by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	2.533	2.07	Silane, trimethyl[5-methyl-2-(1-methylethyl)phenoxy]-		0
2	2.619	2.59	Palladium(II), [(S)-O,N-bis(dicyclohexylphosphino)-2-pyrrolidinemethanol](dichloro)-	C ₂₉ H ₅₃ Cl ₂ NOP ₂ Pd	669
3	2.723	4.29	L-Proline, N-(3-fluoro-5-trifluoromethylbenzoyl)-, octadecyl ester	C ₃₁ H ₄₇ F ₄ NO ₃	557
4	2.91	2.42	Acetamide, N-(2-methoxyphenyl)-		0
5	3.63	0.43	2-(Allylamino)-4,6-bis[2,2,2-trifluoro-1,1-bis(trifluoromethyl)ethyl]-1,3,5-triazine	C ₁₄ H ₆ F ₁₈ N ₄	572
6	4.222	29.07	2-Propanone, 1,3-dihydroxy-	C ₃ H ₆ O ₃	90
7	4.54	2.1	4,7-Bis(4-morpholino-1,2,5-thiadiazol-3-yloxymethyl)-1,3-dioxepane-5,6-diol	C ₁₉ H ₂₈ N ₆ O ₈ S ₂	532
8	4.773	1.25	6-Octadecenoic acid, methyl ester, (Z)-		0
9	5.523	2.65	2,2'-Bis[N-[5-diethylaminopent-2-yl]carbamyldiphenyl disulfide	C ₃₂ H ₅₀ N ₄ O ₂ S ₂	586

10	6.233	1.34	Maytansine	C ₃₄ H ₄₆ ClN ₃ O ₁₀	691
11	6.483	0.62	Phenanthrene-10-ethanamine, 3-bromo- β -hydroxy-N,N-diheptyl-, hydrosulfate	C ₃₀ H ₄₂ BrNO ₄ S	591
12	6.836	4.01	Cyclopentane, 1-acetyl-1,2-epoxy-	C ₇ H ₁₀ O ₂	126
13	7.047	0.97	Tetracontane, 3,5,24-trimethyl-	C ₄₃ H ₈₈	604
14	7.764	1.08	4H-Pyran-4-one, 2,3-dihydro-3,5-dihydroxy-6-methyl-	C ₆ H ₈ O ₄	144
15	8.802	1.16	Benzene, 1-isocyanato-2-methoxy-		0
16	8.984	8.65	2-Furancarboxaldehyde, 5-(hydroxymethyl)-	C ₆ H ₆ O ₃	126
17	9.192	0.93	14-Oxa-1,11-diazatetracyclo[7.4.1.0(2,7).0(10,12)]tetradeca-2,4,6-triene, 11-acetyl-6,9-bis(acetyloxy)-4-formyl-8-[(aminocarbonyloxy)methyl]-	C ₂₀ H ₂₁ N ₃ O ₉	447
18	9.816	1.55	Rhodium, chloro[ethenylidene(2,2-dimethylpropylidene)phosphinidyne]bis[tris(1-methylethyl)phosphine]-,(tb-5-22)-	C ₂₅ H ₅₃ ClP ₃ Rh	584
19	10.05	1.28	3-Heptanol	C ₇ H ₁₆ O	116
20	10.185	3.74	2-Methoxy-4-vinylphenol	C ₉ H ₁₀ O ₂	150
21	11.289	0.53	Isoquinoline, 3-methyl-, 2-oxide	C ₁₀ H ₉ NO	159
22	11.893	0.7	3,5-Dimethylanisole	C ₉ H ₁₂ O	136
23	11.977	0.68	1,2-Benzenedicarboxylic acid, diisooctyl ester		0
24	12.196	3.97	Sucrose	C ₁₂ H ₂₂ O ₁₁	342
25	12.602	4.53	β -D-Glucopyranose, 1,6-anhydro-	C ₆ H ₁₀ O ₅	162
26	13.402	0.78	3-tert-Butyl-4-hydroxyanisole	C ₁₁ H ₁₆ O ₂	180
27	13.737	0.87	Diethyl Phthalate	C ₁₂ H ₁₄ O ₄	222
28	14.101	0.43	1-Hexadecene		0
29	14.768	1.14	9 α -Hydroxysolasodine, o,N-diacetate	C ₃₁ H ₄₇ NO ₅	513
30	15.429	1.28	4-((1E)-3-Hydroxy-1-propenyl)-2-methoxyphenol	C ₁₀ H ₁₂ O ₃	180
31	16.305	0.98	Octadecanal	C ₁₈ H ₃₆ O	268
32	17.564	2.97	n-Hexadecanoic acid	C ₁₆ H ₃₂ O ₂	256
33	18.037	0.44	Phenol, 2,6-dimethyl-4-nitro-	C ₈ H ₉ NO ₃	167
34	19.023	0.44	Phytol	C ₂₀ H ₄₀ O	296
35	19.219	1.62	9-Octadecynoic acid	C ₁₈ H ₃₂ O ₂	280
36	23.177	0.42	13-Methyltriacontane	C ₃₄ H ₇₀	478
37	26.075	2.58	β -Sitosterol	C ₂₉ H ₅₀ O	414
38	26.375	0.63	Androst-4,6-dien-3,11,17-trione, 9-mercapto-	C ₁₉ H ₂₂ O ₃ S	330

39	29.231	2.06	4H-1-Benzopyran-4-one, 5-hydroxy-2-(4- hydroxyphenyl)-7-methoxy-	C ₁₆ H ₁₂ O ₅	284
----	--------	------	--	--	-----

4.7.5 GC-MS Analysis of *Rhododendron lepidotum*

Rhododendron lepidotum afforded the identification of compounds such as Benzoyl chloride, 4-pentyl- (13.12%), Verrucarol (8.4%), 2H-1-Benzopyran, 7-methoxy-2,2-dimethyl- (7.42%), alpha.-D-Glucopyranoside, O- α -D-glucopyranosyl-(1.fwdarw.3)- β -D-fructofuranosyl (5.27%), 1-[3-(1-Adamantyl)-1-methylpropylidene]thiosemicarbazide (5.04%), Benzene, 1-[1,1-dimethylethyl]-4-[2-propenyloxy]- (3.64%), Acetate, (2,4a,5,8a-tetramethyl-1,2,3,4,4a,7,8,8a-octahydro-1-naphthalenyl) ester (3.25%), 1-Buten-1-ol, 2-methyl-4-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, formate, (*E*-) (2.91%), *n*-Hexadecanoic acid (2.64%), and Octadecanal (2.49%).




Figure 4.14 GC-MS Chromatogram of methanolic extract of *Rhododendron lepidotum* leaf and stem.

Table 4.6 Phytochemicals identified in the methanolic extract of *Rhododendron lepidotum* leaf and stem by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	2.523	0.73	3-Carbethoxy-6- <i>n</i> -butyl-7-octadecylmercapto-4-quinolone	C ₃₄ H ₅₅ NO ₃ S	557
2	2.62	0.52	2-(N-Methyl-N-nitroamino)-4,6-bis[2,2,2-trifluoro-1,1-bis(trifluoromethyl)ethyl]-1,3,5-triazine	C ₁₂ H ₃ F ₁₈ N ₅ O ₂	591

3	4.068	1.57	Methylperfluoro[3-(4-ethylpiperazinyl)butyrate]	C ₁₁ H ₃ F ₁₉ N ₂ O ₂	556
4	5.583	0.89	2-Methyliminoperhydro-1,3-oxazine	C ₅ H ₁₀ N ₂ O	114
5	6.206	0.43	Benzene-1,3-diamine, N,N-diacetyl-5-[3,4,4,4-tetrafluoro-2-(1,2,2,2-tetrafluoro-1-trifluoromethylethyl)-1,3-di(trifluoromethyl)-1-butenyloxy]-	C ₁₉ H ₁₁ F ₁₇ N ₂ O ₃	638
6	6.771	0.49	Pseudosolasodine diacetate	C ₃₁ H ₄₉ NO ₄	499
7	8.079	1.37	Octadecanoic acid, octadecyl ester	C ₃₆ H ₇₂ O ₂	536
8	8.572	0.96	1,2-Benzenediol	C ₆ H ₆ O ₂	110
9	8.798	0.76	Benzofuran, 2,3-dihydro-	C ₈ H ₈ O	120
10	9.41	0.39	Hexatriacontane	C ₃₆ H ₇₄	506
11	9.472	0.55	1,2-Benzenediol, 3-methoxy-	C ₇ H ₈ O ₃	140
12	10.18	0.91	2-Methoxy-4-vinylphenol	C ₉ H ₁₀ O ₂	150
13	10.959	0.76	1,4-Benzenediol, 2-methyl-	C ₇ H ₈ O ₂	124
14	12.478	0.81	Hexatriacontyl trifluoroacetate	C ₃₈ H ₇₃ F ₃ O ₂	618
15	13.734	0.59	Diethyl Phthalate	C ₁₂ H ₁₄ O ₄	222
16	14.01	5.27	alpha.-D-Glucopyranoside, O- α -D-glucopyranosyl-(1.fwdarw.3)- β -D-fructofuranosyl	C ₁₈ H ₃₂ O ₁₆	504
17	14.09	1.45	1-Hexadecene		0
18	14.137	2.48	<i>n</i> -Decanoic acid	C ₁₀ H ₂₀ O ₂	172
19	14.182	5.04	1-[3-(1-Adamantyl)-1-methylpropylidene]thiosemicarbazide	C ₁₅ H ₂₅ N ₃ S	279
20	14.538	0.46	.tau.-Muurolol	C ₁₅ H ₂₆ O	222
21	15.13	0.42	Serverogenin acetate	C ₂₉ H ₃₆ O ₁₀	544
22	15.421	0.54	4-((1 <i>E</i>)-3-Hydroxy-1-propenyl)-2-methoxyphenol	C ₁₀ H ₁₂ O ₃	180
23	15.671	0.42	Aromadendrene oxide-(2)	C ₁₅ H ₂₄ O	220
24	16.187	0.52	2-Hydroxy-5-methylisophthalaldehyde	C ₉ H ₈ O ₃	164
25	16.304	2.49	Octadecanal	C ₁₈ H ₃₆ O	268
26	16.752	0.79	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C ₂₀ H ₄₀ O	296
27	17.134	8.4	Verrucarol	C ₁₅ H ₂₂ O ₄	266
28	17.197	0.41	Hexadecanoic acid, methyl ester		0
29	17.569	2.64	<i>n</i> -Hexadecanoic acid	C ₁₆ H ₃₂ O ₂	256
30	19.022	0.41	Phytol	C ₂₀ H ₄₀ O	296
31	19.298	2.07	3-Tetradecyn-1-ol	C ₁₄ H ₂₆ O	210
32	22.77	0.51	1-Heptacosanol	C ₂₇ H ₅₆ O	396

33	23.025	0.84	Hexadecanoic acid, 2,3-dihydroxypropyl ester	C ₁₉ H ₃₈ O ₄	330
34	23.186	2.2	Methylphosphonic acid, anhydride, bis(2,2-dimethylcyclohexyl) ester	C ₁₈ H ₃₆ O ₅ P ₂	394
35	23.247	2.05	6-Octadecenoic acid, methyl ester, (Z)-		0
36	23.436	0.42	1,1-Propanedicarbonitrile, 1,2-dicyclohexyl-	C ₁₇ H ₂₆ N ₂	258
37	23.531	13.12	Benzoyl chloride, 4-pentyl-	C ₁₂ H ₁₅ ClO	210
38	23.771	3.64	Benzene, 1-[1,1-dimethylethyl]-4-[2-propenyloxy]-	C ₁₃ H ₁₈ O	190
39	23.943	1.36	Cyclononasiloxane, octadecamethyl-		0
40	24.15	0.57	Succinimide, N-phenyl-3-(2-vinylcyclohexyl)-	C ₁₈ H ₂₁ NO ₂	283
41	24.537	2.18	2-Pentenamide, N-phenyl-	C ₁₁ H ₁₃ NO	175
42	24.6	2.91	1-Buten-1-ol, 2-methyl-4-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, formate, (E)-	C ₁₅ H ₂₄ O ₂	236
43	24.757	1.48	Neoisolongifolene, 8-oxo-	C ₁₅ H ₂₂ O	218
44	24.891	3.25	Acetate, (2,4a,5,8a-tetramethyl-1,2,3,4,4a,7,8,8a-octahydro-1-naphthalenyl) ester	C ₁₆ H ₂₆ O ₂	250
45	25.238	7.42	2H-1-Benzopyran, 7-methoxy-2,2-dimethyl-	C ₁₂ H ₁₄ O ₂	190
46	25.361	0.55	Di- <i>n</i> -octyl phenyl phosphate	C ₂₂ H ₃₉ O ₄ P	398
47	25.978	0.5	Squalene	C ₃₀ H ₅₀	410
48	29.132	0.58	γ-Tocopherol	C ₂₈ H ₄₈ O ₂	416
49	29.748	0.9	Stigmasteryl tosylate	C ₃₆ H ₅₄ O ₃ S	566
50	30.587	1.4	Vitamin E	C ₂₉ H ₅₀ O ₂	430
51	30.934	1.22	4-Hexen-1-ol, 6-(2,6,6-trimethyl-1-cyclohexenyl)-4-methyl-, (E)-	C ₁₆ H ₂₈ O	236

4.7.6 GC-MS Analysis of *Geranium wallichianum*

Geranium wallichianum afforded the identification of compounds such as 1,2,3-Benzenetriol (57.73%), 1,2-Benzenediol, 3-methoxy- (3.81%), 1,3,4,5-Tetrahydroxy-cyclohexanecarboxylic acid (3.5%), Pentadecanoic acid (3.48%), D-Allose (3.15%), Stigmast-5-en-3-ol, (3β)- (2.94%), 9,12-Octadecadienoic acid (Z,Z)-, methyl ester (2.72%), Hexadecanoic acid, methyl ester (2.33%), *cis*-Vaccenic acid (2.25%), and 9-Octadecenoic acid (Z)-, methyl ester (2.19%).




Figure 4.15 GC-MS Chromatogram of methanolic extract of *Geranium wallichianum* root.

Table 4.7 Phytochemicals identified in the methanolic extract of *Geranium wallichianum* root by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	5.337	1.11	Dimethyl dl-malate	C ₆ H ₁₀ O ₅	162
2	5.716	0.37	2,3-Dihydro-3,5-dihydroxy-6-methyl-4H-pyran-4-one	C ₆ H ₈ O ₄	144

3	7.004	2	5-Hydroxymethylfurfural	C ₆ H ₆ O ₃	126
4	7.701	3.81	1,2-Benzenediol, 3-methoxy-	C ₇ H ₈ O ₃	140
5	9.449	57.73	1,2,3-Benzenetriol	C ₆ H ₆ O ₃	126
6	10.863	3.15	D-Allose	C ₆ H ₁₂ O ₆	180
7	12.539	3.5	1,3,4,5-Tetrahydroxy-cyclohexanecarboxylic acid	C ₇ H ₁₂ O ₆	192
8	14.272	0.24	Benzoic acid, 4-hydroxy-3,5-dimethoxy-, hydrazide	C ₉ H ₁₂ N ₂ O ₄	212
9	14.752	1.35	Benzoic acid, 4-hydroxy-3,5-dimethoxy-	C ₉ H ₁₀ O ₅	198
10	15.783	2.33	Hexadecanoic acid, methyl ester	C ₁₇ H ₃₄ O ₂	270
11	16.146	3.48	Pentadecanoic acid	C ₁₅ H ₃₀ O ₂	242
12	17.47	2.72	9,12-Octadecadienoic acid (Z,Z)-, methyl ester	C ₁₉ H ₃₄ O ₂	294
13	17.519	2.19	9-Octadecenoic acid (Z)-, methyl ester	C ₁₉ H ₃₆ O ₂	296
14	17.735	0.36	Octadecanoic acid, methyl ester	C ₁₉ H ₃₈ O ₂	298
15	17.835	1.19	2-Chloroethyl linoleate	C ₂₀ H ₃₅ ClO ₂	342
16	17.877	2.25	<i>cis</i> -Vaccenic acid	C ₁₈ H ₃₄ O ₂	282
17	18.067	0.79	Octadecanoic acid	C ₁₈ H ₃₆ O ₂	284
18	19.78	0.3	Triacotanoic acid, methyl ester	C ₃₁ H ₆₂ O ₂	466
19	19.993	0.24	Cyclohexanone, 3-methyl-2-(1-methylethyl)-, <i>cis-trans</i>	C ₁₀ H ₁₈ O	154
20	20.521	0.49	1,3-Benzenediol, 5-pentadecyl-	C ₂₁ H ₃₆ O ₂	320
21	21.12	0.72	Succinic acid, dec-4-enyl dodecyl ester	C ₂₆ H ₄₈ O ₄	424
22	21.668	0.3	Abieta-8,11,13-trien-18-oic acid	C ₂₀ H ₂₈ O ₂	300
23	22.381	0.25	Abietic acid	C ₂₀ H ₃₀ O ₂	302
24	22.75	1.87	9-Octadecenal, (Z)-	C ₁₈ H ₃₄ O	266
25	23.139	0.23	Cyclopentane, 1-hexyl-3-methyl-	C ₁₂ H ₂₄	168
26	26.849	0.7	Squalene	C ₃₀ H ₅₀	410
27	29.282	0.12	9-Oxabicyclo[3.3.1]nonane, 2,6-diiodo-	C ₈ H ₁₂ I ₂ O	378
28	32.48	0.61	Vitamin E	C ₂₉ H ₅₀ O ₂	430
29	37.755	2.94	Stigmast-5-en-3-ol, (3 β)-	C ₂₉ H ₅₀ O	414
30	38.86	0.6	4,4,6A,6B,8A,11,11,14B-Octamethyl-1,4,4A,5,6,6A,6B,7,8,8A,9,10,11,12,12A,14,14A,14B-octadecahydro-2H-picen-3-one \$\$ Olean-12-en-3-one	C ₃₀ H ₄₈ O	424
31	40.481	0.77	Urs-12-ene	C ₃₀ H ₅₀	410
32	42.639	0.47	5 β -Androstan-3 α ,11 α ,17 β -triol	C ₁₉ H ₃₂ O ₃	308
33	45.934	0.84	Cyclohexane-1-methanol, 3,3-dimethyl-2-(3-methyl-1,3-butadienyl)-	C ₁₄ H ₂₄ O	208

Table 4.8 Phytocomponents identified in the methanolic extract of *Tsuga dumosa* bark by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	6.076	8.58	Benzoic acid	C ₇ H ₆ O ₂	122
2	8.931	1.57	6,6-Dimethylbicyclo[3.1.1]hept-2-ene-2-carboxylic acid	C ₁₀ H ₁₄ O ₂	166
3	10.932	4.56	D-Allose	C ₆ H ₁₂ O ₆	180
4	12.211	10.18	4-O-Methylmannose	C ₇ H ₁₄ O ₆	194
5	12.5	7.96	1,3,4,5-Tetrahydroxy-cyclohexanecarboxylic acid	C ₇ H ₁₂ O ₆	192
6	12.692	0.76	Phloroglucinol, trimethylsilyl ether	C ₉ H ₁₄ O ₃ Si	198
7	12.89	3.11	Methyl-(2-hydroxy-3-ethoxy-benzyl)ether	C ₁₀ H ₁₄ O ₃	182
8	13.42	10	Mome inositol	C ₇ H ₁₄ O ₆	194
9	15.783	0.36	Eicosanoic acid, methyl ester	C ₂₁ H ₄₂ O ₂	326
10	15.963	0.24	Cyclohexane, (2-methylpropyl)-	C ₁₀ H ₂₀	140
11	16.141	3.78	Pentadecanoic acid	C ₁₅ H ₃₀ O ₂	242
12	17.355	9.55	1-Naphthalenepropanol, α -ethenyldecahydro- α ,5,5,8a-tetramethyl-2-methylene-, [1S-[1 α (R*),4 α β ,8 α]]-	C ₂₀ H ₃₄ O	290
13	17.518	0.32	9-Octadecenoic acid (Z)-, methyl ester	C ₁₉ H ₃₆ O ₂	296
14	17.872	6.97	<i>cis</i> -Vaccenic acid	C ₁₈ H ₃₄ O ₂	282
15	18.065	1.17	Octadecanoic acid	C ₁₈ H ₃₆ O ₂	284
16	18.604	0.35	Undecanoic acid, 11-fluoro-, trimethylsilyl ester	C ₁₄ H ₂₉ FO ₂ Si	276
17	19.347	1.31	<i>n</i> -Nonadecanol-1	C ₁₉ H ₄₀ O	284
18	19.775	0.31	Octadecanoic acid, methyl ester	C ₁₉ H ₃₈ O ₂	298
19	20.221	1.82	Eicosanoic acid	C ₂₀ H ₄₀ O ₂	312
20	21.122	0.58	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C ₂₀ H ₄₀ O	296
21	21.671	1.02	Abieta-8,11,13-trien-18-oic acid	C ₂₀ H ₂₈ O ₂	300
22	22.384	0.87	Abietic acid	C ₂₀ H ₃₀ O ₂	302
23	22.755	0.69	16-Heptadecenal	C ₁₇ H ₃₂ O	252
24	23.147	0.34	1,2-Benzenedicarboxylic acid, diisooctyl ester	C ₂₄ H ₃₈ O ₄	390
25	25.156	1.74	1-Eicosanol	C ₂₀ H ₄₂ O	298
26	26.173	0.74	Hexanal, 3,3-dimethyl-	C ₈ H ₁₆ O	128
27	29.178	1.35	Docosanoic acid	C ₂₂ H ₄₄ O ₂	340
28	33.47	1.72	Tetrahydro geranyl acetate	C ₁₂ H ₂₄ O ₂	200

29	35.035	0.83	(-)-5-Oxatricyclo[8.2.0.0(4,6)]dodecane,12-trimethyl-9-methylene-, [1R-(1R*,4R*,6R*,10S*)]-	C ₁₅ H ₂₄ O	220
30	37.025	0.51	Dodecanoic acid, 2-phenylethyl ester	C ₂₀ H ₃₂ O ₂	304
31	37.732	3.66	Stigmast-5-en-3-ol, (3β)-	C ₂₉ H ₅₀ O	414
32	42.633	1.07	Androst-4-en-3-one, 17-hydroxy-, (17β)-	C ₁₉ H ₂₈ O ₂	288
33	51.876	8.89	Phenol, 2-methoxy-4-(2-propenyl)-, acetate	C ₁₂ H ₁₄ O ₃	206

4.7.8 GC-MS Analysis of *Anemone rivularis*

Anemone rivularis afforded the identification of compounds such as Diethyl Phthalate (7.35%), Cyclohexanecarboxylic acid, 2-hydroxy-1,6-dimethyl-, [1S-(1α,2β,6α)]- (6.01%), Decanoic acid, octyl ester (5.93%), 2(3H)-Furanone, dihydro-4-hydroxy- (4.7%), 3-Phenylpropanoic acid, 1-adamantylmethyl ester (4.69%), 1,3-Propanediol, 2-(hydroxymethyl)-2-nitro- (4.21%), Pentadecanoic acid (3.65%), β-D-Glucopyranose, 1,6-anhydro- (3.46%), Phenol, 4-(3-hydroxy-1-propenyl)-2-methoxy- (2.81%), 1,4-Benzenedicarboxylic acid, diethyl ester (2.69%).




Table 4.9 Phytocomponents identified in the methanolic extract of *Anemone rivularis* whole plant by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	4.521	0.85	Isosorbide Dinitrate	C ₆ H ₈ N ₂ O ₈	236
2	4.57	0.68	Acetic acid, 2-(N-methyl-N-phosphonomethyl)amino-	C ₄ H ₁₀ NO ₅ P	183
3	4.612	2.4	2,5-Dimethyl-4-hydroxy-3(2H)-furanone	C ₆ H ₈ O ₃	128
4	4.735	1.63	Pentanal	C ₅ H ₁₀ O	86
5	4.82	0.51	Butanamine, 2,2-dinitro-N-methyl-	C ₅ H ₁₁ N ₃ O ₄	177
6	4.922	1.43	3-Ethyl-1,3-dimethyldiaziridine (<i>trans</i>)	C ₅ H ₁₂ N ₂	100
7	5.045	1.03	<i>n</i> -Pentanol, 5-[2-methyl-1-cycloazapropyl]-	C ₈ H ₁₇ NO	143
8	5.108	1.01	4H-Pyran-4-one, 2,3-dihydro-3,5-dihydroxy-6-methyl-	C ₆ H ₈ O ₄	144
9	5.15	1.05	3-Heptanone, 5-ethyl-4-methyl-	C ₁₀ H ₂₀ O	156
10	5.287	4.7	2(3H)-Furanone, dihydro-4-hydroxy-	C ₄ H ₆ O ₃	102
11	5.47	0.68	Ethanedial, bis(dimethylhydrazone)	C ₆ H ₁₄ N ₄	142
12	5.527	0.77	2-Butanone, 4-hydroxy-3-methyl-	C ₅ H ₁₀ O ₂	102
13	5.587	1.59	1-Oxaspiro[3.5]nona-5,8-dien-7-one, 3-methylene-	C ₉ H ₈ O ₂	148
14	5.79	1.25	2-Furancarboxaldehyde, 5-(hydroxymethyl)-	C ₆ H ₆ O ₃	126
15	5.913	0.41	1,2,3-Propanetriol, 1-acetate	C ₅ H ₁₀ O ₄	134
16	6.352	1.37	2-Methyl-9-β-d-ribofuranosylhypoxanthine	C ₁₁ H ₁₄ N ₄ O ₅	282
17	6.645	2.15	Propanoic acid, 2-[(tetrahydro-2H-pyran-2-yl)oxy]-	C ₈ H ₁₄ O ₄	174
18	6.807	1.42	1,2:5,6-Dianhydrogalactitol	C ₆ H ₁₀ O ₄	146
19	7.46	0.1	1-(4-Hydroxy-6-methyl-2-pyrimidinyl)-2-aziridinone	C ₇ H ₇ N ₃ O ₂	165
20	7.626	0.88	Cycloheptasiloxane, tetradecamethyl-	C ₁₄ H ₄₂ O ₇ Si ₇	518
21	7.774	1.07	4-(2,4,4-Trimethyl-cyclohexa-1,5-dienyl)-but-3-en-2-one	C ₁₃ H ₁₈ O	190
22	7.975	4.21	1,3-Propanediol, 2-(hydroxymethyl)-2-nitro-	C ₄ H ₉ NO ₅	151
23	8.247	3.46	β-D-Glucopyranose, 1,6-anhydro-	C ₆ H ₁₀ O ₅	162
24	8.581	2.24	2,5-Dimethoxy-4-ethylamphetamine	C ₁₃ H ₂₁ NO ₂	223
25	8.78	2.69	1,4-Benzenedicarboxylic acid, diethyl ester	C ₁₂ H ₁₄ O ₄	222
26	8.976	7.35	Diethyl Phthalate	C ₁₂ H ₁₄ O ₄	222
27	9.247	1.71	Cyclohexanone, 2,3-dimethyl-2-(3-oxobutyl)	C ₁₂ H ₂₀ O ₂	196

28	9.293	2.16	Cyclooctasiloxane, hexadecamethyl-	$C_{16}H_{48}O_8Si_8$	592
29	9.563	5.93	Decanoic acid, octyl ester	$C_{18}H_{36}O_2$	284
30	9.787	4.69	3-Phenylpropanoic acid, 1-adamantylmethyl ester	$C_{20}H_{26}O_2$	298
31	9.865	2.58	4,6-Dimethyl-2(1H)pyridone, 3-(4,6-dibromo-2-hydroxyphenylmethylenamino)	$C_{14}H_{12}Br_2N_2O_2$	398
32	9.969	2.43	Hexadecanal	$C_{16}H_{32}O$	240
33	10.093	6.01	Cyclohexanecarboxylic acid, 2-hydroxy-1,6-dimethyl-, [1S-(1 α ,2 β ,6 α)]-	$C_9H_{16}O_3$	172
34	10.413	0.68	2-Amino-8-[3-d-ribofuranosyl]imidazo[1,2-a]-s-triazin-4-one	$C_{10}H_{13}N_5O_5$	283
35	10.483	1.7	Butanoic acid, 4-(2-oxocyclopentyl)- 4-(2-Oxocyclopentyl)butanoic acid	$C_9H_{14}O_3$	170
36	10.553	2.81	Phenol, 4-(3-hydroxy-1-propenyl)-2-methoxy-	$C_{10}H_{12}O_3$	180
37	10.813	0.64	Carbamodithioic acid, diethyl-, ethyl ester	$C_7H_{15}NS_2$	177
38	10.9	0.66	3-Buten-2-one, 4-(6,6-dimethyl-1-cyclohexen-1-yl)-	$C_{12}H_{18}O$	178
39	11.019	0.89	Cyclononasiloxane, octadecamethyl-	$C_{18}H_{54}O_9Si_9$	666
40	11.153	0.37	Tridecanoic acid, 12-methyl-, methyl ester	$C_{15}H_{30}O_2$	242
41	11.213	0.16	3-[3-Bromophenyl]-7-chloro-3,4-dihydro-10-hydroxy-1,9(2H,10H)-acridinedione	$C_{19}H_{13}BrClNO_3$	417
42	11.283	1.01	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	$C_{20}H_{40}O$	296
43	11.382	0.62	2-Pentadecanone, 6,10,14-trimethyl-	$C_{18}H_{36}O$	268
44	11.467	0.12	2-Butanone, 4-(2,2-dimethyl-6-methylenecyclohexyl)-	$C_{13}H_{22}O$	194
45	11.653	0.13	Benzoic acid, 2,5-bis(trimethylsiloxy)-, trimethylsilyl ester	$C_{16}H_{30}O_4Si_3$	370
46	12.267	0.72	Hexadecanoic acid, methyl ester	$C_{17}H_{34}O_2$	270
47	12.719	3.65	Pentadecanoic acid	$C_{15}H_{30}O_2$	242
48	12.821	0.79	Dibutyl phthalate	$C_{16}H_{22}O_4$	278
49	13.396	0.08	Methyl 10-methyl-hexadecanoate	$C_{18}H_{36}O_2$	284
50	14.183	0.55	10,13-Octadecadienoic acid, methyl ester	$C_{19}H_{34}O_2$	294
51	14.27	0.54	9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)-	$C_{19}H_{32}O_2$	292
52	14.374	1.39	Phytol	$C_{20}H_{40}O$	296
53	14.658	1.89	9,12-Octadecadienoic acid (Z,Z)-	$C_{18}H_{32}O_2$	280
54	14.742	1.64	9,12,15-Octadecatrienoic acid, (Z,Z,Z)-	$C_{18}H_{30}O_2$	278

55	14.921	0.61	Octadecanoic acid	C ₁₈ H ₃₆ O ₂	284
56	15.027	0.38	2-Isopropenyl-4,4,7a-trimethyl- 2,4,5,6,7,7a-hexahydro-benzofuran-6-ol	C ₁₄ H ₂₂ O ₂	222
57	15.26	0.13	Tricyclo[4.3.1.1(3,8)]undecan-3-amine	C ₁₁ H ₁₉ N	165
58	15.974	0.12	1-Bromo-4-bromomethyldecane	C ₁₁ H ₂₂ Br ₂	312
59	16.593	0.12	1-Tetradecanol, 14-chloro-	C ₁₄ H ₂₉ ClO	248
60	16.763	0.11	Retinoic acid	C ₂₀ H ₂₈ O ₂	300
61	17.454	0.14	Aromadendrene oxide-(2)	C ₁₅ H ₂₄ O	220
62	18.538	0.11	Hexadecanoic acid, 2-hydroxy-1- (hydroxymethyl)ethyl ester	C ₁₉ H ₃₈ O ₄	330
63	18.6	0.09	Octadecanoic acid, 5,9,13,17-tetramethyl-, methyl ester, [5R-(5R*,9R*,13R*)]-	C ₂₃ H ₄₆ O ₂	354
64	20.155	0.11	1-Nonadecene	C ₁₉ H ₃₈	266
65	20.246	0.17	<i>E,Z</i> -1,3,12-Nonadecatriene	C ₁₉ H ₃₄	262
66	20.347	0.09	7-Octynoic acid, methyl ester	C ₉ H ₁₄ O ₂	154
67	21.369	0.09	2,6,10,14,18,22-Tetracosahexaene, 2,6,10,15,19,23-hexamethyl-, (all- <i>E</i>)-	C ₃₀ H ₅₀	410
68	22.81	0.1	Anthraergostatetraenol hexahydrobenzoate	C ₃₅ H ₅₂ O ₂	504
69	23.218	0.12	Stigmasta-4,7,22-trien-3 β -ol	C ₂₉ H ₄₆ O	410
70	23.425	0.15	Tetracosamethyl-cyclododecasiloxane	C ₂₄ H ₇₂ O ₁₂ Si ₁₂	888
71	23.647	0.1	Isophthalic acid, allyl pentadecyl ester	C ₂₆ H ₄₀ O ₄	416
72	23.722	0.26	7-Dehydrosiosgenin 3-acetate	C ₂₉ H ₄₂ O ₄	454
73	24.148	0.36	17-(1,5-Dimethylhexyl)-10,13-dimethyl- 2,3,4,7,8,9,10,11,12,13,14,15,16,17- tetradecahydro-1H- cyclopenta[a]phenanthren-3-ol	C ₂₇ H ₄₆ O	386

4.7.9 GC-MS Analysis of *Scurrula elata*

Scurrula elata afforded the identification of compounds such as 1,2,3-Benzenetriol (25.12%), Diethyl Phthalate (18.14%), Benzoic acid, 2-amino-4-hydroxymethyl-, methyl ester (8.56%), Phthalic acid, di-(1-hexen-5-yl) ester (3.76%), γ -Sitosterol (3.15%), D-Allose (2.98%), 2-Cyclopenten-1-one, 2-butyl-3-methoxy- (2.34%), *trans*-3-Cyclopentene-1,2-dicarboxylic acid, 3-acetyl- 4-methyl-, diethyl ester (2.19%), 1-(3,3,3-Trifluoro-2-oxopropyl)-2(1H)-pyridinone (1.86%), and 3,4,5-Trimethyl-1H-pyrano[2,3-c]pyrazol-6-one (1.65%).




Figure 4.18 GC-MS Chromatogram of methanolic extract of *Scurrula elata* branches.

Table 4.10 Phytochemicals identified in the methanolic extract of *Scurrula elata* branches by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	4.523	0.28	Imidazole, 2-amino-5-[(2-carboxy)vinyl]-	C ₆ H ₇ N ₃ O ₂	153
2	6.455	0.22	5-O-Methyl-d-gluconic acid dimethylamide	C ₉ H ₁₉ NO ₆	237
3	8.22	1.2	1,3-Propanediol, 2-(hydroxymethyl)-2-nitro-	C ₄ H ₉ NO ₅	151
4	8.537	0.47	3-Chloropropionic acid, 2-isopropoxyphenyl ester	C ₁₂ H ₁₅ ClO ₃	242
5	8.569	0.27	2-Amino-2-[3-methyl-2-furyl]acetic acid	C ₇ H ₉ NO ₃	155
6	10.787	0.11	Phthalic acid, 2-ethoxyethyl pentadecyl ester	C ₂₇ H ₄₄ O ₅	448
7	10.889	25.12	1,2,3-Benzenetriol	C ₆ H ₆ O ₃	126
8	11.133	2.34	2-Cyclopenten-1-one, 2-butyl-3-methoxy-	C ₁₀ H ₁₆ O ₂	168
9	11.197	1.32	Imidazole, 5-methoxymethyl-4-nitro-	C ₅ H ₇ N ₃ O ₃	157
10	11.22	1.86	1-(3,3,3-Trifluoro-2-oxopropyl)-2(1H)-pyridinone	C ₈ H ₆ F ₃ NO ₂	205
11	11.283	0.67	1-(2-Pyrazinyl)-3-methyl-1-butanone	C ₉ H ₁₂ N ₂ O	164
12	11.36	0.87	Pyrazole-5-carboxylic acid, 3-methyl-	C ₅ H ₆ N ₂ O ₂	126

13	11.427	0.62	3-[(Cyclohexyl-methyl-amino)-methyl]-3H-benzooxazol-2-one	C ₁₅ H ₂₀ N ₂ O ₂	260
14	11.457	0.44	5-Furfurylidene hydantoin	C ₈ H ₆ N ₂ O ₃	178
15	11.51	0.53	2-Octenoic acid, 4-isopropylidene-7-methyl-6-methylene-, methyl ester	C ₁₄ H ₂₂ O ₂	222
16	11.597	0.17	Nicotinyl Alcohol	C ₆ H ₇ NO	109
17	12.09	0.08	Furazanamine, 4-azido-	C ₂ H ₂ N ₆ O	126
18	12.157	0.32	4-(2,4,4-Trimethyl-cyclohexa-1,5-dienyl)-but-3-en-2-one	C ₁₃ H ₁₈ O	190
19	12.55	2.98	D-Allose	C ₆ H ₁₂ O ₆	180
20	12.648	3.76	Phthalic acid, di-(1-hexen-5-yl) ester	C ₂₀ H ₂₆ O ₄	330
21	12.95	1.65	3,4,5-Trimethyl-1H-pyrano[2,3-c]pyrazol-6-one	C ₉ H ₁₀ N ₂ O ₂	178
22	13.057	0.87	1-Propanone, 1-(1,3-benzodioxol-5-yl)-3-(dimethylamino)-	C ₁₂ H ₁₅ NO ₃	221
23	13.097	0.2	Tricyclo[4.3.1.1(3,8)]undecane-1-carboxylic acid	C ₁₂ H ₁₈ O ₂	194
24	13.227	0.22	Phthalic acid, monoamide, N-ethyl-N-phenyl-, ethyl ester	C ₁₈ H ₁₉ NO ₃	297
25	13.664	18.14	Diethyl Phthalate	C ₁₂ H ₁₄ O ₄	222
26	13.853	8.56	Benzoic acid, 2-amino-4-hydroxymethyl-, methyl ester	C ₉ H ₁₁ NO ₃	181
27	14.003	1.08	Benzonitrile, 2-amino-4,5-diethoxy-	C ₁₁ H ₁₄ N ₂ O ₂	206
28	14.117	2.19	<i>trans</i> -3-Cyclopentene-1,2-dicarboxylic acid, 3-acetyl- 4-methyl-, diethyl ester	C ₁₄ H ₂₀ O ₅	268
29	14.174	0.89	Tricyclo[4.3.1.1(3,8)]undecane-1-carboxylic acid, methyl ester	C ₁₃ H ₂₀ O ₂	208
30	14.237	0.97	2,4,7,14-Tetramethyl-4-vinyl-tricyclo[5.4.3.0(1,8)]tetradecan-6-ol	C ₂₀ H ₃₄ O	290
31	14.3	0.59	1-(3,6,6-Trimethyl-1,6,7,7a-tetrahydrocyclopenta[c]pyran-1-yl)ethanone	C ₁₃ H ₁₈ O ₂	206
32	14.433	0.1	Acetic acid, 2,6,6-trimethyl-3-methylene-7-(3-oxobutylidene)oxepan-2-yl ester	C ₁₆ H ₂₄ O ₄	280
33	15.45	0.12	Furan-2-carboxylic acid, 5-(1-hexynyl)-	C ₁₁ H ₁₂ O ₃	192
34	16.016	0.79	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C ₂₀ H ₄₀ O	296
35	16.07	0.1	2-(1-Cyclohexenyl)cyclohexanone	C ₁₂ H ₁₈ O	178
36	16.464	0.33	1,4-Eicosadiene	C ₂₀ H ₃₈	278
37	16.918	0.21	Hexadecanoic acid, methyl ester	C ₁₇ H ₃₄ O ₂	270

38	17.31	0.36	1,2-Benzenedicarboxylic acid, butyl 2-ethylhexyl ester	C ₂₀ H ₃₀ O ₄	334
39	17.411	0.47	1,2-Benzenedicarboxylic acid, bis(2-methylpropyl) ester	C ₁₆ H ₂₂ O ₄	278
40	18.353	0.1	Carbonic acid, 2-ethoxyethyl neopentyl ester	C ₁₀ H ₂₀ O ₄	204
41	18.457	0.23	Pentadecyl trifluoroacetate	C ₁₇ H ₃₁ F ₃ O ₂	324
42	18.587	0.24	Cyclopropaneoctanoic acid, 2-[[2-[(2-ethylcyclopropyl)methyl]cyclopropyl)methyl]-, methyl ester	C ₂₂ H ₃₈ O ₂	334
43	18.623	0.19	l-(+)-Ascorbic acid 2,6-dihexadecanoate	C ₃₈ H ₆₈ O ₈	652
44	18.657	0.25	9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)-	C ₁₉ H ₃₂ O ₂	292
45	18.739	0.52	Phytol	C ₂₀ H ₄₀ O	296
46	18.96	0.13	2R,3S-9-[[1,3-Dihydroxy-4-fluoro-3-butoxy)methyl]guanine	C ₁₀ H ₁₄ FN ₅ O ₄	287
47	19.033	0.03	L-Lyxose	C ₅ H ₁₀ O ₅	150
48	20.621	0.21	trans-2-Dodecenoic acid	C ₁₂ H ₂₂ O ₂	198
49	24.383	0.16	Cyclohexane, 1,1'-(1,2-dimethyl-1,2-ethanediyl)bis-, (R*,R*)-(./-.-)	C ₁₆ H ₃₀	222
50	25.543	0.26	2,6,10,14,18,22-Tetracosahexaene, 2,6,10,15,19,23-hexamethyl-, (all-E)-	C ₃₀ H ₅₀	410
51	28.436	0.22	β-Tocopherol	C ₂₈ H ₄₈ O ₂	416
52	29.747	0.9	dl-α-Tocopherol succinate	C ₃₃ H ₅₄ O ₅	530
53	33.712	3.15	γ-Sitosterol	C ₂₉ H ₅₀ O	414
54	34.652	0.62	β-Amyrin	C ₃₀ H ₅₀ O	426

4.7.10 GC-MS Analysis of *Terminalia bellerica*

Terminalia bellerica afforded the identification of compounds such as 2-Methyl-9-β-d-ribofuranosylhypoxanthine (18.73%), Diethyl Phthalate (16.72%), 1,2,3-Benzenetriol (14.43%), Phenol, 4,4'-methylenebis[2,6-dimethyl- (10.94%), 1,4-Benzenedicarboxylic acid, diethyl ester (10.63%), Spiro[cyclopenta[d]-1,3,2-dioxaborin-4(5H),1'-cyclopentane], 2-ethyl-6,7-dihydro- (5.12%), Phthalic acid, ethyl 2-pentyl ester (2.73%), Phenol, 2-[(1-methylpropyl)thio]- (2.07%), Phthalic acid, ethyl non-5-yn-3-yl ester (1.87%), and 1-(6-Hydroxyimidazo[2,1-b]thiazol-5-yl)ethanone (1.06%).




Figure 4.19 GC-MS Chromatogram of methanolic extract of *Terminalia bellerica* fruit.

Table 4.11 Phytochemicals identified in the methanolic extract of *Terminalia bellerica* fruit by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	4.52	0.32	Imidazole, 2-amino-5-[(2-carboxy)vinyl]-	C ₆ H ₇ N ₃ O ₂	153
2	4.557	0.28	N-[2,2,2-Trifluoro-1-(isopropylamino)-1-(trifluoromethyl)ethyl]isovaleramide	C ₁₁ H ₁₈ F ₆ N ₂ O	308
3	4.589	0.66	Tricyclo[4.3.1.1(3,8)]undecane-1-carboxylic acid	C ₁₂ H ₁₈ O ₂	194
4	10.657	0.22	Menthone (2,4-dinitrophenyl)hydrazine	C ₁₆ H ₂₂ N ₄ O ₄	334
5	10.743	0.36	4-Hexenoic acid, 6-hydroxy-4-methyl-, methyl ester, (E)-	C ₈ H ₁₄ O ₃	158
6	10.757	0.33	Menthol, 1'-(butyn-3-one-1-yl)-, (1R,2S,5R)-	C ₁₄ H ₂₂ O ₂	222
7	10.81	1.06	1-(6-Hydroxyimidazo[2,1-b]thiazol-5-yl)ethanone	C ₇ H ₆ N ₂ O ₂ S	182
8	11.038	14.43	1,2,3-Benzenetriol	C ₆ H ₆ O ₃	126
9	11.197	10.94	Phenol, 4,4'-methylenebis[2,6-dimethyl-	C ₁₇ H ₂₀ O ₂	256
10	11.413	2.07	Phenol, 2-[(1-methylpropyl)thio]-	C ₁₀ H ₁₄ OS	182

11	11.493	0.61	2,2-Diethoxy-2-pyrazinyl-1-ethanol	C ₁₀ H ₁₆ N ₂ O ₃	212
12	11.543	0.81	2,5-Cyclohexadiene-1,4-dione, 2-hydroxy-6-(hydroxymethyl)-3,5-dimethyl-	C ₉ H ₁₀ O ₄	182
13	11.59	0.34	1-(4-Ethoxy-2-nitroanilino)-1-deoxy-β-d-mannopyranose	C ₁₄ H ₂₀ N ₂ O ₈	344
14	11.66	0.27	Pyrazole-5-carboxylic acid, 3-methyl-	C ₅ H ₆ N ₂ O ₂	126
15	13.013	0.13	5,6,7,8-Tetrahydro-5-oxo-2-hydroxyquinolin-4-yl acetic acid	C ₁₁ H ₁₁ NO ₄	221
16	13.04	0.28	Phenethylamine, 2,4,5-trimethoxy-α-methyl-	C ₁₂ H ₁₉ NO ₃	225
17	13.117	0.87	1-Carbazol-9-yl-3-(2-phenoxy-ethylamino)-propan-2-ol	C ₂₃ H ₂₄ N ₂ O ₂	360
18	13.443	10.63	1,4-Benzenedicarboxylic acid, diethyl ester	C ₁₂ H ₁₄ O ₄	222
19	13.644	16.72	Diethyl Phthalate	C ₁₂ H ₁₄ O ₄	222
20	13.823	18.73	2-Methyl-9-β-d-ribofuranosylhypoxanthine	C ₁₁ H ₁₄ N ₄ O ₅	282
21	14.143	5.12	Spiro[cyclopenta[d]-1,3,2-dioxaborin-4(5H),1'-cyclopentane], 2-ethyl-6,7-dihydro-	C ₁₂ H ₁₉ BO ₂	206
22	14.343	1.87	Phthalic acid, ethyl non-5-yn-3-yl ester	C ₁₉ H ₂₄ O ₄	316
23	14.53	2.73	Phthalic acid, ethyl 2-pentyl ester	C ₁₅ H ₂₀ O ₄	264
24	14.677	1.03	Perfluorooctanoic acid, 3,4-dimethylphenyl ester	C ₁₆ H ₉ F ₁₅ O ₂	518
25	14.75	0.77	Benzonitrile, 2-amino-4,5-diethoxy-	C ₁₁ H ₁₄ N ₂ O ₂	206
26	14.837	0.33	Thioxane-3-carbonitrile, 3-piperidyl-	C ₁₁ H ₁₈ N ₂ S	210
27	14.915	0.13	1,2-Benzenedicarboxylic acid, dihexyl ester	C ₂₀ H ₃₀ O ₄	334
28	16.942	0.13	Dodecanoic acid, 10-methyl-, methyl ester	C ₁₄ H ₂₈ O ₂	228
29	17.412	0.86	1,2-Benzenedicarboxylic acid, bis(2-methylpropyl) ester	C ₁₆ H ₂₂ O ₄	278
30	18.622	0.19	10-Octadecenoic acid, methyl ester	C ₁₉ H ₃₆ O ₂	296
31	23.464	0.15	2-Pentene, 3-diethylboryl-2-(methyl-1-propynylsilyl)-	C ₁₃ H ₂₅ BSi	220
32	23.817	0.15	[1,3,5]Triazine-2-carboxylic acid, 4-ethylamino-6-morpholin-4-yl-, amide	C ₁₀ H ₁₆ N ₆ O ₂	252
33	23.983	0.17	Silane, dimethyl(dimethyl(but-3-enyloxy)silyloxy)isobutoxy-	C ₁₂ H ₂₈ O ₃ Si ₂	276
34	24.161	0.32	Cyclononasiloxane, octadecamethyl-	C ₁₈ H ₅₄ O ₉ Si ₉	666
35	24.508	0.21	2-Myristinoyl-glycinamide	C ₁₆ H ₂₈ N ₂ O ₂	280
36	25.929	0.13	Tricyclo[3.3.1.1(3,7)]decane-2-ol-1-carboxylic acid, methyl ester	C ₁₂ H ₁₈ O ₃	210

4.7.11 GC-MS Analysis of *Ranunculus scleratus*

Ranunculus scleratus afforded the identification of compounds such as Diethyl Phthalate (19.41%), 2-Thiazolamine, 5-nitro- (18.4%), N-Aminopyrrolidine (8.07%), 1-Methylamino-1-deoxy-d-glucitol-N-thiocarboxylic acid 2-[1-[2-pyridyl]ethylidene]hydrazide (3.26%), 9,12,15-Octadecatrienoic acid, (Z,Z,Z)- (3.08%), 1,2,3,4,5,6,7,8-Octahydro-2-naphthol, 4-methylene-2,5,5-trimethyl- (3.06%), 9,12-Octadecadienoic acid (Z,Z)-, 2,3-dihydroxypropyl ester (3.04%), 1,4-Benzenedicarboxylic acid, diethyl ester (3%), Pentadecanoic acid (2.99%), and Phthalic acid, 3-methylphenyl tridecyl ester (2.92%).




Figure 4.20 GC-MS Chromatogram of methanolic extract of *Ranunculus scleratus* whole plant.

Table 4.12 Phytocomponents identified in the methanolic extract of *Ranunculus scleratus* whole plant by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	4.529	0.36	Imidazole, 2-amino-5-[(2-carboxy)vinyl]-	C ₆ H ₇ N ₃ O ₂	153
2	4.587	1.01	Diethylene glycol monododecyl ether	C ₁₆ H ₃₄ O ₃	274
3	4.697	0.2	Isoxazolidine, 4-ethyl-2,5-dimethyl-, <i>trans</i> -	C ₇ H ₁₅ NO	129
4	5.633	0.12	1,2-Dideoxy-l-erythro-pentitol	C ₅ H ₁₂ O ₃	120

5	5.783	0.12	2H-Azepin-2-one, hexahydro-1-methyl-	C ₇ H ₁₃ NO	127
6	7.048	2.25	1-Butanol, 3-methyl-, acetate	C ₇ H ₁₄ O ₂	130
7	7.244	0.56	1,4-Dioxaspiro[2.4]heptan-5-one	C ₅ H ₆ O ₃	114
8	7.586	0.11	4H-Pyran-4-one, 2,3-dihydro-3,5-dihydroxy-6-methyl-	C ₆ H ₈ O ₄	144
9	8.186	0.15	7,8,9,10-Tetrahydro-4-hydroxy-3-piperidinomethyl-11H,13H-benzo[4',5']thieno[3',2':4,5]pyrimidino[1,2-a]azepin-13-one	C ₂₁ H ₂₅ N ₃ O ₂ S	383
10	8.305	8.07	N-Aminopyrrolidine	C ₄ H ₁₀ N ₂	86
11	8.5	0.33	3(2H)-Furanone, dihydro-5-isopropyl-	C ₇ H ₁₂ O ₂	128
12	8.57	0.13	Acetamide, N-(syn-7-chloro-exo-bicyclo[2.2.1]hept-2-yl)-	C ₉ H ₁₄ ClNO	187
13	10.321	0.14	Tripropylpropoxysilane	C ₁₂ H ₂₈ OSi	216
14	12.429	0.77	D-Allose	C ₆ H ₁₂ O ₆	180
15	12.493	0.13	Adenine-9-propanoic acid, α -t-butoxycarbonylamino-	C ₁₃ H ₁₈ N ₆ O ₄	322
16	12.547	0.21	3-(Dimethylhydrazono)-2,4-dimethylhexane-2,5-diol	C ₁₀ H ₂₂ N ₂ O ₂	202
17	12.63	0.21	1-(3-Cyano-4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)-3-(4-fluorocinnamoyl)-2-thiourea	C ₁₉ H ₁₆ FN ₃ OS ₂	385
18	13.201	0.47	Benzo[b]dioxan, 4,4-dimethyl-5-nitro-3-methoxy-	C ₁₁ H ₁₃ NO ₅	239
19	13.363	3	1,4-Benzenedicarboxylic acid, diethyl ester	C ₁₂ H ₁₄ O ₄	222
20	13.56	19.41	Diethyl Phthalate	C ₁₂ H ₁₄ O ₄	222
21	14.06	3.26	1-Methylamino-1-deoxy-d-glucitol-N-thiocarboxylic acid 2-[1-[2-pyridyl]ethylidene]hydrazide	C ₁₅ H ₂₄ N ₄ O ₅ S	372
22	14.103	0.53	Phthalic acid, 2-ethoxyethyl isobutyl ester	C ₁₆ H ₂₂ O ₅	294
23	14.13	2.92	Phthalic acid, 3-methylphenyl tridecyl ester	C ₂₈ H ₃₈ O ₄	438
24	14.291	3.06	1,2,3,4,5,6,7,8-Octahydro-2-naphthol, 4-methylene-2,5,5-trimethyl-	C ₁₄ H ₂₂ O	206
25	14.41	1.38	Pyridine-2,6-dicarboxylic acid, diethylboryl ester	C ₁₁ H ₁₄ BNO ₄	235
26	14.511	1.2	Phthalic acid, 2-cyclohexylethyl ethyl ester	C ₁₈ H ₂₄ O ₄	304
27	14.568	1.15	2-Decen-1-ol, (Z)-	C ₁₀ H ₂₀ O	156
28	14.657	0.9	5-Amino-1,4-dihydro-quinoxaline-2,3-dione	C ₈ H ₇ N ₃ O ₂	177

29	14.735	2.42	Octadecanal	C ₁₈ H ₃₆ O	268
30	14.99	0.48	Tricyclo[4.3.1.1(3,8)]undecane-1-carboxylic acid	C ₁₂ H ₁₈ O ₂	194
31	15.063	0.42	4-Pyridinecarboxamide, 2-ethyl-	C ₈ H ₁₀ N ₂ O	150
32	15.13	0.22	2(3H)-Benzofuranone, 3,3-dimethyl-5-[(methylsulfonyl)oxy]-	C ₁₁ H ₁₂ O ₅ S	256
33	16.014	2.43	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C ₂₀ H ₄₀ O	296
34	16.077	0.25	2-Hexadecene, 3,7,11,15-tetramethyl-, [R-[R*,R*-(E)]]-	C ₂₀ H ₄₀	280
35	16.917	0.24	Hexadecanoic acid, methyl ester	C ₁₇ H ₃₄ O ₂	270
36	17.147	0.25	Gamolenic Acid	C ₁₈ H ₃₀ O ₂	278
37	17.29	2.99	Pentadecanoic acid	C ₁₅ H ₃₀ O ₂	242
38	17.403	1.12	1,2-Benzenedicarboxylic acid, bis(2-methylpropyl) ester	C ₁₆ H ₂₂ O ₄	278
39	18.572	0.11	10,13-Octadecadienoic acid, methyl ester	C ₁₉ H ₃₄ O ₂	294
40	18.646	0.12	9,12,15-Octadecatrienoic acid, methyl ester,	C ₁₉ H ₃₂ O ₂	292
41	18.735	1.27	Phytol	C ₂₀ H ₄₀ O	296
42	18.949	1.34	9,12-Octadecadienoic acid (Z,Z)-	C ₁₈ H ₃₂ O ₂	280
43	19.022	3.08	9,12,15-Octadecatrienoic acid, (Z,Z,Z)-	C ₁₈ H ₃₀ O ₂	278
44	19.175	0.85	Octadecanoic acid	C ₁₈ H ₃₆ O ₂	284
45	19.416	0.73	Hexanoic acid, 2-tetradecyl ester	C ₂₀ H ₄₀ O ₂	312
46	20.233	0.1	3-Cyclopentylpropionic acid, 2-dimethylaminoethyl ester	C ₁₂ H ₂₃ NO ₂	213
47	20.612	0.19	E-11-Hexadecenal	C ₁₆ H ₃₀ O	238
48	20.822	0.09	N,N'-Dicycloheptyl-1,2,4,5-tetrazine-3,6-diamine	C ₁₆ H ₂₈ N ₆	304
49	22.134	0.1	Hexanoic acid, 2-dimethylaminoethyl ester	C ₁₀ H ₂₁ NO ₂	187
50	22.537	0.34	1H-3a,7-Methanoazulen-5-ol, octahydro-3,8,8-trimethyl-6-methylene-	C ₁₅ H ₂₄ O	220
51	22.677	1.97	Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester	C ₁₉ H ₃₈ O ₄	330
52	22.93	0.98	Phosphorochloridic acid, diethyl ester	C ₄ H ₁₀ ClO ₃ P	172
53	23.302	18.4	2-Thiazolamine, 5-nitro-	C ₃ H ₃ N ₃ O ₂ S	145
54	24.023	0.27	Quinoline-1-carboxaldehyde, 1,2,3,4-tetrahydro-4,8-dimethyl-2-spirocyclohexane-	C ₁₇ H ₂₃ NO	257
55	24.103	0.98	Bicyclo[6.4.0]dodeca-1(8),4-diene-10,11-dicarboxylic acid	C ₁₄ H ₁₈ O ₄	250

Table 4.13 Phytocomponents identified in the methanolic extract of *Symplocos lucida* bark by GC-MS.

SN	R.Time	Area%	Name	Formula	MolWeight
1	4.526	0.3	Pterin-6-carboxylic acid	C ₇ H ₅ N ₅ O ₃	207
2	4.607	1.17	Phthalic acid, ethyl tridec-2-yn-1-yl ester	C ₂₃ H ₃₂ O ₄	372
3	5.01	0.23	3-Phenpropanol, 2'-hydroxy-3',4',6'-trimethyl	C ₁₂ H ₁₈ O ₂	194
4	6.098	0.5	Benzeneacetic acid, 4-hydroxy-3-methoxy-, methyl ester	C ₁₀ H ₁₂ O ₄	196
5	6.133	0.39	N-(4,6-Dimethyl-pyrimidin-2-yl)-N'-(3-methyl-butyl)-guanidine	C ₁₂ H ₂₁ N ₅	235
6	6.347	9.35	4-((1E)-3-Hydroxy-1-propenyl)-2-methoxyphenol	C ₁₀ H ₁₂ O ₃	180
7	6.767	0.93	Phthalic acid, ethyl 4-isopropylphenyl ester	C ₁₉ H ₂₀ O ₄	312
8	7.541	2.07	2,5,5,8a-Tetramethyl-6,7,8,8a-tetrahydro-5H-chromen-3-one	C ₁₃ H ₂₀ O ₂	208
9	7.747	3.71	Benzoic acid, 3-formyl-4,6-dihydroxy-2,5-dimethyl-, methyl ester	C ₁₁ H ₁₂ O ₅	224
10	7.958	1.2	Methyl tetra-p-nitrobenzate-β-O-galactopyranoside	C ₃₅ H ₂₆ N ₄ O ₁₈	790
11	8.372	2.44	1H-2-Indenone,2,4,5,6,7,7a-hexahydro-3-(1-methylethyl)-7a-methyl	C ₁₃ H ₂₀ O	192
12	8.572	0.79	Benzonitrile, 2-amino-4,5-diethoxy-	C ₁₁ H ₁₄ N ₂ O ₂	206
13	8.635	1.71	Phthalic acid, 2-ethoxyethyl dodecyl ester	C ₂₄ H ₃₈ O ₅	406
14	8.813	2.9	1,4-Benzenedicarboxylic acid, diethyl ester	C ₁₂ H ₁₄ O ₄	222
15	9.007	1.9	Diethyl Phthalate	C ₁₂ H ₁₄ O ₄	222
16	9.161	1.36	2,5-Dimethoxy-4-propoxybenzaldehyde	C ₁₂ H ₁₆ O ₄	224
17	9.361	0.85	Phthalic acid, decyl isobutyl ester	C ₂₂ H ₃₄ O ₄	362
18	9.48	0.24	Benzoic acid, 2,5-bis(trimethylsiloxy)-, trimethylsilyl ester	C ₁₆ H ₃₀ O ₄ Si ₃	370
19	11.195	0.48	1,2-Benzenedicarboxylic acid, bis(2-methylpropyl) ester	C ₁₆ H ₂₂ O ₄	278
20	11.518	0.45	Methyl (3,4-dimethoxyphenyl)(hydroxy)acetate	C ₁₁ H ₁₄ O ₅	226
21	11.732	1.1	3,5-Dimethoxy-4-hydroxycinnamaldehyde	C ₁₁ H ₁₂ O ₄	208
22	11.842	6.88	Benzenemethanol, 2,5-dimethoxy-, acetate	C ₁₁ H ₁₄ O ₄	210
23	15.804	2.83	Stigmasta-4,22-dien-3β-ol	C ₂₉ H ₄₈ O	412
24	15.941	0.53	Stigmasta-7,16-dien-3-ol, (3β,5α)-	C ₂₉ H ₄₈ O	412
25	16.021	0.8	Stigmasterol	C ₂₉ H ₄₈ O	412

26	17.148	0.16	Cedran-diol, 8S,14-	C ₁₅ H ₂₆ O ₂	238
27	17.767	0.19	2-(1,3-Dihydro-3,3-dimethyl-1-phenyl-3,4-benzofuran-1-yl)acetic acid	C ₁₈ H ₁₈ O ₃	282
28	18.127	0.48	Silanol, trimethyl-, phosphite (3:1)	C ₉ H ₂₇ O ₃ PSi ₃	298
29	18.207	0.22	1-[3,4-Dichlorophenyl]-3-[4,6-dimethyl-2-pyrimidinyl]guanidine	C ₁₃ H ₁₃ Cl ₂ N ₅	309
30	18.253	0.41	2-Isopropenyl-4,4,7a-trimethyl-2,4,5,6,7,7a-hexahydro-benzofuran-6-ol	C ₁₄ H ₂₂ O ₂	222
31	18.368	0.52	1-Bromo-3,7-dimethyloctane	C ₁₀ H ₂₁ Br	220
32	18.413	0.3	1-[N,N'-Bis(2-methylphenyl)amidino]aziridine	C ₁₇ H ₁₉ N ₃	265
33	18.483	0.44	16-Heptadecenal	C ₁₇ H ₃₂ O	252
34	18.693	2.96	Benzenemethanol, α -(1-phenylaminoethyl)-	C ₁₅ H ₁₇ NO	227
35	18.775	4.8	γ -Sitosterol	C ₂₉ H ₅₀ O	414
36	19.251	2.39	Cyclononasiloxane, octadecamethyl-	C ₁₈ H ₅₄ O ₉ Si ₉	666
37	19.43	3.06	Cholest-5-en-3-ol, 24-propylidene-, (3 β)	C ₃₀ H ₅₀ O	426
38	19.527	1.48	26,26-Dimethyl-5,24(28)-ergostadien-3 β -ol	C ₃₀ H ₅₀ O	426
39	19.587	1.86	Tyramine, N-formyl-	C ₉ H ₁₁ NO ₂	165
40	19.733	1.34	3 β ,5 α ,6 β -Trihydroxyandrostane-17-one	C ₁₉ H ₃₀ O ₄	322
41	19.88	4.26	3-(Prop-2-en-1-onyl)-2,4,4-trimethylcyclohex-2-en-1-one	C ₁₂ H ₁₆ O ₂	192
42	20.147	4.96	4-Hydroxy-3,5-dimethylbenzoic acid	C ₉ H ₁₀ O ₃	166

4.8 Antimicrobial Effects

The methanolic extracts of twelve medicinal plants displayed a dose dependent antimicrobial activity against one or more of the microorganisms tested (Table 4.14 to

Table 4.19). Maximum zone of inhibition against *S. aureus* and MRSA both were shown by *T. bellerica* while *K. pneumoniae* and *S. Typhi* were inhibited maximally by *L. involuta* and *T. dumosa* respectively. *L. involuta* did not show zone of inhibition against any of the gram positive bacteria taken. *S. elata* and *T. bellerica* were active against all the tested microorganisms while *S. lucida* only against MRSA. Ten of the twelve extracts inhibited at least three of the six organisms tested while eight extracts inhibited at least four organisms. *G. wallichianum* demonstrated effectiveness against five of the organisms tested except *K. pneumoniae*. *R. scleratus* did not inhibit any of the bacteria taken. Maximum zone of inhibition against *S. cerevisiae* and *Pichia sp.* were shown by *H. javanica* and *A. rivularis* respectively while *S. lucida* did not show the zone of inhibition against any of the fungi taken.

Most of the plant extracts were found active against gram positive bacteria and fungi while only some were found active to gram negative bacteria.




Figure 4.22 Antimicrobial activity of *T. dumosa* by agar well diffusion method as shown by the zone of inhibition against methicillin-resistant *Staphylococcus aureus* and *Salmonella Typhi*, respectively.

Table 4.14 Antibacterial activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against *Staphylococcus aureus* using agar well diffusion method.

Plant extracts in mg/ml	100	50	25	12.5	DMSO	PI100
<i>L. involuta</i>	-	-	-	-	-	24
<i>H. javanica</i>	8	7	6	6	-	24
<i>P. emblica</i>	13	9	8	7	-	23
<i>C. aciculatus</i>	-	-	-	-	-	23
<i>R. lepidotum</i>	11	10	9	8	-	23
<i>G. wallichianum</i>	10	8	7	6	-	22
<i>T. dumosa</i>	11	10	9	8	-	22
<i>A. rivularis</i>	7	6	-	-	-	23
<i>S. elata</i>	10	9	8	7	-	23
<i>T. bellerica</i>	16	15	14	13	-	20
<i>R. scleratus</i>	-	-	-	-	-	21
<i>S. lucida</i>	-	-	-	-	-	22

Table 4.15 Antibacterial activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against methicillin-resistant *Staphylococcus aureus* using agar well diffusion method.

Plant extracts in mg/ml	100	50	25	12.5	DMSO	MET5
<i>L. involuta</i>	-	-	-	-	-	-
<i>H. javanica</i>	7	6	-	-	-	-
<i>P. emblica</i>	20	18	17	15	-	-
<i>C. aciculatus</i>	8	7	6	-	-	-
<i>R. lepidotum</i>	11	10	9	7	-	-
<i>G. wallichianum</i>	13	12	11	9	-	-
<i>T. dumosa</i>	13	12	11	10	-	-
<i>A. rivularis</i>	-	-	-	-	-	-
<i>S. elata</i>	12	11	11	11	-	-
<i>T. bellerica</i>	21	20	19	17	-	-
<i>R. scleratus</i>	-	-	-	-	-	-
<i>S. lucida</i>	8	7	-	-	-	-

Table 4.16 Antibacterial activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against *Klebsiella pneumoniae* using agar well diffusion method.

Plant extracts in mg/ml	100	50	25	12.5	DMSO	PIT100/10
<i>L. involuta</i>	10	8	7	7	-	
<i>H. javanica</i>	-	-	-	-	-	
<i>P. emblica</i>	-	-	-	-	-	
<i>C. aciculatus</i>	9	8	7	6	-	
<i>R. lepidotum</i>	-	-	-	-	-	
<i>G. wallichianum</i>	-	-	-	-	-	
<i>T. dumosa</i>	-	-	-	-	-	9
<i>A. rivularis</i>	-	-	-	-	-	
<i>S. elata</i>	8	7	6	-	-	
<i>T. bellerica</i>	8	7	6	-	-	
<i>R. scleratus</i>	-	-	-	-	-	
<i>S. lucida</i>	-	-	-	-	-	8

Table 4.17 Antibacterial activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against *Salmonella* Typhi using agar well diffusion method.

Plant extracts in mg/ml	100	50	25	12.5	DMSO	PI100
<i>L. involuta</i>	-	-	-	-	-	15
<i>H. javanica</i>	-	-	-	-	-	15
<i>P. emblica</i>	-	-	-	-	-	15
<i>C. aciculatus</i>	11	10	9	8	-	15
<i>R. lepidotum</i>	-	-	-	-	-	13
<i>G. wallichianum</i>	8	7	-	-	-	15
<i>T. dumosa</i>	19	16	14	13	-	16
<i>A. rivularis</i>	8	7	-	-	-	12
<i>S. elata</i>	13	12	11	10	-	15
<i>T. bellerica</i>	11	9	7	6	-	16
<i>R. scleratus</i>	-	-	-	-	-	15
<i>S. lucida</i>	-	-	-	-	-	16

Table 4.18 Antifungal activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against *Saccharomyces cerevisiae* using agar well diffusion method.

Plant extracts in mg/ml	100	50	25	12.5	DMSO	Fluconazole (2.5mg/ml)
<i>L. involuta</i>	7	6	-	-	-	
<i>H. javanica</i>	11	11	9	6	-	
<i>P. emblica</i>	9	8	7	6	-	
<i>C. aciculatus</i>	6	-	-	-	-	24
<i>R. lepidotum</i>	-	-	-	-	-	25
<i>G. wallichianum</i>	9	7	7	-	-	25
<i>T. dumosa</i>	-	-	-	-	-	25
<i>A. rivularis</i>	10	8	8	7	-	26
<i>S. elata</i>	7	6	-	-	-	27
<i>T. bellerica</i>	9	9	8	7	-	25
<i>R. scleratus</i>	8	7	-	-	-	24
<i>S. lucida</i>	-	-	-	-	-	25

Table 4.19 Antifungal activity of methanolic extracts of 12 medicinal plants as shown by zone of inhibition (mm) against *Pichia sp.* using agar well diffusion method.

Plant extracts in mg/ml	100	50	25	12.5	DMSO	Fluconazole (2.5 mg/ml)
<i>L. involuta</i>	12	11	10	9	-	16
<i>H. javanica</i>	16	11	9	8	-	12
<i>P. emblica</i>	9	9	8	7	-	11
<i>C. aciculatus</i>	-	-	-	-	-	11
<i>R. lepidotum</i>	8	7	6	6	-	13
<i>G. wallichianum</i>	9	8	7	6	-	10
<i>T. dumosa</i>	8	7	-	-	-	15
<i>A. rivularis</i>	22	16	13	9	-	10
<i>S. elata</i>	8	7	-	-	-	12
<i>T. bellerica</i>	8	7	6	6	-	12
<i>R. scleratus</i>	9	9	8	7	-	16
<i>S. lucida</i>	-	-	-	-	-	12

4.9 Brine Shrimp Cytotoxicity

The degree of lethality was found to be directly proportional to the concentration of the extract (Figure 4.23). Of the twelve medicinal plant extracts tested nine extracts exhibited significant toxicity towards *Artemia salina* (brine shrimp) in which *A. rivularis* (58.11 µg/ml), *L. involuta* (91.72 µg/ml) and *T. bellerica* (94.65 µg/ml) with LC₅₀ value less than 100 µg/ml were highly toxic while *G. wallichianum* (135.90 µg/ml), *H. javanica* (195.68 µg/ml), *C. aciculatus* (252.82 µg/ml), *S. lucida* (292.19 µg/ml), *P. emblica* (306.27 µg/ml) and *S. elata* (400.35 µg/ml) with LC₅₀ value less than 500 µg/ml were moderately toxic. *T. dumosa* (9523.10 µg/ml) exhibited highest LC₅₀ value followed by *R. scleratus* (2069.84 µg/ml) and *R. lepidotum* (1607.17 µg/ml) which were found to be practically non-toxic (Figure 4.24).




Figure 4.23 Percentage mortality of *Artemia salina* (brine shrimp) larvae on treatment of methanolic extracts of 12 medicinal plants.




Figure 4.24 LC₅₀ of methanolic extracts of 12 medicinal plants on *Artemia salina* (brine shrimp) larvae.

4.10 MTT Cell Proliferation Assay

The methanol extracts of twelve medicinal plants exerted a dose dependant cytotoxicity against HeLa and MDCK cell lines. Among the plant extracts tested for growth inhibition (EC₅₀) of HeLa cells, *S. lucida* (52.22 µg/ml) was found to be the most effective followed by *T. dumosa* (122.3 µg/ml), *C. aciculatus* (177.6 µg/ml), *R. scleratus* (249.4 µg/ml), *S. elata* (304 µg/ml), *A. rivularis* (345.6 µg/ml), and *R. lepidotum* (389.7 µg/ml) while *T. bellerica* (1048 µg/ml) was found to be the least effective. In case of MDCK cells, *L. involuta* (240 µg/ml) was found to be most effective followed by *T. dumosa* (305 µg/ml), *A. rivularis* (407.7 µg/ml), *S. elata* (751.1 µg/ml), and *P. emblica* (925.3 µg/ml) while others with EC₅₀ value more than 1000 µg/ml were found to be less effective. Ten of the twelve medicinal plants including *S. lucida*, *C. aciculatus*, *R. scleratus*, *S. elata*, *R. lepidotum*, *G. wallichianum*, and *H. javanica* were found to be more effective against HeLa cells, however *L. involuta* and *P. emblica* inhibited MDCK cells more effectively.




Figure 4.25 Effects of *H. javanica* methanol extract on HeLa cell line with its decreasing concentration.

4.10.1 HeLa Cell Survival




Figure 4.26 Effects of methanolic extracts of 12 medicinal plants on HeLa cell survival.




Figure 4.27 Effects of methanolic extracts of 12 medicinal plants on HeLa cell survival.

Table 4.20 EC₅₀ of methanolic extracts of 12 medicinal plants on HeLa cell line.

Plant extracts	HeLa EC ₅₀ in µg/ml
<i>L. involuta</i>	412.9
<i>H. javanica</i>	574.9
<i>P. emblica</i>	991.6
<i>C. aciculatus</i>	177.6
<i>R. lepidotum</i>	389.7
<i>G. wallichianum</i>	560.1
<i>T. dumosa</i>	122.3
<i>A. rivularis</i>	345.6
<i>S. elata</i>	304
<i>T. bellerica</i>	1048
<i>R. scleratus</i>	249.4
<i>S. lucida</i>	52.22

4.10.2 MDCK Cell Survival




Figure 4.28 Effects of methanolic extracts of 12 medicinal plants on MDCK cell survival.




Figure 4.29 Effects of methanolic extracts of 12 medicinal plants on MDCK cell survival.

Table 4.21 EC₅₀ of methanolic extracts of 12 medicinal plants on MDCK cell line.

Plant extracts	MDCK EC ₅₀ in µg/ml
<i>L. involuta</i>	240
<i>H. javanica</i>	>1000
<i>P. emblica</i>	925.3
<i>C. aciculatus</i>	3521
<i>R. lepidotum</i>	4448
<i>G. wallichianum</i>	1105
<i>T. dumosa</i>	305
<i>A. rivularis</i>	407.7
<i>S. elata</i>	751.1
<i>T. bellerica</i>	1239
<i>R. scleratus</i>	1360
<i>S. lucida</i>	1389

5 Discussion

Nepal enjoys the richness of highly valued medicinal plants with varieties of local uses in different parts of the country. Other parts of the world also report their use in one or the other ways. Some researches have been made globally to test for the presence of novel bioactive compounds and their effects *in vitro* and *in vivo*. Our part of the world, however, has remained far behind in the race. This excludes us from the the benefits mother nature has provided.

Nature has preserved the plants with the myriads of biochemicals evolved during their struggle for survival – either against the adverse environmental conditions or against the rival organisms. The plants are thus the self made biochemical factories. Each chemical can find their use somewhere for human wellbeing. In medicine, the phytochemicals have an immense value. They find wider applications in modern day health care system.

Phytomedicine is in transistion from fringe to the mainstream medicine due to the presence of active principles in plants capable of promoting human health. WHO has also recognized its importance in the safe treatment of infectious as well as non-infectious diseases. The global acceptance of phytopharmaceuticals can also be advocated by their higher rate of FDA approval.

Traditional use of medicinal plants need some scientific validation for their continued and diversified use. Present study is the preliminary one to identify the bioactive compounds and biological activities of some of the underutilized medicinal plants of Nepal. The presence of a diverse range of phytocompounds, as indicated by GC-MS analysis and other phytochemical tests, and the presentation of important biological activities – such as antioxidant and anticancer – by the crude extracts in different concentrations suggest the great prospects of our medicinal plants for their wider applications in phytotherapy and phytopharmaceuticals.

5.1 Methanol Extraction

Medicinal plants possess a vast number of compounds which differ in their polarities. Different solvents with different polarities could have been used for their extraction such as water, ethanol, or acetone. We used methanol as an extraction solvent. Methanol is polar than water and ethanol. It is known to extract most of the active constituents from the plant material. It has been reported as an effective solvent to extract bioactive compounds such as phenolics (Siddhuraju & Becker, 2003). Another advantage is its easy evaporation resulting in quick concentration of the extracts.

The extraction yield of twelve medicinal plants in methanol came in the range from 4.39% to 32.48%, the highest of which was obtained from *T. bellerica* and the lowest from *C. aciculatus*.

H. javanica whole plant in methanol (9.85 %) in our study presented an extract yield somewhat similar to that previously reported in other solvents such as water (11.55%) and ethanol (12.95%) (Huang et al., 2008). Percentage yield of methanol extract of *P. emblica* bark (20.76 %) in our study has been found to be similar to the minimum of that reported from methanol extract of *P. emblica* fruits from six regions in China (21.0% to 39.4%) (Liu et al., 2008). However, it was found to be higher than that reported from *P. emblica* fruit in water (8.76%) (Ngamkitidechakul et al., 2010), and 70% methanol (4.2%) (Hazra et al., 2010). Percentage yield of methanol extract of *R. lepidotum* leaf and stem (7.12 %) in our study has been found to be somewhat similar to that reported from methanol extract of *R. lepidotum* leaves and twigs (6.07%) (Bhattarai et al., 2010). However, it was lower than that reported from methanol extract of *R. lepidotum* aerial parts (18.9%) (Rajbhandari et al., 2009), and higher than that reported from chloroform:methanol extract of *R. lepidotum* leaves and twigs (4.397%) (Bhattarai et al., 2010). Extract yield of *G. wallichianum* root (19.34 %) in methanol in our study has been found to be higher than that reported from *G. wallichianum* in 70% ethanol (2%) (Ahmad et al., 2003). Extract yield of *T. bellerica* fruit in methanol (32.48 %) in our study has been found to be somewhat similar to that reported from *T. bellerica* fruit in 70% ethanol (33% w/w) (Choudhary, 2008), and *T. bellerica* fresh leaves in 70% ethanol (30%) (Prabhu & Gopal, 2011; Vinoth et al., 2012). However it was found to be higher than that reported from *T. bellerica* fruit in 70% methanol (3.7%) (Hazra et al., 2010), ethanol (15%) (Manjunatha et al., 2011).

The choice of the solvent influences the extraction yield of medicinal plants. Compound to be contained in the extract is determined by the solvent used which in turn affects the biological activities of plant extracts (Gong et al., 2012; Jang et al., 2007). Other factors that may influence the biological activities include plant parts used, geography, harvest season, drying process, storage conditions, and post-harvest processing (Wendakoon et al., 2012). These factors also affect the extraction yield.

5.2 Total Phenolic Content

Phenolics are the most abundant structures in plants. Antioxidant compounds in plants are usually in the phenolic form. These important plant components can give up hydrogen atoms from their hydroxyl groups to radicals and form stable phenoxyl radicals, hence play an important role in antioxidant activity. Phenolics from intake of fruit and vegetable has been

found to reduce the risk of oxidative stress associated diseases such as cardiovascular diseases, cancer or osteoporosis (Hollman & Katan, 1999; Scalbert et al., 2005). Phenolic compounds can exhibit many biological activities such as an antioxidant, antimutagenic, anticarcinogenic, anti-inflammatory and antimicrobial. They can exert chemopreventive effects due to their free radical scavenging and metal chelating properties (Kampa et al., 2007).

We determined the total phenolic content of plant extracts by Folin-Ciocalteu method. It is a preferred method for the determination of total phenolic content (Singleton et al., 1999). This method allows the estimation of all the phenolic compounds including flavonoids, anthocyanins and nonflavonoids (Benvenuti et al., 2004). This assay is based on the transfer of electrons in alkaline medium from phenolic compounds to phosphomolybdic/phosphotungstic acid complexes to form the other type of blue complexes (possibly $(\text{PMoW}_{11}\text{O}_{40})^{4-}$) that are determined spectroscopically at approximately 765 nm (Singleton et al., 1999; Singleton & Rossi, 1965).

Other oxidation substrates present in the extract can interfere the total phenolics measurement in an inhibitory, additive or enhancing manner (Singleton et al., 1999; Singleton & Rossi, 1965). The inhibitory effects could be due to the oxidants competing with F-C reagent and/or air oxidation after the sample is made alkaline. For this reason, the F-C reagent is added ahead of alkali (Singleton et al., 1999). Additive effects occur from unanticipated phenols, aromatic amines, high sugar levels or ascorbic acid in the samples. However, despite these disadvantages, the F-C assay is simple and reproducible and has been widely used for quantification of phenolic compounds in plant extracts.

Total phenolic content shown by the plant extracts ranged from 6.74 mgGAE/g to 55.66 mgGAE/g. The highest amount of phenolic content was found in *P. emblica* (55.66 ± 0.84 mgGAE/g) followed by *G. wallichianum* (54.99 ± 0.74 mgGAE/g), *S. elata* (54.71 ± 0.68 mgGAE/g), *T. bellerica* (54.69 ± 0.77 mgGAE/g), and *T. dumosa* (54.19 ± 0.93 mgGAE/g) while the lowest was observed in *R. scleratus* (6.74 ± 0.62 mgGAE/g).

Total phenolic content in methanolic extract of *P. emblica* bark (55.66 ± 0.84 mgGAE/g) in our study has been found to be lower than that reported in methanolic extracts of *P. emblica* fruit from six regions of China (81.5 to 120.9 mgGAE/g) (Liu et al., 2008) in addition to that in water and 70% methanol extract of *P. emblica* fruits (34.22 g gallic acid/100g and 215.60 mg/ml GAE/100 mg respectively) (Charoenteeraboon et al., 2010; Hazra et al., 2010). Similarly, TPC observed in methanolic extract *T. bellerica* fruit (54.69 ± 0.77 mgGAE/g) in our study has been lower than that reported in 70% methanol extract of *T. bellerica* fruit (133.00 mg/ml GAE/100

mg) (Hazra et al., 2010). Similarly, TPC in methanolic extract of *R. scleratus* whole plant (6.74 ± 0.62 mgGAE/g) in our study has been lower than that previously reported from whole plant of *R. sceleratus* ethyl acetate fraction (97.1 GAE /mg/g), *n*-butanol fraction (79.6 GAE/mg/g), chloroform fraction (73.3 GAE mg/g), aqueous fraction (64.6 GAE/mg/g), and *n*-hexane fraction (23.5 GAE/mg/g) (Shahid et al., 2015). Similarly, water and ethanol extract of *H. nepalensis* whole plant (115.77 and 74.53 μ g CE/mg respectively) (Huang et al., 2008) have been reported to contain higher amount of total phenolics.

We already know that phenolic compounds are responsible for antioxidant activities. Phenolic compounds present in our plant extracts may help prevent cancer, diabetes, heart disease, neurodegeneration, and other age related degenerative diseases. Further *in vitro* and *in vivo* tests after isolation and elucidation of specific phenolic compounds may open new doors for other infinite possibilities.

5.3 Total Flavonoid Content

Flavonoids are the most abundant polyphenolic compounds ubiquitously found in plants. Plants rich in flavonoids are good source of antioxidants which effectively scavenge the reactive oxygen species due to the structure and the substitution pattern of phenolic hydroxyl groups (Cao et al., 1997; Miller & Ruiz-Larrea, 2002; Sharififar et al., 2008; Subhasree et al., 2009). Flavonoids have been reported to exhibit a wide range of biological activities such as antioxidant, anti-inflammatory, antimicrobial, anti-angiogenic, anticancer and anti-allergic (Anyasor et al., 2010; Ayoola et al., 2008; Barile et al., 2007; Chao et al., 2002; Hossain et al., 2011; Igbinsosa et al., 2009; Kokate, 2008).

We determined the total flavonoid content in plant extracts by spectrophotometric method using aluminium chloride in the presence of potassium acetate. This is one of the most commonly used approaches for the determination of total flavonoid content in food or medicinal plant samples. It involves the formation of aluminium-flavonoid complexes, with flavonols (quercetin, morin, kaempferol and rutin) and flavones luteolin exhibiting the maximum absorbance at 415–425 nm (Pełal & Pyrzynska, 2014). The variation of this method in the presence of NaNO₂ shows specificity for rutin, luteolin and catechins (Pełal & Pyrzynska, 2014).

Total flavonoid content for each plant extract was observed at a point within the narrow range from 7.51 mgQE/g to 25.65 mgQE/g. The highest amount of flavonoid content was found in *P. emblica* (25.65 ± 0.77 mgQE/g) followed by *S. elata* (22.26 ± 0.69 mgQE/g), *C. aciculatus* (19.83 ± 0.94 mgQE/g), *G. wallichianum* (18.53 ± 0.67 mgQE/g), *R. lepidotum* (18.35

± 0.76 mgQE/g), and *H. javanica* (18.24 ± 0.83 mgQE/g) while the lowest was observed in *T. bellerica* (7.51 ± 0.54 mgQE/g).

Similar study on flavonoid content in methanol extracts of *P. emblica* fruit from six regions in China has reported it to vary from 20.3 to 38.7mgQE/g (Liu et al., 2008) which is similar to our result. However, TFC previously reported in 70% methanol extract of *P. emblica* fruit (176.00 mg/ml QE/100 mg) (Hazra et al., 2010), 70% methanol extract of *T. bellerica* fruit (138.30 mg/ml QE/100 mg) (Hazra et al., 2010) have been found to be higher than that in our study. Similarly, water and ethanol extracts of *H. nepalensis* whole herb (31.48 and 66.07 μ g RE/mg respectively) (Huang et al., 2008) have been reported to contain high amount of total flavonoids.

We detected a lower amount of flavonoid content in our plant extracts which might be due to the presence of a lower amount flavonols (quercetin, morin, kaempferol and rutin) and flavones luteolin in the plant extracts since this method is specific for these compounds in the presence of potassium acetate (Pękal & Pyrzynska, 2014). However, there is still the chance that they display the higher amount of rutin, luteolin and catechins when tested in the presence of NaNO₂ (Pękal & Pyrzynska, 2014).

We observed that our plant extracts demonstrated antimicrobial, cytotoxic and anticancer activities. Though present in a small amount, flavonoids have a part to play whatever be the degree of their activities. Other in-depth studies are needed to further confirm their presence, and to isolate and elucidate them to perform *in vitro* and *in vivo* studies.

5.4 Antioxidant Activity

An organism's metabolism fights against oxidative effects with its own antioxidant defence systems. Elimination and neutralisation of reactive oxygen species is handled by both enzymatic and non-enzymatic antioxidant mechanisms. Natural antioxidants have a diversity of biochemical activities, some of which include the inhibition of reactive oxygen species generation, direct or indirect scavenging of free radicals, and alteration of intracellular redox potential (Finkel & Holbrook, 2000). Many antioxidant compounds are known to have anticancer effects (Dragsted et al., 1993; Johnson et al., 1994). Plants rich in secondary metabolites, including phenolics, flavonoids and carotenoids, have antioxidant activity due to their redox properties and chemical structures.

We examined the antioxidant activity of the plant extracts by their ability to scavenge DPPH free radicals and presented it as IC₅₀ value which is the concentration of the extract necessary

to inhibit 50% of the free radicals. Thus, an extract with the lower IC₅₀ value possess the higher capacity to neutralize the free radicals. DPPH is an organic free radical that does not disintegrate in water, methanol, or ethanol. The odd electron in the DPPH radical is responsible for its deep purple colour and the maximum absorbance at 517nm (Kumarasamy et al., 2007). DPPH can easily receive an electron or hydrogen from antioxidant molecules to become a stable diamagnetic molecule (Soares et al., 1997). It results in discoloration from purple to yellow due to the decreasing quantity of DPPH radicals which indicates the radical scavenging nature of the extract (Guo et al., 2007; Molyneux, 2004).

DPPH assay is a quick, reliable and reproducible method widely used to test the ability of compounds to act as free radical scavengers or hydrogen donors, and to evaluate antioxidant activity of plant extracts (Mosquera et al., 2007). The free radical scavenging activities of extracts depend on the ability of antioxidant compounds to lose hydrogen and the structural conformation of these components (Fukumoto & Mazza, 2000; Shimada et al., 1992).

Plant extracts displayed a dose dependent antioxidant activity with their IC₅₀ values ranging from 4.54 µg/ml to 195.55 µg/ml. *T. dumosa* (4.54 ± 0.18 µg/ml) exhibited the highest antioxidant activity as shown by the lowest IC₅₀ value – lower than that observed in ascorbic acid (4.57 ± 00 µg/ml) – followed by *P. emblica* (6.64 ± 0.71 µg/ml), *G. wallichianum* (7.24 ± 0.48 µg/ml), *T. bellerica* (12.08 ± 0.76 µg/ml), and *S. elata* (14.20 ± 0.94 µg/ml). *R. scleratus* (195.55 ± 2.45 µg/ml) showed the highest IC₅₀ value with the lowest antioxidant activity.

Methanol extract of *G. wallichianum* root (6.9 µg/ml) in our study has been found to possess higher scavenging activity (IC₅₀) than that previously reported in ethanol, *n*-butanol, and water extracts of *G. wallichianum* whole plant (19.05, 24.133 and 25.35 µg/mL) (Ismail et al., 2009). Methanol extract of *P. emblica* bark (7.14 µg/ml) in our study has been found to possess higher scavenging activity (IC₅₀) than that previously reported in water extract of *P. emblica* fruits (51.3 µg/ml) (Charoenteeraboon et al., 2010) and lower scavenging activity than that previously reported in 70% methanol extract of *P. emblica* fruits (1.43 µg/ml) (Hazra et al., 2010). DPPH radical scavenging activity (EC₅₀) of methanolic extracts of *P. emblica* fruit from six regions in China have been reported to vary from 11.23 to 45.44 µg/mL (Liu et al., 2008). Methanol extract of *T. bellerica* fruit (11.37 µg/ml) in our study has been found to possess lower scavenging activity (IC₅₀) than that previously reported in 70% methanol extract of *T. bellerica* fruit (IC₅₀ 1.45µg/ml) (Hazra et al., 2010). Methanolic extract of *H. javanica* whole plant (144.39 µg/ml) in our study has been found to possess higher scavenging activity (IC₅₀) than that previously reported in ethanol extract of *H. nepalensis* whole plant (314.51 µg/mL) and lower scavenging activity than that previously reported in water extract

of *H. nepalensis* whole plant (84.20 µg/mL) (Huang et al., 2008). DPPH radical scavenging activity (% inhibition at a given concentration) has been reported for *n*-hexane (84.6% at 1000 µg/ml), chloroform (77.9% at 500 µg/ml), ethyl acetate (80.9% at 30 µg/ml), *n*-butanol (80.8% at 60 µg/ml), and remaining aqueous (77.1% at 1000 µg/ml) fractions of whole plant of *R. sceleratus* (Shahid et al., 2015).

We identified a number of antioxidant compounds by GC-MS analyses. Phytochemical investigations also showed the presence of phenolic and flavonoid compounds. These compounds are known to display antioxidant activities due to their redox properties (Rice-Evans et al., 1995; Rice-Evans et al., 1996; Soobrattee et al., 2005). This indicates that our plants can have great implication in isolation of therapeutic agents to prevent oxidative stress related degenerative diseases.

5.4.1 Correlation of TPC and TFC with Antioxidant Activity

Several studies have reported a positive correlation between total phenolic content and antioxidant activity in plants (Bouayed et al., 2007; Djeridane et al., 2006; Kim et al., 2003; Lim & Quah, 2007; Tawaha et al., 2007). There are also studies which report that there is no positive relationship (Hesam et al., 2012; Rafat et al., 2010). A study on methanolic extracts of *P. emblica* fruit from six regions in China has presented good correlations of antioxidant activity with total phenolic content and total flavonoid content (Liu et al., 2008). Such correlations have also been reported from fruit extracts of *P. emblica* and *T. belerica* (Hazra et al., 2010).

In our study, we observed a significant correlation between the antioxidant activity and the total phenolic content while comparing the TPC and the IC₅₀ of DPPH RSA of twelve medicinal plants ($r = -0.9684$, $p < 0.0001$). However, we could not establish a good correlation between antioxidant activity and total flavonoid content. Our results suggest that total phenolic contents of these plants highly attributed their antioxidant activities.

5.5 Antimicrobial Activity

In recent years, multiple drug resistant pathogenic microorganisms have been developed due to the indiscriminate use of commercial antimicrobial drugs commonly used in the treatment of infectious diseases, making it a growing problem worldwide. In addition to this problem, antibiotics are sometimes associated with adverse effects on host including hypersensitivity, immune suppression and allergic reactions (Khanahmadi et al., 2010). Therefore, there is a need to develop alternative antimicrobial drugs from various sources such as medicinal plants for the treatment of infections.

We analysed antimicrobial effects of plant extracts against six different microbial strains viz. *Staphylococcus aureus*, methicillin resistant *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Salmonella Typhi*, *S. cerevisiae* and *Pichia sp.* by agar well diffusion method. This method is widely used to examine the antimicrobial effects of plant extracts. Plant extracts are introduced into the wells in the agar plate whose agar surface is already swabbed with uniform microbial inoculum. The antimicrobial compounds diffuse during incubation to inhibit the growth of microbial strains showing the zone of inhibition on the agar surface. The size of inhibition zone is proportional to the activity shown by the extracts.

We observed that maximal inhibition of *S. aureus* and MRSA both were exerted by *T. bellerica*. Maximum inhibition of *K. pneumoniae* and *S. Typhi* were shown by *L. involuta* and *T. dumosa*. However, *L. involuta* was found ineffective against any of the gram positive bacteria. Antimicrobial activity of *S. lucida* was observed only against MRSA while *S. elata* and *T. bellerica* inhibited all the tested microorganisms. It was found that inhibition of at least three of the six microbes were exerted by ten of the twelve medicinal plants. Eight of the extracts inhibited at least four of the total organisms. *G. wallichianum* was ineffective against *K. pneumoniae* but it inhibited other five organisms. *R. scleratus* could not inhibit any of the bacteria. *S. lucida* did not inhibit any of the fungi. Maximum inhibition of *S. cerevisiae* and *Pichia sp.* were exerted by *H. javanica* and *A. rivularis*, respectively.

Methanol extract of *H. javanica* whole plant in our study showed antimicrobial activity against *Staphylococcus aureus* (8mm) similar to that previously reported from methanol extracts of *H. javanica* leaf (81 and 162 mg/mL, 30 μ l) against *S. aureus* (7.30 and 8.50 mm) (Mandal et al., 2016). Antibacterial activity of methanol extract of *P. emblica* bark against *Staphylococcus aureus* (13 mm) in our study has been found to be supported by previous studies in methanol extract of leaves and fruits of *P. emblica* against *S. aureus* (31.00 and 35.66 mm) (Javale & Sabnis, 2010), methanol extract of fruits of *P. emblica* (10mg/ml) against *Staphylococcus aureus* (18mm) (Raghu & Ravindra, 2010). Methanol extract of *P. emblica* bark in our study was found to be inactive against *Klebsiella pneumoniae*. However, previous studies have reported significant activities of methanol extract of leaves and fruits of *P. emblica* against *K. pneumoniae* (30.00, and 26.17 mm) (Javale & Sabnis, 2010), and methanol extract of fruits of *P. emblica* (10mg/ml) against *Klebsiella pneumoniae* (19mm) (Raghu & Ravindra, 2010). Antibacterial activity of methanol extract of *P. emblica* bark in our study against *Staphylococcus aureus* (13 mm) has been supported by a previous report on antibacterial activity of flavonoids and saponins from *Emblia officinalis* fruit and *Emblia officinalis* leaf against *S. aureus* (Javale & Sabnis, 2010). Methanol extract of *P. emblica* bark in our study was found to be inactive against *Klebsiella pneumoniae*. Flavonoids and saponins from

Emblica officinalis leaf have also been reported to be inactive against *K. pneumoniae*, however, that from *Emblica officinalis* fruit have been reported to be active (Javale & Sabnis, 2010).

Methanol extract of *G. wallichianum* root in our study was found to be active against *Staphylococcus aureus* (10mm) which have been supported by similar studies on 70% ethanolic extract of *G. wallichianum* against *Staphylococcus aureus* (15mm) (Ahmad et al., 2003), and methanol extract of *G. wallichianum* rhizomes (1mg/ml, 100 and 200 μ L) against *S. aureus* (14.33 mm) (Ismail et al., 2012). Methanol extract of *G. wallichianum* root in our study showed very low activity against *Salmonella* Typhi (8mm) which is somewhat similar to the previous reports on inactivities of 70% ethanolic extract of *G. wallichianum* against *Salmonella* Typhi (Ahmad et al., 2003), and inactivity of methanol extract of *G. wallichianum* rhizomes (1mg/ml, 100 and 200 μ L) against *S. Typhi* (Ismail et al., 2012). Methanol extract of *G. wallichianum* leaves (1mg/ml, 100 and 200 μ L) has been reported to be inactive against *S. aureus* and, *S. Typhi* (Ismail et al., 2012). Methanol extract of *G. wallichianum* root in our study showed no activity against *Klebsiella pneumoniae* which is similar to the previous report on inactivity of 70% ethanolic extract of *G. wallichianum* against *Klebsiella pneumoniae* (Ahmad et al., 2003).

Antimicrobial activity of methanol extract of *T. bellerica* fruit against *Staphylococcus aureus* (16mm) in our study has been supported by the previous study on methanol extract of *T. bellirica* fruit pericarp (50 and 100mg/ml) against *S. aureus* (18 and 30mm) by paper disc diffusion (Nagendraswamy et al., 2013). Antimicrobial activity of methanol extract of *R. lepidotum* leaf and stem against *Staphylococcus aureus* (11mm) in our study has been supported by previous study on methanol extract of *R. lepidotum* leaves and twigs against *S. aureus* (13mm) by filter disc diffusion (Bhattarai et al., 2010). Antibacterial activities of methanol extract of *R. lepidotum* leaf and stem against *Staphylococcus aureus* (11mm), and methicillin resistant *Staphylococcus aureus* (11mm) in our study has been found to be supported by a report on antibacterial activity of coumarin compounds from methanolic extract of *R. lepidotum* against *Staphylococcus aureus*, and methicillin resistant *Staphylococcus aureus* (Shakeel-U-Rehman et al., 2010).

Methanol extract of *R. scleratus* whole plant was found to be inactive against *Staphylococcus aureus* and *Salmonella* Typhi in our study, however on a previous report, chloroform, ethyl acetate, *n*-butanol and aqueous fractions of whole plant of *R. sceleratus* have been reported to inhibit *S. aureus* (12.0, 10.8, 11.2, and 10.5 μ g/200 μ l), and *S. Typhi* (10.4, 10.8, 10.4, and 11.3 μ g/200 μ L) by 96-well microplate assay (Shahid et al., 2015).

There are studies which report the inhibitory activities of some of these medicinal plants against some other fungal strains such as *G. wallichianum* against *Trichophyton Schoenleinii*, *Pseudallescheria Boydii*, *Candida albicans*, *Aspergillus niger*, *Microsporium canis*, and *Fusarium solani* (Ahmad et al., 2003; Ismail et al., 2012), *R. sceleratus* against *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Trichophyton tonsurans*, *Microsporium gypseum*, and *Microsporium fulvum* (Sharma et al., 2012), and *H. javanica* against *Candida albicans* (Sood & Yadav, 2014).

We found our plant extracts to be more effective against the gram-positive bacteria than to the gram-negative. It might be described based on the fact that gram-negative bacteria have an effective permeability barrier, comprised of a thin lipopolysaccharide outer membrane which render the cell wall impermeable to lipophilic solutes including the plant extracts. In addition, gram-negative bacteria also have inherent overexpressed or multiple efflux pumps that prevent the intracellular accumulation of antibacterial agents. Gram-positive bacteria, on the other hand, have a mesh-like peptidoglycan layer which is more accessible to permeation by the extracts (Burt, 2004; Qadan et al., 2005; Rameshkumar et al., 2007; Stefanello et al., 2008).

It can be said of the plants that they are what they contain. Antimicrobial activities shown by the plants must have been contributed by the compounds they possess. We identified a variety of antimicrobial compounds by GC-MS analyses. Phytochemical investigations also showed the presence of phenolic and flavonoid contents. Phenolics can inhibit hydrolytic enzymes or other interactions that inactivate microbial adhesins, cell envelope transport proteins and non-specific interactions with carbohydrates (Pyla et al., 2010). Flavonoids can form complex with extracellular and soluble proteins and with bacterial cell walls or disrupt microbial membranes (Cowan, 1999). Other compounds have similar inhibitory actions. Tannins can bind to proline rich proteins and interfere with the protein synthesis (Shimada, 2006). Saponins can cause leakage of proteins and certain enzymes from the cell (Zablotowicz et al., 1996). Steroids may associate with membrane lipid and cause leakages from liposomes (Epanand et al., 2007).

The results clearly show that our plant extracts can find potential therapeutic uses in isolation of broad-spectrum antimicrobial compounds to fight the war against drug resistant microbial pathogens. The presumed safety of the plant derived drugs signifies their great market potential.

5.6 Brine Shrimp Cytotoxicity

Toxicity of pharmaceutical compounds possess a great risk on public health. Brine shrimp lethality has been used as a very useful tool for testing general toxicity (Almeida et al., 2002). It is used in determination of bioactivity of plant products (Lhullier et al., 2006; McLaughlin et al., 1991; Meyer et al., 1982; Stefanello et al., 2006), and synthetic compounds (Almeida et al., 2002). It has found an application in detection and isolation of bioactive compounds from plant extracts (Mackeen et al., 2000; Meyer et al., 1982; Solis et al., 1993; Krishnaraju et al., 2005; Ogunnusi & Dosumu, 2008). It has been known to have a good correlation with cytotoxic, antitumor and pesticidal activities (McLaughlin et al., 1998; Pelka et al., 2000).

Standard brine shrimp lethality bioassay stipulates that an LC_{50} value < 1000 $\mu\text{g/ml}$ is considered bioactive in toxicity evaluation of plant extracts (Clarkson et al., 2004). Based on this benchmark, we found that nine of the twelve medicinal plants exhibited significant and dose dependent toxicity to brine shrimps in which *A. rivularis* (58.11 $\mu\text{g/ml}$), *L. involuta* (91.72 $\mu\text{g/ml}$) and *T. bellerica* (94.65 $\mu\text{g/ml}$) presented high toxicity with LC_{50} value less than 100 $\mu\text{g/ml}$ while *G. wallichianum* (135.90 $\mu\text{g/ml}$), *H. javanica* (195.68 $\mu\text{g/ml}$), *C. aciculatus* (252.82 $\mu\text{g/ml}$), *S. lucida* (292.19 $\mu\text{g/ml}$), *P. emblica* (306.27 $\mu\text{g/ml}$) and *S. elata* (400.35 $\mu\text{g/ml}$) exerted moderate toxicity with LC_{50} value less than 500 $\mu\text{g/ml}$. *T. dumosa* (9523.10 $\mu\text{g/ml}$), *R. scleratus* (2069.84 $\mu\text{g/ml}$), and *R. lepidotum* (1607.17 $\mu\text{g/ml}$) identified themselves as non-toxic showing the LC_{50} values higher 1000 $\mu\text{g/ml}$.

Cytotoxicity (LC_{50}) of methanol extract *G. wallichianum* root (136 $\mu\text{g/ml}$) in our study has been found to be higher than that previously reported in methanol and 70% ethanol extracts of *G. wallichianum* rhizome (>1000 $\mu\text{g/ml}$) (Ismail et al., 2012; Ahmad et al., 2003), and ethyl acetate fraction of *G. wallichianum* rhizome (LC_{50} 333.256 $\mu\text{g/ml}$) (Ismail et al., 2012). Cytotoxicity of methanolic extract of *P. emblica* bark (306 $\mu\text{g/ml}$) in our study has been found to be lower than that reported in ethanolic extract of *P. emblica* fruit (60 $\mu\text{g/ml}$) (Hossen et al., 2014). Cytotoxicity of methanol extract of *R. lepidotum* leaf and stem (1607 $\mu\text{g/ml}$) in our study has been found to be lower than that reported in methanol extract of *R. lepidotum* leaves and twigs (111.59 $\mu\text{g/ml}$) (Bhattarai et al., 2010).

We already know that brine shrimp lethality shows a good correlation with insecticidal, cytotoxic, and anticancer properties. Higher lethality shown by our plant extracts may suggest that they contain a number of bioactive compounds with insecticidal and anticancer properties. Higher cytotoxicity shown by the plants may also pose a risk on human health. Their suitability for human consumption needs to be ascertained by several in-depth researches.

5.7 Anticancer Activities

Cancer is one of the most important diseases of the 21st century. The incidence of cancer is increasing during the last decades in most countries. In spite of numerous efforts to combat this disease, world medicine is facing with the lack of perfect cure for it. Many chemotherapeutic drugs are presently placed in a predicament of reduced therapeutic effect due to the problem of drug resistance (Peters et al., 2002). Chemotherapeutic drugs also exert toxicity to normal cells, which in turn causes the unpleasant side effects to the patients.

In the last decades, there has been a great emphasis towards the researches on complementary and alternative medicine for cancer management (Guevara et al., 1999). Several studies have been conducted on medicinal herbs to find new antitumor compounds. It has been reported that about 3000 plants with anticancer properties have been used for anticancer drugs (Hartwell, 1971; Pandey, 2002).

Numerous compounds are reported from plants with anticancer properties such as phenolics, flavonoids, alkaloids, phenylpropanoids, and terpenoids (Kintzios, 2006; Park et al., 2008). Plants contain almost unlimited capacity to generate compounds that fascinates researchers in the quest for novel chemotherapeutics (Reed & Pellecchia, 2005). The persistent search for new anticancer compounds in medicinal plants is a realistic and promising strategy for its prevention (Hu et al., 2009).

We determined the antiproliferative activities of plant extracts against HeLa and MDCK cell lines by MTT assay. This assay is based on the conversion of yellow MTT tetrazolium salt to purple insoluble formazan crystals. Dehydrogenase enzymes from mitochondria and endoplasmic reticulum in viable cells are involved in the reaction (Fotakis & Timbrell, 2006; Slater et al., 1963). Cells which have undergone death due to toxic damage cannot transform MTT into formazan. So, the amount of formazan produced is directly proportional to the number of viable cells.

We observed that the methanol extracts of twelve medicinal plants demonstrated a dose dependant growth inhibition of HeLa and MDCK cell lines. HeLa cell line was most effectively inhibited (EC_{50}) when exposed to *S. lucida* (52.22 $\mu\text{g/ml}$) followed by *T. dumosa* (122.3 $\mu\text{g/ml}$), *C. aciculatus* (177.6 $\mu\text{g/ml}$), *R. scleratus* (249.4 $\mu\text{g/ml}$), *S. elata* (304 $\mu\text{g/ml}$), *A. rivularis* (345.6 $\mu\text{g/ml}$), and *R. lepidotum* (389.7 $\mu\text{g/ml}$) while it was least affected by the presence of *T. bellerica* (1048 $\mu\text{g/ml}$). MDCK cell line was susceptible to the highest inhibition when exposed to *L. involuta* (240 $\mu\text{g/ml}$) followed by *T. dumosa* (305 $\mu\text{g/ml}$), *A. rivularis* (407.7 $\mu\text{g/ml}$), *S. elata* (751.1 $\mu\text{g/ml}$), and *P. emblica* (925.3 $\mu\text{g/ml}$) while it was less affected by the presence

of other extracts with their EC_{50} values higher than 1000 $\mu\text{g}/\text{ml}$. Ten of the twelve medicinal plants including *S. lucida*, *C. aciculatus*, *R. scleratus*, *S. elata*, *R. lepidotum*, *G. wallichianum*, and *H. javanica* exhibited higher inhibition against HeLa cell line in compared to the MDCK cell line to which only *L. involuta* and *P. emblica* showed higher inhibition. This suggests the usefulness of these plants in cancer chemoprevention.

In our study, methanol extract of leaf and stem of *R. lepidotum* was found to be effective (EC_{50}) against HeLa cell line (389.7 $\mu\text{g}/\text{ml}$) but not much effective against MDCK cell line (4448 $\mu\text{g}/\text{ml}$). In a similar study, inhibitory effects of *R. lepidotum* aerial parts or leaves and twigs have been reported against MDCK cell line (CC_{50} of >100 $\mu\text{g}/\text{ml}$) (Rajbhandari et al., 2009), and HeLa cell line (Bhattarai et al., 2010) by using MTT assay. Methanol extracts of *P. emblica* bark in our study was found to show lower inhibitory effects against HeLa cell line (EC_{50} 991.6 $\mu\text{g}/\text{ml}$) using MTT assay than that previously reported from water extracts of *P. emblica* (LC_{50} of 87.5 $\mu\text{g}/\text{ml}$) using Sulforhodamine B assay (Ngamkitidechakul et al., 2010). These plants have also been reported to show activities against other cell lines such as *H. nepalensis* against human hepatoma 3B cell line (Huang et al., 2008), *R. lepidotum* against Vero cell line (Rajbhandari et al., 2009), and *P. emblica* against A549, HepG2, MDA-MB-231, SKOV3, and SW620 human cancer cell lines (Ngamkitidechakul et al., 2010).

Compounds isolated from these plants have been reported to exert inhibitory effects against different cancer cell lines such as triterpenes and triterpene saponins from *H. nepalensis* against HL-60, SMMC-7721, A-549, MCF-7 and SW480 (Ma et al., 2016), flavonoid named aciculatin from *C. aciculatus* against HCT116 (Lai et al., 2012), apigenin derivatives and other compounds from *Chrysopogon aciculatis* against MCF-7, H460, HT-29, and CEM (Shen et al., 2012), saponins and other compounds from *A. rivularis* against HL-60, HepG2, A549 and HeLa (Wang et al., 2014).

We identified a number of anticancer compounds by GC-MS analyses. We also demonstrated that phenolic compounds possessed by plants attributed their antioxidant activities which in turn may unveil anticancer effects. Further in-depth research may help unravel the compounds responsible for anticancer activities shown by plants.

5.8 GC-MS analysis

The GC-MS analysis of twelve medicinal plants afforded the identification of a wide range fatty acids, esters, sugars, steroidal, phenolic, and heterocyclic compounds with bioactivities reported in most of them by different researchers.

1,2,3-Benzenetriol (Syn. Pyrogallol) was identified from *G. wallichianum* (57.73%), *P. emblica* (57.39%), *S. elata* (25.12%), and *T. bellerica* (14.43%). It is a polyphenol compound and a hydrolysable tannin (Khanbabaee & van Ree, 2001) obtained through ingestion of tea and coffee (Müller et al., 2006) and degradation of gallic acid in colon (Yasuda et al., 2000). It exhibits anticancer activity against MCF-7, PC-3, LNCaP, DU145 and HCT-116 cell lines (Chew et al., 2014). It has been reported as an antioxidant which increases H₂O₂ resistance associated with a reduction in intracellular oxidation and protein carbonylation (Mendes et al., 2015). It also acts as a superoxide generator which has been reported to enhance iNOS gene expression (Kuo et al., 2000).

9,12-Octadecadienoic acid (Z,Z)- was identified from *H. javanica* (29.21%), *A. rivularis* (1.89%), and *R. scleratus* (1.34%). It is a linoleic acid ester. It has been reported to exhibit anti-inflammatory, hypocholesterolemic, cancer preventive, hepatoprotective, nematocide, insectifuge, antihistaminic antieczemic, antiacne, 5-alpha reductase inhibitor antiandrogenic, antiarthritic, anticoronary activities (Maruthupandian & Mohan, 2011; Parthipan et al., 2015; Rajeswari et al., 2012).

2-Propanone, 1,3-dihydroxy- was identified from *C. aciculatus* (29.07%). It is an aliphatic ketone. It has been reported to be used in artificial respiration, and found to possess antioxidant and analgesic activities (Sivakumar & Gajalakshmi, 2014).

Diethyl phthalate was identified from *R. scleratus* (19.41%), *S. elata* (18.14%), *T. bellerica* (16.72%), *A. rivularis* (7.35%), *S. lucida* (1.9%), *L. involuta* (1.11%), *C. aciculatus* (0.87%), *P. emblica* (0.75%), and *R. lepidotum* (0.59%). It has been reported to be used for the preparation of about 67 consumer formulations including bath preparations (oils, tablets, and salts), eye shadow, toilet waters, perfumes, other fragrance preparations, skin care preparations and also as a component in insecticide sprays, mosquito repellents and camphor substitute (Jayaraman et al., 2011). It has been reported to possess antifouling, antimicrobial (Diaz & Thilaga, 2016), acetylcholinesterase and neurotoxic activities (Velanganni et al., 2011). It has also been found to reduce growth rate, food consumption and increase organ weights in human (Hauser & Calafat, 2005).

2-Thiazolamine, 5-nitro- (Syn. Thiazole, 2-amino-5-nitro-) was identified from *R. scleratus* (18.4%) in this study. It has been found to be evaluated as a hypoxic radiosensitizer using bacteria (Rockwell et al., 1982). It has been found to be effective as antihistomonad in turkeys and against trichomoniasis in pigeons (O'Neil, 2006).

n-Hexadecanoic acid (Syn. palmitic acid) was identified from *H. javanica* (15.53%), *C. aciculatus* (2.97%), *L. involuta* (2.78%), *R. lepidotum* (2.64%), and *P. emblica* (0.65%). It is a fatty acid reported to exhibit antioxidant, hypocholesterolemic, antiinflammatory, antibacterial (Konovalova et al., 2013; Sermakkani & Thangapandian, 2012), nematicide, pesticide, lubricant, antiandrogenic, flavor, hemolytic and 5- α reductase inhibitor activities (Parthipan et al., 2015; Rajeswari et al., 2012).

Phenol, 4,4'-methylenebis[2,6-dimethyl- was identified from *T. bellerica* (10.94%). It is an antioxidant compound which has also been reported to be used in fuel, polymers, and as an additive in petroleum-based lubricants (Revathi et al., 2013).

4-O-Methylmannose was identified from *T. dumosa* (10.18%). This methylated sugar has been reported to possess antibacterial activity (Kumar & Bhaskar, 2012).

Mome inositol was identified from *T. dumosa* (10%). It has been reported to display antiallopecic, anticirrhotic, antineuropathic, cholesterolytic, lipotropic (Kumar et al., 2012) and antiproliferative activities (Neda et al., 2013).

1-Naphthalenepropanol, α -ethenyldecahydro- α ,5,5,8a-tetramethyl-2-methylene-, [1S-[1 α (R*),4 α β ,8 $\alpha\alpha$]]- (Syn. 13-Epimanool) was identified from *T. dumosa* (9.55%). This labdane type diterpene finds its use as rosin or sizing agents (Salem et al., 2015).

4-((1E)-3-Hydroxy-1-propenyl)-2-methoxyphenol was identified from *S. lucida* (9.35%), *C. aciculatus* (1.28%), and *R. lepidotum* (0.54%). This phenolic compound possess antimicrobial, antioxidant, antiinflammatory and analgesic activities (Gopalakrishnan & Vadivel, 2011).

A furan aldehyde named 2-Furancarboxaldehyde, 5-(hydroxymethyl)- is a useful product found in the fossil fuel resources (Andreia et al., 2011). It was identified it from *L. involuta* (9%), *C. aciculatus* (8.65%), and *A. rivularis* (1.25%) in this study. It has been reported to exhibit antifungal, antibacterial (Oskoueian et al., 2011), anti-inflammatory (Brustugun et al., 2005; Li & Lu, 2005; Xu et al., 2007), clastogenic, uterotonic (Gopalakrishnan & Udayakumar, 2014) and preservative properties (Gopalakrishnan & Vadivel, 2011; Lilhore & Pande, 2016). It has been found to display protective effects against CCl₄-induced injury to the liver and vascular endothelium (Graff & Pollack, 2005) and cytotoxicity against BGC823 cell line (Guo L et al., 2016).

Phenol, 2-methoxy-4-(2-propenyl)-, acetate (Syn. eugenol acetate or eugenyl acetate) was identified from *T. dumosa* (8.89%). It has been reported to show antioxidant and antibacterial

activity (Vanin et al., 2014). It has been known to be toxic and repellent to red imported fire ants *Solenopsis invicta* (Kafle & Shih, 2013).

Benzoic acid was identified from *T. dumosa* (8.58%). It is an aromatic carboxylic acid which has been reported to be used in perfumes, flavorings and as an anti-fungal agent (Rajeswari & Rani, 2014-15).

Verrucarol is a sesquiterpene portion of several macrocyclic dilactones which was identified from *R. lepidotum* (8.4%) in this study. It has been reported as a mycotoxin (Johanning et al., 2002; Tuomi et al., 2000) with potent anticancer activity (Kraus & Frazier, 1980). Certain derivatives of verrucarol are reported to inhibit viral infection (Coochon et al., 1980; Snider & Amin, 1978).

1,3,4,5-tetrahydroxy-cyclohexanecarboxylic acid (Syn. Quinic acid) is a phenolic acid which was identified in this study from *T. dumosa* (7.96%), and *G. wallichianum* (3.5%). It is used in the synthesis of anti-influenza (anti-swine flu) medication and shows anti-aging effects. It is also used in treatment of prostate Cancer (Inbathamizh & Padmini, 2013; Pero & Lund, 2009). It exhibits anti-inflammatory activities (Kumar et al., 2015; Zeng et al., 2009). It is reported to exert a potent antibacterial and antifungal activities against typical microbial strains affecting the human respiratory or urinary tract, including *Candida* species and also exhibits anti-PI-3, anti-herpes effects and cytotoxicity against MDBK cells (Ozçelik et al., 2011). It induces antioxidant activity enhancing the synthesis of tryptophan and nicotinamide (Pero et al., 2009).

2H-1-Benzopyran, 7-methoxy-2,2-dimethyl- (syn. Precocene I) was identified from *R. lepidotum* (7.42%). It is a naturally occurring chromonoid (Bowers et al., 1976) known to display insect antijvenile hormonal activity (Bowers, 1985). It has been reported to exhibit insecticidal activities against *Drosophila melanogaster* (Wilson et al., 1983), *Aedes albopictus* (Liu & Liu, 2014), *Liposcelis bostrychophila* (Lu et al., 2014), and a malathion-resistant *Oryzaephilus surinamensis* (Saleem & Wilkins, 1984). It has also been known to inhibit mycotoxin 3-acetyldeoxynivalenol in *Fusarium graminearum* (Yaguchi et al., 2009).

Cyclopentane, 1-acetyl-1,2-epoxy- is a steroid that was identified from *L. involuta* (7.13%), and *C. aciculatus* (4.01%) in this study. It has been found as a precursor for cyclopentane monoterpene synthesis (Prabu et al., 2013).

cis-Vaccenic acid is a monounsaturated and non-essential ω -7 fatty acid which was identified from *T. dumosa* (6.97%), and *G. wallichianum* (2.25%) in this study. It has been reported to exhibit anti-inflammatory and antioxidant activities (Anyasor et al., 2014). It has been found

to inhibit the growth of *Bacillus subtilis* (Laser, 1951). It has been reported to suppress VCAM-1 and ICAM-1 expression (Abbasia et al., 2015) and exhibit hypolipidaemic and antihypertensive activities (Bhattacharya et al., 2014). It has been found to reduce the risk of myocardial infarction (Djousse et al., 2012) and heart failure with antecedent coronary heart disease (Djousse et al., 2014). However, it has been positively associated with reduced kidney function (Block et al., 2012).

Benzenemethanol, 2,5-dimethoxy-, acetate was identified from *S. lucida* (6.88%). It has been reported to possess antibacterial activity (Jebastella & Reginald, 2015).

Decanoic acid, octyl ester (Syn. octyl decanoate) was identified from *A. rivularis* (5.93%). It has also been reported from honey bees such as *Apis mellifera* queens (Bernasconi et al., 1999) and *Trigona silvestriana* workers (Johnson et al., 1985).

A sugar named β -D-Glucopyranose, 1,6-anhydro- (Syn. levoglucosan) is a pyrolysis product of cellulose and is one of the major organic components in the biomass combustion of particulate matter. It was identified from *P. emblica* (5.49%), *C. aciculatus* (4.53%), and *A. rivularis* (3.46%) in this study. It has been frequently used as an environmental tracer for biomass burning (Fraser & Lakshmanan, 2000; Simoneit et al., 1999).

2-Butenal, 2-methyl-, (E)- (Syn. tiglic aldehyde) was detected from *P. emblica* (5.47%). This hemiterpene molecule displays properties such as fuel and energy storage, cell signaling, and membrane integrity (HMDB, 2016; Yannai, 2004).

α -D-Glucopyranoside, O- α -D-glucopyranosyl-(1,6)- β -D-fructofuranosyl (Syn. Melezitose) was identified from *R. lepidotum* (5.27%). It is a trisaccharide molecule that has been obtained from plant exudations or honey made from such exudations, and also incorporated by some mushrooms (Barros et al., 2008).

1-[3-(1-Adamantyl)-1-methylpropylidene]thiosemicarbazide was found in *R. lepidotum* (5.04%). It has been reported to possess antimicrobial activity (Kumaravel et al., 2016).

γ -Sitosterol was identified from *S. elata* (4.8%), and *S. lucida* (3.15%). It has been suggested to exert hypolipidemic (Balamurugan et al., 2015), antihyperglycemic (Balamurugan et al., 2011), anti-diabetic, anti-angiogenic, anticancer, antimicrobial, anti-inflammatory, antidiarrhoeal and antiviral properties (Venkata et al., 2012). It shows anticancer activity against MCF-7 and A549 cells exhibiting growth inhibition, cell cycle arrest and apoptosis through c-Myc suppression (Sundarraaj et al., 2012).

D-Allose is a rare sugar that was identified from *T. dumosa* (4.56%), *G. wallichianum* (3.15%), *S. elata* (2.98%), and *R. scleratus* (0.77%) in this study. It has been reported to possess antioxidant (Murata et al., 2003), immunosuppressive (Hossain et al., 2000) and anti-inflammatory activities (Gao et al., 2011). It has also been reported to exhibit anticancer effects against human ovarian (Sui et al., 2005a), prostate (Naha et al., 2008), liver (Sui et al., 2005b), leukemia (Hirata et al., 2009), and head and neck (Mitani et al., 2009) cancer cell lines.

1,3-Propanediol, 2-(hydroxymethyl)-2-nitro- is a glycerol/nitrogen compound which was identified from *A. rivularis* (4.21%), and *S. elata* (1.2%) in this study. It has been reported to have anticataract, anti-tear-wax, antiketotic, antineuralgic, arrhythmogenic, emollient, hyperglycemic activities (Kumar et al., 2012). It exhibits antimicrobial activities (Gunasekarana et al., 2013), and thus used as a bacteriostat in disinfectants (Popendorf et al., 1995).

Benzaldehyde, 2-hydroxy-6-methyl- (Syn. 6-Methylsalicylaldehyde or 2,6-Cresotaldehyde) is an aromatic aldehyde detected from *L. involuta* (4%). It has been reported to exhibit antimicrobial (Rajeswari et al., 2013), anticancer, antimutagenic, antitumor and immunostimulant activities (Balasubramanian et al., 2014).

Sucrose is a non reducing sugar formed by combination of glucose and fructose. It was identified from *C. aciculatus* (3.97%). It has been reported that diets very high in sucrose decrease insulin sensitivity (Thresher et al., 2000), induce hypertriglyceridemia and hyperinsulinemia (Daly et al., 1997). Dietary sucrose has also been found as a promotor of colon tumor (Caderni et al., 1997; Caderni et al., 1996; Kristiansen et al., 1996; Luceri et al., 1996; Stamp et al., 1993).

3,7,11,15-Tetramethyl-2-hexadecen-1-ol was identified from *H. javanica* (3.96%), *R. scleratus* (2.43%), *A. rivularis* (1.01%), *R. lepidotum* (0.79%), *S. elata* (0.79%), *P. emblica* (0.75%), and *T. dumosa* (0.58%). It is a terpene alcohol which has been reported to possess antimicrobial, anti-inflammatory (Rajeswari et al., 2012), anticancer and diuretic effects (Gopinath et al., 2013).

1,2-Benzenediol (Syn. catechol) was identified from *L. involuta* (3.87%), and *R. lepidotum* (0.96%). It is a phenolic compound reported to possess antioxidant, fungicidal, antibacterial, pesticidal (Saravanan et al., 2014), antiseptic (Cowan, 1999), anti-complementary (Chen, 2011) and anti-inflammatory activities (Park & Lee, 2012). It has been reported to inhibit NO production, tyrosinase activity and melanin production (Park & Lee, 2012). It is a powerful

reducing agents with potential application against oxidative stress (Kalt & Cock, 2014). It shows anticancer activity against breast cancer (Manorenjitha et al., 2013).

1,2-Benzenediol, 3-methoxy- was identified from *G. wallichianum* (3.81%), and *R. lepidotum* (0.55%). It is an aromatic compound which has been reported to exhibit a potent anti-complementary activity (Chen, 2011).

9-Octadecynoic acid (Syn. Stearolic acid) is the acetylenic analogue oleic acid identified from *L. involuta* (3.81%), and *C. aciculatus* (1.62%). It has been implicated as a hypotensive compound (Teres et al., 2008). It has also been reported as a DNA binding agent which binds to DNA and weakly inhibits DNA polymerase activity (Berry et al., 1991).

Pentadecanoic acid is a lauric acid identified from *T. dumosa* (3.78%), *A. rivularis* (3.65%), *G. wallichianum* (3.48%), and *R. scleratus* 2.99%). It has been reported to have antioxidant activity (Lawal et al., 2015).

Phthalic acid, di-(1-hexen-5-yl) ester is an organic acid ester identified from *S. elata* (3.76%) which possess antifungal properties (Sridevi et al., 2014).

Octadecane, 2,2,4,15,17,17-hexamethyl-7,12-bis(3,5,5-trimethylhexyl)- was identified from *L. involuta* (3.75%). It has been reported to exert anticancer activities (Babar et al., 2016; Camerman et al., 1978; Corbett et al., 1982).

2-Methoxy-4-vinylphenol was identified from *C. aciculatus* (3.74%), *R. lepidotum* (0.91%), and *H. javanica* (0.33%). It is a phenolic compound used as a flavouring agent (Jeong et al., 2011) which has been reported to possess antimicrobial, antioxidant and analgesic effects (Gopalakrishnan & Vadivel, 2011; Rajeswari & Rani, 2014). It exhibits strong anti-inflammatory activity through the elimination of NF- κ B and MAPK activation (Jeong et al., 2011). It has been reported to arrest abnormal cell cycle progression by preventing the hyperphosphorylation of retinoblastoma protein in NIH 3T3 cells (Jeong & Jeong, 2010).

Stigmast-5-en-3-ol, (3 β)- (Syn. β -Sitosterol) was identified from *T. dumosa* (3.66%), and *G. wallichianum* (2.94%). It has been reported to exert phosphorylation of extracellular-signal regulating kinase and p38 mitogen-activated protein kinase (Moon et al., 2007) and promote apoptosis by increasing FAS levels and caspase-8 activity (Awad et al., 2007). It has been found to modulate antioxidant enzyme (Vivancos & Moreno, 2005), decrease free radical generation (Moreno, 2003), and inhibit colon cancer (Baskar et al., 2010; Jayaprakash et al., 2007) and prostate cancer (Awad et al., 2005). It exhibits analgesic, anti-inflammatory (Dighe et al., 2016) and cholesterol lowering effect (Wang & Ng, 1999). It has also been

reported to possess antimicrobial, antiarthritic, antiasthma, diuretic (Mujeeb et al., 2014), antidiabetic, antiangiogenic, antidiarrhoeal and antiviral activities (De Oliveira et al., 2014).

2-Deoxy-D-galactose was identified from *L. involuta* (3.41%). It has been reported as a substrate for the Galactose Transport System of *Escherichia coli* (Henderson & Giddens, 1977) which has been found to inhibit tumors in mice (Laszlo et al., 1960) and prevent accumulation of toxic phosphorylated metabolites of 2-deoxygalactose (Nagelkerke & Postma, 1978).

9,12,15-Octadecatrienoic acid, (Z,Z,Z)- is an unsaturated fatty acid identified from *R. scleratus* (3.08%), and *A. rivularis* (1.64%). It has been reported to exhibit antimicrobial, antiallopecic, antianaphylactic, antiarteriosclerotic, antileukotriene, antimenorrhagic, antiprosthetic, carcinogenic, antigranular, metastatic, immunomodulatory (Mickymaray et al., 2016) as well as nematicide and antieczemic activities (Kavitha et al., 2014). It has also been reported to display antiandrogenic (5-Alpha-reductase-inhibitory), antiacne, antiarthritic, anticoronary, antihistaminic, antiinflammatory, cancer preventive, hepatoprotective, hypocholesterolemic and insectifuge activities (Kavitha et al., 2014; Mickymaray et al., 2016). It has also been reported to be applicable as drugs for genital or sexual disorders and in contraceptives (Rawal & Sonawani, 2016).

Cholest-5-en-3-ol, 24-propylidene-, (3 β) (Syn. 24-Propylidenecholest-5-en-3 β -ol) was identified from *S. lucida* (3.06%). It has been reported to exhibit antibacterial and antioxidant activities (Thanigaivel et al., 2015).

9,12-Octadecadienoic acid (Z,Z)-, 2,3-dihydroxypropyl ester was identified from *R. scleratus* (3.04%). It has been reported to exhibit antipyretic, anticonvulsant, antiseptic, analgesic (Edewor et al., 2016), hypocholesterolemic, antieczemic, nematicide, hepatoprotective, antiarthritic, antiacne and antihistaminic activities (Gnanavel & Saral, 2013).

Maytansine was identified from *L. involuta* (2.86%), and *C. aciculatus* (1.34%). It is an ansa macrolide structure attached to a chlorinated benzene ring which has been found to suppress microtubule dynamics (Lopus et al., 2010) and exhibit antimitotic effect (Bhattacharyya & Wolff, 1977; Mandelbaum-Shavit et al., 1976; Remillard et al., 1975). It has been reported to be effective against breast cancer, lung carcinoma and murine melanocarcinoma solid tumors and has an antileukemic activity against lymphocytic leukemia (Cassady et al., 2004; Lopus et al., 2009). It inhibits microtubule assembly and kills cancer cells, but has severe side effects (Cassady et al., 2004). It shows systemic toxicity such as neurotoxicity and gastrointestinal tract adverse reaction (Haddley, 2013; They et al., 2014).

Phytol was identified from *H. javanica* (2.75%), *L. involuta* (2.68%), *A. rivularis* (3.19%), *R. scleratus* (1.27%), *S. elata* (0.52%), *C. aciculatus* (0.44%), and *R. lepidotum* (0.41%). It is an acyclic monounsaturated diterpene alcohol known to be the precursor for synthetic forms of vitamin E (Qiu et al., 2014) and vitamin K1 (Daines et al., 2003). It has been found in chlorophyll and tocopherols and has been known to be a strong immunomodulator but it exerts strong toxicity (Kagoura & Morohashi, 1999; Lim et al., 2006). It has been reported to exhibit antischistosomal (de Moraes et al., 2014), antinociceptive, antioxidant (Santos et al., 2013), antimicrobial, diuretic (Jananie et al., 2011; Sudha et al., 2013), antiinflammatory (Silva et al., 2014), antiallergic (Ryu et al., 2011) and anticancer activities (Guo et al., 2014; Islam et al., 2015; Kim et al., 2015; Pejin et al., 2014).

We identified a number of bioactive compounds with proven bioactivities. The presence of these and other bioactive compounds either alone or in combination are responsible for medicinal properties of these plants. Further isolation and elucidation of safe therapeutic phytochemicals may help address the problem of drug resistance and toxicity. Novel phytopharmaceuticals can bring infinite possibilities in modern medicine.

6 Summary and Conclusion

Preliminary study was conducted on bioactive compounds and biological activities of twelve medicinal plants with some history of use as traditional medicine around the world. The results from total phenolic content, total flavonoid content, GC-MS analysis, radical scavenging activity, antimicrobial activity, brine shrimp lethality, and cancer cell inhibition advocated that the bioactive compounds afforded by plants display one or more of the biological activities either individually or synergistically.

The methanolic extracts of *T. dumosa*, *P. emblica*, *G. wallichianum*, *T. bellerica*, *S. elata*, *S. lucida*, and *R. lepidotum* showed a good antioxidant activity with IC_{50} value less than 50 $\mu\text{g/ml}$. Phenolic compounds were attributed to be the major contributor of antioxidant activity as suggested by a significant correlation between TPC and IC_{50} of DPPH RSA. These plants and probably others could be the good source of natural antioxidants. The higher antimicrobial activities against *S. aureus* and MRSA both were displayed by *T. bellerica*, *P. emblica*, *R. lepidotum*, *T. dumosa*, *G. wallichianum*, and *S. elata*. *S. Typhi* was inhibited maximally by *T. dumosa* followed by *S. elata*, *C. aciculatus*, and *T. bellerica*. Minimum number of plants showed activity against *K. pneumoniae* with higher effect exerted by *L. involuta* followed by *C. aciculatus*. Ten of the twelve plant extracts inhibited at least three of the six organisms tested while eight extracts inhibited at least four organisms. *G. wallichianum* demonstrated effectiveness against five of the organisms tested except *K. pneumoniae*. *S. elata* and *T. bellerica* were active against all the tested microorganisms. Plants inhibited gram positive bacteria more effectively. Nine of the twelve medicinal plants showed significant toxicity to brine shrimps with LC_{50} value less than 500 $\mu\text{g/ml}$ including the three viz. *A. rivularis*, *L. involuta*, and *T. bellerica* showing high toxicity with LC_{50} less than 100 $\mu\text{g/ml}$. Maximum growth inhibition on HeLa was shown by *S. lucida* followed by *T. dumosa*, *C. aciculatus*, *R. scleratus*, *S. elata*, *A. rivularis*, *R. lepidotum*, and *L. involuta* while that on MDCK was exerted by *L. involuta* followed by *T. dumosa*, and *A. rivularis* with EC_{50} value less than 500 $\mu\text{g/ml}$. Higher number of plants were effective against HeLa as compared to MDCK. The major bioactive compounds identified from GC-MS analysis of plant extracts have been found to have previously been reported to exert different biological activities including antioxidant, antimicrobial, and anticancer.

Multidrug resistant microbial pathogens are on rise. Cancer treatment still remains a big challenge. The lack of efficient treatment in addition to the side effects resulting from current therapies need to be effectively addressed. At this stage, our study offers a scientific basis for traditional use of medicinal plants. Plants showed the rich presence of bioactive compounds.

The significant inhibition of *S. aureus* and MRSA, multidrug resistant *S. Typhi* and *K. pneumoniae* by some of the plant extracts suggest their use in isolation of compounds with antimicrobial activity and as the substitute for antibiotics. The significant toxicity of some of the medicinal plants signifies their potential use as the source of anticancer drugs and at the same time demands their careful analysis before use in medical formulations. Hence, the medicinal plants could be used as an important source of therapeutic components in pharmaceutical and medical industries. For this reason, further work need to be conducted to isolate, identify and characterize bioactive compounds and test them rigorously *in vitro* and *in vivo* and assess their safety which helps discover novel drugs.

7 Recommendation

Further investigations are required for:

- isolation, identification, and characterization of the bioactive compounds from different solvent fractions
- observation of biological activities of isolated compounds either alone or in combination with other compounds *in vivo* and *in vitro*
- examination of therapeutic mechanisms of plant extracts and their compounds with the focus to explore their specific cellular and molecular targets

References

- Abbasi MA (2004) Bioactive chemical constituents of *Symplocos recemosa* and *Comiphora mukul* (pp. 139). Karachi, Pakistan: Thesis Submitted in University of Karachi.
- Abbasia A, Mostafaiea A, Bahramia G, Mansouria K, & Sisakhtnejada S (2015) The effect of *cis* and *trans* vaccenic acids on expression of ICAM-1 And VCAM-1 in human microvascular endothelial cells (HMEC). *Journal of Reports in Pharmaceutical Sciences*. **4**(1): 65-74
- Acharya R, & Acharya KP (2009) Ethnobotanical study of medicinal plants used by Tharu community of Parroha VDC, Rupandehi District, Nepal. *Scientific World*. **7**(9): 80-84
- Ahmad B, Ismail M, Iqbal Z, & Chaudhry MI (2003) Biological Activities of *Geranium wallichianum*. *Asian Journal of Plant Sciences*. **2**(13): 971-973
- Ahmad K, Shakeel-u-Rehman, Chisti AM, Shawl AS, & Taneja SC (2010) Chemical constituents of *Rhododendron lepidotum*. *Chemistry of Natural Compounds*. **46**(2): 195-197
- Ahmad VU, Zubair M, Abbasi MA, Kousar F, Ullah F, Fatima N, & Choudhary MI (2005) Phenolic glycosides from *Symplocos racemosa*. *Zeitschrift für Naturforschung*. **60b**: 1101-1104
- Ahmed Y, Sohrab H, Al-Reza SM, Tareq S, F, Hasan CM, & Sattar MA (2010) Antimicrobial and cytotoxic constituents from leaves of *Sapium baccatum*. *Food and Chemical Toxicology*. **48**(2): 549-552
- Ajani EO, Salako AA, Sharlie PD, Akinleye WA, Adeoye AO, Salau BA, & Adebawo OO (2009) Chemopreventive and remediation effect of *Hydrocotyl bonariensis* Comm. Ex Lam (Apiaceae) leave extract in galactose-induced cataract. *Journal of Ethnopharmacology*. **123**(1): 134-142
- Aleksandrova MS (1975) *Rhododendrons of USSR Natural Flora*. Moscow: Nauka.
- Ali M, Bhutani KK, & Srivastava TN (1990) Triterpenoids from *Symplocos racemosa* bark. *Phytochemistry*. **29**(11): 3601-3604
- Almeida PA, Silva TMS, & Echevarria A (2002) Mesoionic 5-alkyl-1,3-dithiolium-4-thiolates: synthesis and brine shrimp toxicity. *Heterocyclic Communications*. **8**(6): 593-600
- Alves RRN, & Rosa IL (2007) Biodiversity, traditional medicine and public health: where do they meet? *Journal of Ethnobiology and Ethnomedicine*. **3**(14): 1–9
- Ambasta SP, Ramchandran K, Kashyapa K, & Chand R (1992) *The useful plant of India*. New Delhi, India: Council of Science and Industrial Research (CSIR).
- Amel OH, Malek BH, Hichem BJ, Ali L, Mahjoub A, & Boulbaba S (2013) Antioxidant and anti-acetylcholinesterase activities of extracts from *Rapistrum rugosum* in Tunisia. *Asian Pacific Journal of Tropical Disease*. **3**(5): 367–374
- Andreia A, Rosatella A, Svilen P, Simeonov A, Rraquel FM, Frade A, & Carlos AMA (2011) 5-Hydroxymethylfurfural (HMF) as a building block platform: Biological properties, synthesis and synthetic applications. *Green Chemistry*. **13**: 754-793
- Anyasor GN, Aina DA, Olushola M, & Aniyikaye AF (2011) Phytochemical constituent, proximate analysis, antioxidant, antibacterial and wound healing properties of leaf extracts of *Chromolaena odorata*. *Annals of Biological Research*. **2**(2): 441-451

- Anyasor GN, Funmilayo O, Odutola O, Olugbenga A, & Oboutor EM (2014) Chemical constituents in n-butanol fractions of *Costus afer* Ker Gawl leaf and stem. *Journal of Intercultural Ethnopharmacology*. **3**(2): 78-84
- Anyasor GN, Ogunwenmo KO, Oyelana OA, & Akpofunure BE (2010) Phytochemical constituents and antioxidant activities of aqueous and methanol stem extracts of *Costus afer* Ker Gawl. (Costaceae). *African Journal of Biotechnology*. **9**(31): 4880-4884
- Arslanyolu M, & Erdemgil FZ (2006) Evaluation of the antibacterial activity and toxicity of isolated arctiin from the seeds of *Centaurea sclerolepis*. *Journal Faculty of Pharmacy Ankara*. **35**(2): 103-109
- Arteel GE (2003) Oxidants and antioxidants in alcohol induced liver disease. *Gastroenterology*. **124**(3): 778–790
- Arulmozhi A, & Surya C (2016) Antidiabetic and hypolipidemic activity of *Terminalia bellirica* on alloxan induced diabetic rats. *Asian Journal of Innovative Research*. **1**(1): 16-21
- Aslokar LV, Kakkar KK, & Chakre OJ (1992) *Glossary of Indian Medicinal Plants with active principles. Part I*. New Delhi: CSIR.
- Athiroh N, Permatasari N, Sargowo D, & Widodo MA (2014) Effect of *Scurrula atropurpurea* on nitric oxide, endothelial damage, and endothelial progenitor cells of DOCA-salt hypertensive rats. *Iranian Journal of Basic Medical Sciences*. **17**(8): 622-625
- Atta-ur-Rahman, Choudhary MI, & Thomsen JW (1991) *Manual of bioassay techniques for natural product research*. Amsterdam: Harward academic press.
- Awad AB, Andrew TB, & Carol SF (2005) Effect of resveratrol and β -sitosterol in combination on reactive oxygen species and prostaglandin release by PC-3 cells. *Prostaglandins, Leukotrienes and Essential Fatty Acids*. **72**(3): 219-226
- Awad AB, Chinnam M, Fink CS, & Bradford PG (2007) β -sitosterol activates Fas signaling in human breast cancer cells. *Phytomedicine*. **14**(11): 747-754
- Ayensu ES, & DeFilipps RA (1978) *Endangered and Threatened Plants of the United States*. Washington, DC: Smithsonian Institution Press and World Wildlife Fund.
- Ayoola GA, Coker HAB, Adesegun SA, Adepoju-Bello AA, Obaweya K, Ezennia EC, & Atangbayila TO (2008) Phytochemical screening and antioxidant activities of some selected medicinal plants used for malaria therapy in Southwestern Nigeria. *Tropical Journal of Pharmaceutical Research*. **7**(3): 1019-1024
- Babar AG, Pande A, & Kulkarni BG (2016) Antifungal activity and investigation of bioactive compounds of marine intertidal bivalve *Gafrarium divaricatum* from West coast of India. *International Journal of Pure & Applied Bioscience*. **4**(2): 211-217
- Balamurugan R, Duraipandiyar V, & Ignacimuthu S (2011) Antidiabetic activity of γ -sitosterol isolated from *Lippia nodiflora* L. in streptozotocin induced diabetic rats. *European Journal of Pharmacology*. **667**(1-3): 410-418
- Balamurugan R, Stalin A, Aravinthan A, & Kim J-H (2015) γ -sitosterol a potent hypolipidemic agent: *In silico* docking analysis. *Medicinal Chemistry Research*. **24**(1): 124-130
- Balasubramanian S, Ganesh D, Kiran KS, Prakash KJM, & Surya NVVS (2014) GC-MS analysis of phytocomponents in the methanolic extract of *Mentha arvensis* (corn mint). *International Journal of Chemistry and Pharmaceutical Sciences*. **2**(6): 926-929

- Baliga MS, & Dsouza JJ (2011) Amla (*Embllica officinalis* Gaertn), a wonder berry in the treatment and prevention of cancer. *European Journal of Cancer Prevention*. **20**(3): 225–239
- Baral SR, & Kurmi PP (2006) *Compendium of Medicinal Plants in Nepal*. Chabahil, Kathmandu, Nepal: Rachana Sharma publishers.
- Barile E, Bonanomi G, Antignani V, Zolfaghari B, Sajjadi SE, Scala F, & V L (2007) Saponins from *Allium minutiflorum* with antifungal activity. *Phytochemistry*. **68**(5): 596-603
- Barlow B (1991) Privisional key to the genera of Loranthaceae and Viscaceae of the Flora Malesiana region. *Flora Malesiana Bulletin*. **10**(4): 335-338
- Barros L, Cruz T, Baptista P, Estevinho LM, & Ferreira IC (2008) Wild and commercial mushrooms as source of nutrients and nutraceuticals. *Food and Chemical Toxicology*. **46**(8): 2742-2747
- Barukial J, & Sarmah JN (2011) Ethnomedicinal plants used by the people of Golaghat district, Assam, India. *International Journal of Medicinal and Aromatic Plants*. **1**(3): 203-211
- Baskar AA, Ignacimuthu S, Paulraj GM, & Al-Numair KS (2010) Chemopreventive potential of β -Sitosterol in experimental colon cancer model - an *In vitro* and *In vivo* study. *BMC Complementary and Alternative Medicine*. **10**(1): 1-10
- Becker EM, Nissen LR, & Skibsted LH (2004) Antioxidant evaluation protocols: Food quality or health effects. *European Food Research and Technology*. **219**(6): 561-571
- Benvenuti S, Pellati F, Melegar M, & Bertelli D (2004) Polyphenols, anthocyanins, ascorbic acid and radical scavenging activity of Rubus, Ribes and Aronia. *Journal of Food Science*. **69**: 164-169
- Bernasconi G, Bigler L, Hesse M, & Ratnieks FLW (1999) Characterization of queen-specific components of the fluid released by fighting honey bee queens. *Chemoecology*. **9**(4): 161-167
- Berry DE, Chan JA, MacKenzie L, & Hecht SM (1991) 9-Octadecynoic acid: a novel DNA binding agent. *Chemical Research in Toxicology*. **4**(2): 195–198
- Bhargava KP, Kishor K, Pant MC, & Saxena PR (1965) Identification of tryptamine derivatives in *Ranunculus sceleratus* L. *British Journal of Pharmacology*. **25**(3): 743-750
- Bhattacharya A, Ghosh G, Agrawal D, Sahu PK, Kumar S, & Mishra SS (2014) GC-MS profiling of ethanolic extract of *Moringa oleifera* leaf. *Int J Pharm Bio Sci*. **5**(4): 263 - 275
- Bhattacharyya A, LaMarche VC, & Hughes MK (1992) Tree-ring chronologies from Nepal. *Tree-Ring Bulletin*. **52**: 59-66
- Bhattacharyya B, & Wolff J (1977) Maytansine binding to the vinblastine sites of tubulin. *FEBS Letters*. **75**(1): 159–162
- Bhattarai K, Shrestha TM, Bajracharya R, Jain SC, & Lamichhane J (2010) Biological activities of three different medicinal plants from Himalayan region of Nepal. *Nepal Journal of Science and Technology*. **11**: 139-146
- Bhuvan RP, Jignesh PD, Bhavik PA, & Ashok GL (2009) Potent *in vitro* anticancer activity of *Symplocos racemosa* bark. *Romanian Journal of Biology-Plant Biology*. **54**(2): 135-140
- Birt DF, Hendrich S, & Wang W (2001) Dietary agents in cancer prevention: flavonoids and isoflavonoids. *Pharmacology & Therapeutics*. **90**(2-3): 157–177
- Blinova KF, Karpovich VN, & Sukkei LK (1961) *Proceedings of the 2nd All-Union Conference on Pharmaceuticals, Moscow*.

- Block R, Kakinami L, Liebman S, Shearer GC, Kramer H, & Tsai M (2012) *Cis*-vaccenic acid and the Framingham Risk Score predict chronic kidney disease: the multi-ethnic study of atherosclerosis (MESA). *Prostaglandins, Leukotrienes and Essential Fatty Acids*. **86**(4-5): 175-182
- Bolton JL, Trush MA, Penning TM, Dryhurst G, & Monks TJ (2000) Role of quinones in toxicology. *Chemical Research in Toxicology*. **13**(3): 135–160
- Borowski E, Bontemps-Gracz MM, & Piwkowska A (2005) Strategies for overcoming ABC-transporters-mediated multidrug resistance (MDR) of tumor cells. *Acta Biochimica Polonica*. **52**(3): 609–627
- Borris RP (1996) Natural products research: perspectives from a major pharmaceutical company. *Journal of Ethnopharmacology*. **51**(1-3): 29-38
- Bouayed J, Khosro PB, Rammal H, Dicko A, Desor F, Younos C, & Soulimani R (2007) Comparative evaluation of the antioxidant potential of some Iranian medicinal plants. *Food Chemistry*. **104**(1): 364-368
- Bowers WS (1985) Antihormones. In Kerkut GA & Gilbert LI (Eds.), *Comprehensive Insect Physiology, Biochemistry and Pharmacology* (Vol. 8, pp. 551-564). New York: Pergamon Press.
- Bowers WS, Ohta T, Cleere JS, & Marsella PA (1976) Discovery of insect anti juvenile hormones in plants. *Science*. **193**(4253): 542–547
- Bravo L (1998) Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. *Nutrition Reviews*. **56**(11): 317-333
- Brustugun J, Tonnesen HH, Edge R, & Navaratnam S (2005) Formation and reactivity of free radicals in 5-hydroxymethyl-2-furaldehyde—the effect on isoprenaline photostability. *Journal of Photochemistry and Photobiology B: Biology*. **79**(2): 109–119
- Burt S (2004) Essential oils: their antibacterial properties and potential applications in foods—a review. *International Journal of Food Microbiology*. **94**(3): 233-253
- Caderni G, Lancioni L, Luceri C, Giannini A, Lodovici M, Biggeri A, & Dolara P (1997) Dietary sucrose and starch affect dysplastic characteristics in carcinogen-induced aberrant crypt foci in rat colon. *Cancer Letters*. **114**(1-2): 39-41
- Caderni G, Luceri C, Lancioni L, & Dolara P (1996) Dietary sucrose, glucose, fructose, and starches affect colonic functions in rats. *Nutrition and Cancer*. **25**(2): 179–186
- Calvin CL, & Wilson CA (2006) Comparative morphology of epicortical roots in Old and New World Loranthaceae with reference to root types, origin, patterns of longitudinal extension and potential for clonal growth. *Flora - Morphology, Distribution, Functional Ecology of Plants*. **201**(1): 51–64
- Camerman A, Smith HW, & Camerman N (1978) Stereochemistry of dihydrofolate reductase inhibitor antitumor agents: molecular structure of "Baker's antifol" (NSC 139105, triazinate) and "insoluble Baker's antifol" (NSC 113423). *Biochemical and Biophysical Research Communications*. **83**(1): 87-93
- Campbell DL, Lawton LA, Beattie KA, & Codd GA (1994) Comparative assessment of the specificity of the brine shrimp and microtox assays to hepatotoxic (microcystin-LR-containing) cyanobacteria. *Environmental Toxicology*. **9**(1): 71-77
- Cao G, Sofic E, & Prior RL (1997) Antioxidant and prooxidant behavior of flavonoids: structure-activity relationships. *Free Radical Biology and Medicine*. **22**(5): 749–760

- Carballo JL, Hernández-Inda ZL, Pérez P, & García-Grávalos MD (2002) A comparison between two brine shrimp assays to detect *in vitro* cytotoxicity in marine natural products. *BMC Biotechnology*. **2**: 17
- Carte' BK, Carr S, DeBrosse C, Hemling ME, MacKenzie L, Offen P, & Berry DE (1991) Aciculatin, a novel flavone-C-glycoside with DNA binding activity from *Chrysopogon aciculatis*. *Tetrahedron*. **47**(10-11): 1815–1822
- Carter S, & Livingston R (1982) Principle of cancer chemotherapy. In Carter S, Glatstein E & Livingston R (Eds.), *Principles of Cancer Treatment* (pp. 95–110). New York, USA: McGraw-Hill.
- Cassady JM, Chan KK, Floss HG, & Leistner E (2004) Recent developments in the maytansinoid antitumor agents. *Chemical & Pharmaceutical Bulletin (Tokyo)*. **52**(1): 1–26
- CCTHD (1992) *Compendium of Chinese Traditional Herbal Drugs* (Vol. 2). Beijing: People's Health Press.
- Çelik TA, & Aslantürk ÖS (2007) Cytotoxic and genotoxic effects of *Lavandula stoechas* aqueous extracts. *Biologia*. **62**(3): 292–296
- Chang YS, Chen IS, Hsieh WC, & Ou JC (2003) *The catalogue of medicinal plant resources in Taiwan*. Taipei, Taiwan: Committee on Chinese Medicine and Pharmacy, Department of Health, Executive Yuan.
- Chao L, Hsiu S-L, & Hou Y-C (2002) Flavonoids in herbs: biological fates and potential interactions with xenobiotics. *Journal of Food and Drug Analysis*. **10**(4): 219-228
- Charoenteeraboon J, Ngamkitidechakul C, Soonthornchareonnon N, Jaijoy K, & Sireeratawong S (2010) Antioxidant activities of the standardized water extract from fruit of *Phyllanthus emblica* Linn. *Songklanakarinn Journal of Science and Technology*. **32**(6): 599-604
- Chen CW, & Ho CT (1995) Antioxidant properties of polyphenols extracted from green and black teas. *Journal of Food Lipids*. **2**(1): 35-46
- Chen DF (2011) Isolation and characterization of anticomplementary agents from medicinal plants. *Planta Medica*. **77**(5): S9
- Chew YL, Lim YY, Stanslas J, Ee GC, & Goh JK (2014) Bioactivity-guided isolation of anticancer agents from *Bauhinia kockiana* Korth. *African Journal of Traditional, Complementary and Alternative medicines*. **11**(3): 291-299
- Chin YW, Balunas MJ, Chai HB, & Kinghorn AD (2006) Drug discovery from natural sources. *AAPS Journal*. **8**(2): E239-253
- Chitme HR, Chandra M, & Kaushik S (2004) Studies on anti-diarrhoeal activity of *Calotropis gigantea* R.Br. in experimental animals. *Journal of Pharmacy and Pharmaceutical Sciences*. **7**(1): 70-75
- Chopra RN, Chopra IC, Handa KL, & Kapur LD (1958) *Indigenous drugs of India*. Calcutta: Academic Publishers.
- Chopra RN, Nayar SL, & Chopra IC (1956) *Glossary of Indian medicinal plants*. New Delhi: Council of Scientific & Industrial Research.
- Chosson E, Chaboud A, Chulia JA, & Raynaud J (1998a) Dihydroflavonol glycosides from *Rhododendron ferrugineum*. *Phytochemistry*. **49**(5): 1431-1433
- Chosson E, Chaboud A, Chulia JA, & Raynaud J (1998b) A phloracetophenone glucoside from *Rhododendron ferrugineum*. *Phytochemistry*. **47**(1): 87-88

- Choudhary GP (2008) Wound healing activity of the ethanol extract of *Terminalia bellirica* Roxb. fruits. *Natural Product Radiance*. **7**(1): 19-21
- Chuang DY, Chan MH, Zong Y, Sheng W, He Y, Jiang JH, Simonyi A, Gu Z, Fritsche KL, Cui J, Lee JC, Folk WR, Lubahn DB, Sun AY, & Sun GY (2013) Magnolia polyphenols attenuate oxidative and inflammatory responses in neurons and microglial cells. *Journal of Neuroinflammation*. **10**: 15
- Clarkson C, Maharaj VJ, Crouch NR, Grace OM, Pillay P, Matsabisa MG, Bhagwandin N, Smith PJ, & Folb PI (2004) *In vitro* antiplasmodial activity of medicinal plants native to or naturalised in South Africa. *Journal of Ethnopharmacology*. **92**(2-3): 177-191
- Collins AR (2005) Assays for oxidative stress and antioxidant status: applications to research into the biological effectiveness of polyphenols. *American Journal of Clinical Nutrition*. **81**(1): 261S-267S
- Conner DE, & Beuchat LR (1984) Effects of essential oils from plants on growth of food spoilage yeasts. *Journal of Food Science*. **49**(2): 429-434
- Cohon RC, Cramp MC, & Parsons PJ (1980). *Journal of the Chemical Society, Chemical Communications*: 197
- Corbett TH, Leopold WR, Dykes DJ, Roberts BJ, Griswold DPJ, & Schabel FMJ (1982) Toxicity and anticancer activity of a new triazine antifolate (NSC 127755). *Cancer research*. **42**(5): 1707-1715
- Council NR (1982) Ecological aspect of the development in the humid tropics. Washington, DC: National Academy Press.
- Cowan MM (1999) Plant products as antimicrobial agents. *Clinical Microbiology Reviews*. **12**(4): 564-582
- Cronquist A (1981) *An integrated system of classification of flowering plants*. New York: Columbia University Press.
- CSIR (1968) *The wealth of India: A dictionary of Indian raw materials and industrial products* (Vol. IV). New Delhi, India: National Institute of Science Communication.
- Currell G (2015) *Scientific Data Analysis*. Oxford: Oxford University Press.
- Daines AM, Payne RJ, Humphries ME, & Abell AD (2003) The synthesis of naturally occurring vitamin K and vitamin K analogues. *Current Organic Chemistry*. **7**(16): 1625-1634
- Daly ME, Vale C, Walker M, Alberti KG, & Mathers JC (1997) Dietary carbohydrates and insulin sensitivity: a review of the evidence and clinical implications. *American Journal of Clinical Nutrition*. **66**(5): 1072–1085
- Dani DD (1986) Population and society in Nepal: an overview. In Joshi SC (Ed.), *Nepal Himalaya: geo-ecological perspective* (pp. 163-173). Nainital, India: Himalayan Research Group.
- Das MS, & Devi G (2015) *In vitro* cytotoxicity and glucose uptake activity of fruits of *Terminalia bellirica* in Vero, L-6 and 3T3 cell lines. *Journal of Applied Pharmaceutical Science*. **5**(12): 92-95
- Davidson-Hunt IJ (2000) Ecological ethnobotany: stumbling toward new practices and paradigms. *MASA*. **16**: 1–13
- Davies J (1994) Inactivation of antibiotics and the dissemination of resistance genes. *Science*. **264**(5157): 375-382

- Davies KJ (1995) Oxidative stress: the paradox of aerobic life. *Biochemical Society Symposia*. **61**: 1-31
- Davies KJ (2000) Oxidative stress, antioxidant defenses, and damage removal, repair, and replacement systems. *IUBMB Life*. **50**(4-5): 279-289
- de Moraes J, de Oliveira RN, Costa JP, Junior ALG, de Sousa DP, Freitas RM, Allegretti SM, & Pinto PLS (2014) Phytol, a diterpene alcohol from chlorophyll, as a drug against neglected tropical disease *Schistosomiasis mansoni*. *PLoS Neglected Tropical Diseases*. **8**(1): e2617
- De Oliveira GG, De S Pereira Junior JA, Bastos IVGA, Filho RCS, Lopes NP, & De Melo SJ (2014) Phytochemical Investigation of chloroform, dichloromethane and ethyl acetate extracts from the flowers of *Adenocalymma imperatoris-maximiliani* (Bignoniaceae). *International Journal of Pharma and Bio Sciences*. **5**(3): 491-500
- Desai JS, Chaudhari S, Adam A, & Chaudhari H (2014) Effect of alcoholic extract of *Terminalia bellirica* Roxb. in dextran sulphate sodium induced colitis. *World Journal of Pharmacy and Pharmaceutical Sciences*. **3**(3): 2121-2133
- Devmurari VP (2010) Antibacterial evaluation and phytochemical screening of *Symplocos racemosa* Roxb. *International Journal of PharmTech Research*. **2**(2): 1359-1363
- Dhakal RC, Kalauni SK, Awale S, Shrestha K, & Gewali MB (2011) Isolation and characterization of sterols and an aliphatic alcohol from *Tsuga dumosa* D. Don of Nepal. *Scientific World*. **9**(9): 16-17
- Dhaon R, Jain GK, Sarin JPS, & Khanna NM (1989) Symposide: a new antifibrinolytic glycoside from *Symplocos racemosa*. *Indian Journal of Chemistry*. **28**(2): 982-983
- Diallo D, Hveem B, Mahmoud MA, Betge G, Paulsen BS, & Maiga A (1999) An ethnobotanical survey of herbal drugs of Gourma district, Mali. *Pharmaceutical Biology*. **37**(1): 80-91
- Diaz JHJ, & Thilaga RD (2016) GC-MS analysis of methanolic extract from the internal shell of cephalopods. *World Journal of Pharmacy and Pharmaceutical Sciences*. **5**(1): 820-831
- Dighe SB, Kuchekar BS, & Wankhede SB (2016) Analgesic and anti-inflammatory activity of β -sitosterol isolated from leaves of *Oxalis corniculata*. *International Journal of Pharmacological Research*. **6**(3): 109-113
- Dileep P, & Nair GG (2015) Taxonomic and ethnobotanical studies of grasses used by tribals of Wayanad district, Kerala, south western ghats of India. *Journal of Global Biosciences*. **4**(5): 2212-2235
- Ding Y, Tang H-F, Wang J-B, Liu D, Tian X-R, Wang X-Y, & Zhou X-M (2011) Triterpenoid saponins from *Anemone rivularis* var. *flore-minore*. *Biochemical Systematics and Ecology*. **39**(3): 236-239
- Ding Y, Tian X-R, Tang H-F, Feng J-T, Zhang W, Hai W-L, Wang X-Y, & Wang Y (2012) Two new glycosides from the whole plant of *Anemone rivularis* var. *flore-minore*. *Phytochemistry Letters*. **5**(3): 668-672
- Djeridane A, Yousfi M, Nadjemi B, Boutassouna D, Stocker P, & Vidal N (2006) Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. *Food Chemistry*. **97**(4): 654-660
- Djoussé L, Matsumoto C, Hanson NQ, Weir NL, Tsai MY, & Gaziano JM (2014) Plasma *cis*-vaccenic acid and risk of heart failure with antecedent coronary heart disease in male physicians. *Clinical Nutrition*. **33**(3): 478-482

- Djoussé L, Matthan NR, Lichtenstein AH, & Gaziano JM (2012) Red blood cell membrane concentration of *cis*-palmitoleic and *cis*-vaccenic acids and risk of coronary heart disease. *American Journal of Cardiology*. **110**(4): 539–544
- Domb AJ, & Wolnerman JS (2002). *Eur. Pat. Appl. EP*. **137**: 1236-1244
- Dragsted LO, Strube M, & Larsen JC (1993) Cancer-protective factors in fruits and vegetables: biochemical and biological background. *Pharmacology & Toxicology*. **72**(1): 116–135
- Drews J (2000) Drug discovery: a historical perspective. *Science*. **287**(5460): 1960-1964
- Duke JA, Bogenschutz-Godwin MJ, duCellier J, & K DP-A (2002) *Handbook of Medicinal Herbs*. Boca Raton, Florida CRC Press.
- Duwadee NPS, & Kunwar RM (2001) Botanical survey of Khaptad National Park and buffer zone area, far western Nepal. *Botanica orientalis*. **2**: 165-170
- Edewor TI, Owa SO, Ologan AO, & Akinfemi F (2016) Quantitative determination of the saponin content and GC-MS study of the medicinal plant *Cassytha filiformis* (linn.) leaves. *Journal of Coastal Life Medicine*. **4**(2): 154-156
- Eduardo V, de Dios MJ, Isidro H, Analía D, & Christian S (2006) Antimicrobial activity of dichloromethane extracts of eleven plants from the flora of Entre Ríos (Argentina). *Pharmacologyonline*. **3**: 845-849
- El-Desouky SK, Ryu SY, & Kim YK (2008) A new cytotoxic acylated apigenin glucoside from *Phyllanthus emblica* L. *Natural Product Research*. **22**(1): 91–95
- Epand RF, Savage PB, & Epand RM (2007) Bacterial lipid composition and the antimicrobial efficacy of cationic steroid compounds (Ceragenins). *Biochimica et Biophysica Acta*. **1768**(10): 2500-2509
- Fan CQ, Zhao WM, Ding BY, & Qin WG (2001) Constituents from the leaves of *Rhododendron latoucheae*. *Fitoterapia*. **72**(4): 449-452
- Fang JM, Wu SK, & Cheng YS (1985) A study of the constituents of the bark of *Tsuga chinensis* Pritz. var. *formosana* (Hay.). *Journal of the Chinese Chemical Society*. **32**: 477–480
- Farnsworth NR, & Soejarto DD (2009) Global importance of medicinal plants. In Akerele O, Heywood V & Syngé H (Eds.), *Conservation of medicinal plants* (pp. 25-52). Cambridge: Cambridge University Press.
- Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, & Bray F (2015) Cancer incidence and mortality worldwide: sources, methods and major patterns in Globocan 2012. *International Journal of Cancer*. **136**(5): E359-386
- Finkel T, & Holbrook NJ (2000) Oxidants, oxidative stress and the biology of ageing. *Nature*. **408**(6809): 239–247
- Florentino IF, Nascimento MV, Galdino PM, De Brito AF, Da Rocha FF, Tonussi CR, De Lima TC, De Paula JR, & Costa EA (2013) Evaluation of analgesic and anti-inflammatory activities of *Hydrocotyle umbellata* L., Araliaceae (acariçoba) in mice. *Anais da Academia Brasileira de Ciências*. **85**(3): 987-997
- Fotakis G, & Timbrell JA (2006) *In vitro* cytotoxicity assays: comparison of LDH, neutral red, MTT and protein assay in hepatoma cell lines following exposure to cadmium chloride. *Toxicology Letters*. **160**(2): 171–177
- Fraser MP, & Lakshmanan K (2000) Using levoglucosan as a molecular marker for the long-range transport of biomass combustion aerosols. *Environmental Science & Technology*. **34**(21): 4560–4564

- Fukumoto LR, & Mazza G (2000) Assessing antioxidant and prooxidant activities of phenolic compounds. *Journal of Agricultural and Food Chemistry*. **48**(8): 3597–3604
- Gadir SA (2012) Assessment of bioactivity of some Sudanese medicinal plants using brine shrimp (*Artemia salina*) lethality assay. *Journal of Chemical and Pharmaceutical Research*. **4**(12): 5145-5148
- Gangwar KK, & Joshi BD (2008) Diversity of medicinal flora of district Pithoragarh & their uses by the local communities. In Joshi BD, Tripathi CPM & Joshi PC (Eds.), *Biodiversity and Environmental Management* (pp. 136-146). New Delhi, India: APH Publishing House.
- Gao D, Kawai N, & Tamiya T (2011) The anti-inflammatory effects of D-allose contribute to attenuation of cerebral ischemia-reperfusion injury. *Medical Hypotheses*. **76**(6): 911–913
- Gao XZ, Zhou CX, Zhang SL, Yao W, & Zhao Y (2005) Studies on the chemical constituents in herb of *Ranunculus sceleratus*. *Zhongguo Zhong Yao Za Zhi*. **30**(2): 124-126
- Ghimire SK (2008) Medicinal plants in the Nepal Himalaya: current issues, sustainable harvesting, knowledge gaps and research priorities. In Jha PK, Karmacharya SB, Chettri MK, Thapa CB & Shrestha BB (Eds.), *Medicinal Plants in Nepal: an anthology of contemporary research* (pp. 187-193). Kathmandu, Nepal: Ecological Society.
- Ghimire SK, Thapa-Magar S, Shrestha MR, Devkota B, & Gubhaju MR (2008) Identification of important plant areas (IPAs) in Shivapuri-Langtang linkage area of sacred Himalayan landscape, Nepal. Kathmandu: Unpublished Report - WWF Nepal Program.
- Ghosal S, Tripathi VK, & Chauhan S (1996) Active constituents of *Embllica officinalis*: Part 1 - The chemistry and antioxidant effects of two new hydrolysable tannins, emblicanin A and B. *Indian Journal of Chemistry*. **35B**(9): 941-948
- Giday M, Asfaw Z, Woldu Z, & Teklehaymanot T (2009) Medicinal plant knowledge of the Bench ethnic group of Ethiopia: an ethnobotanical investigation. *Journal of Ethnobiology and Ethnomedicine*. **5**(1): 34
- Gilbert D (1981) *Oxygen and living processes: an interdisciplinary approach*. NY: Springer.
- Gnanavel V, & Saral AM (2013) GC-MS analysis of petroleum ether and ethanol extracts from *Abrus precatorius* Linn. *International Journal of Pharmacy and Biological Sciences*. **4**(3): 37-44
- Gong Y, Liu X, He WH, Xu HG, Yuan F, & Gao YX (2012) Investigation into the antioxidant activity and chemical composition of alcoholic extracts from defatted marigold (*Tagetes erecta* L.) residue. *Fitoterapia*. **83**(3): 481–489
- Gopalakrishnan K, & Udayakumar R (2014) GC-MS analysis of phytochemicals of leaf and stem of *Marsilea quadrifolia* (L.). *International Journal of Biochemistry Research & Review*. **4**(6): 517-526
- Gopalakrishnan S, & Vadivel E (2011) GC-MS analysis of some bioactive constituents of *Mussaenda frondosa* Linn. *International Journal of Pharmacy and Biological Sciences*. **2**(1): 313-320
- Gopinath S, Sakthidevi G, Muthukumaraswamy S, & Mohan VR (2013) GC-MS analysis of bioactive constituents of *Hypericum mysorens* (Hypericaceae). *Journal of Current Chemical & Pharmaceutical Sciences*. **3**(1): 6-15
- Graff CL, & Pollack GM (2005) Nasal drug administration: potential for targeted central nervous system delivery. *Journal of Pharmaceutical Sciences*. **94**(6): 1187–1195

- Grever MCB (2001) Cancer drug discovery and development. In DeVita VHS & Rosenberg SA (Eds.), *Cancer: Principles and practice of oncology* (pp. 328-339). Philadelphia: Lippincott Raven.
- Grierson AJC, & Long DG (1984-1999) *Flora of Bhutan*. UK: Royal Botanic Garden Edinburgh.
- Guevara AP, Vargas C, Sakurai H, Fujiwara Y, Hashimoto K, Maoka T, Kozuka M, Ito Y, Tokuda H, & Nishino H (1999) An antitumor promoter from *Moringa oleifera* Lam. *Mutation Research*. **440**(2): 181-188
- Guidi I, Galimberti D, Lonati S, Novembrino C, Bamonti F, Tiriticco M, Fenoglio C, Venturelli E, Baron P, Bresolin N, & Scarpini E (2006) Oxidative imbalance in patients with mild cognitive impairment and Alzheimer's disease. *Neurobiology of Aging*. **27**(2): 262–269
- Gunasekarana S, Vijay T, Sarumathy K, Palanie S, Panneerselvamb RPS, & Srinivasanb V (2013) Phytoconstituents evaluation by GC-MS and therapeutic efficacy of *Grewiaum bellifera* on streptozotocin (STZ)-induced diabetic rats. *International Journal of Pharmacy and Life Sciences*. **4**(2): 2380-2386
- Guo J, Yuan Y, Lu D, Du B, Xiong L, Shi J, Yang L, Liu W, Yuan X, Zhang G, & Wang F (2014) Two natural products, *trans*-phytol and (22E)-ergosta-6,9,22-triene-3 β ,5 α ,8 α -triol, inhibit the biosynthesis of estrogen in human ovarian granulosa cells by aromatase (CYP19). *Toxicology and Applied Pharmacology*. **279**(1): 23–32
- Guo L, Huang T, Hu B, & Wang C (2016) Bioactivity-guided isolation and characterization of antitumor active compound from marine-derived fungus *Penicillium chrysogenum* HGQ6. *Advance Journal of Food Science and Technology*. **10**(5): 353-355
- Guo XY, Wang J, Wang NL, Kitanaka S, & Yao XS (2007) 9, 10-Dihydrophenanthrene derivatives from *Pholidota yunnanensis* and scavenging activity on DPPH free radical. *Journal of Asian Natural Products Research*. **9**(2): 165-174
- Gupta A, & Sharma MC (2015) Biologically active long-chain aliphatic alcohols and esters from the bark of *Symplocos racemosa*. *International Journal of Pharmacognosy and Phytochemical Research*. **7**(5): 1056-1059
- Haddley K (2013) Trastuzumab emtansine for the treatment of HER2-positive metastatic breast cancer. *Drugs Today (Barc)*. **49**(11): 701–715
- Halliwel B (1996) Antioxidants in human health and disease. *Annual Review of Nutrition*. **16**: 33-50
- Halliwel B, & Guteridge JMC (1989) *Free Radicals in Biology and Medicine*. Oxford, UK: Clarendon Press.
- Haque R, Bin-Hafeez B, Ahmad I, Parvez S, Pandey S, & Raisuddin S (2001) Protective effects of *Emblca officinalis* Gaertn. in cyclophosphamide-treated mice. *Human & Experimental Toxicology*. **20**(12): 643-650
- Hartwell JL (1971) Plants used against cancer. A survey. *Lloydia*. **34**(4): 386-425
- Hauser R, & Calafat AM (2005) Phthalates and human health. *Occupational and Environmental Medicine*. **62**(11): 806–818
- Hazra B, Sarkar R, Biswas S, & Mandal N (2010) Comparative study of the antioxidant and reactive oxygen species scavenging properties in the extracts of the fruits of *Terminalia chebula*, *Terminalia belerica* and *Emblca officinalis*. *BMC Complementary and Alternative Medicine*. **10**: 20

- Hegnauer R (1973). In Hegnauer R (Ed.), *Chemotaxonomie der Pflanzen, Band VI* (pp. 478). Basel, Stuttgart: Birkh Nuser Verlag.
- Henderson PJ, & Giddens RA (1977) 2-Deoxy-D-galactose, a substrate for the galactose-transport system of *Escherichia coli*. *Biochemical Journal*. **168**(1): 15-22
- Hertog MG, Feskens EJ, Hollman PC, Katan MB, & Kromhout D (1993) Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen elderly study. *Lancet*. **342**(8878): 1007-1011
- Hesam F, Balali GR, & Tehrani RT (2012) Evaluation of antioxidant activity of three common potato (*Solanum tuberosum*) cultivars in Iran. *Avicenna Journal of Phytomedicine*. **2**(2): 79–85
- Hirata Y, Saito M, Tsukamoto I, Yamaguchi F, Sui L, Kamitori K, Dong Y, Uehara E, Konishi R, Janjua N, & Tokuda M (2009) Analysis of the inhibitory mechanism of D-allose on MOLT-4F leukemia cell proliferation. *Journal of Bioscience and Bioengineering*. **107**(5): 562–568
- HMDB (2016) Showing metabocard for Tiglic aldehyde (HMDB31512). Retrieved from <http://www.hmdb.ca/metabolites/HMDB31512>
- Holland HD (1997) Evidence for life on Earth more than 3,850 million years ago. *Science*. **275**: 38-39
- Hollman PC, & Katan MB (1999) Dietary flavonoids: intake, health effects and bioavailability. *Food and Chemical Toxicology*. **37**(9-10): 937-942
- Hollman PCH, Hertog MGL, & Katan MB (1996) Analysis and health effects of flavonoids. *Food Chemistry*. **57**(1): 43-46
- Hossain MA, Shah MD, Gnanaraj C, & Iqbal M (2011) *In vitro* total phenolics, flavonoids contents and antioxidant activity of essential oil, various organic extracts from the leaves of tropical medicinal plant *Tetrastigma* from Sabah. *Asian Pacific Journal of Tropical Medicine*. **4**(9): 717-721
- Hossain MA, Wakabayashi H, Goda F, Kobayashi S, Maeba T, & Maeta H (2000) Effect of the immunosuppressants FK506 and D-allose on allogenic orthotopic liver transplantation in rats. *Transplantation Proceedings*. **32**(7): 2021–2023
- Hossen SM, Sarkar R, & Mahmud S (2014) Medicinal potential of *Phyllanthus emblica* (Linn.) fruits extracts: biological and pharmacological activities. *British Journal of Pharmaceutical Research*. **4**(12): 1486-1499
- Hsieh IN, Chang AS, Teng CM, Chen CC, & Yang CR (2011) Aciculatin inhibits lipopolysaccharide-mediated inducible nitric oxide synthase and cyclooxygenase-2 expression *via* suppressing NF- κ B and JNK/p38 MAPK activation pathways. *Journal of Biomedical Science*. **18**: 28
- Hu YW, Liu CY, Du CM, Zhang J, Wu WQ, & Gu ZL (2009) Induction of apoptosis in human hepatocarcinoma SMMC-7721 cells *in vitro* by flavonoids from *Astragalus complanatus*. *Journal of Ethnopharmacology*. **123**(2): 293-301
- Huang Q, Huang R, Zhang S, Lin J, Wei L, He M, Zhuo L, & Lin X (2013a) Protective effect of genistein isolated from *Hydrocotyle sibthorpioides* on hepatic injury and fibrosis induced by chronic alcohol in rats. *Toxicology Letters*. **217**(2): 102-110

- Huang Q, Zhang S, Huang R, Wei L, Chen Y, Lv S, Liang C, Tan S, Liang S, Zhuo L, & Lin X (2013b) Isolation and identification of an anti-hepatitis B virus compound from *Hydrocotyle sibthorpioides* Lam. *Journal of Ethnopharmacology*. **150**(2): 568-575
- Huang SS, Huang GJ, Ho YL, Lin YH, Hung HJ, Chang TN, Chang MJ, Chen JJ, & Chang YS (2008) Antioxidant and antiproliferative activities of the four *Hydrocotyle* species from Taiwan. *Botanical Studies*. **49**: 311-322
- Husin F, Chan YY, Gan SH, Sulaiman SA, & Shueb RH (2015) The effect of *Hydrocotyle sibthorpioides* Lam. extracts on *in vitro* dengue replication. *Evidence-Based Complementary and Alternative Medicine*. **2015**
- Hyun DH, Hernandez JO, Mattson MP, & de Cabo R (2006) The plasma membrane redox system in aging. *Ageing Research Reviews*. **5**(2): 209–220
- Igbinosa OO, Igbinosa QC, & Aiyegoro OA (2009) Antimicrobial activity and phytochemical screening of stem bark extracts from *Jatropha curcas* (Linn). *African Journal of Pharmacy and Pharmacology*. **3**(2): 58-62
- Inbathamizh L, & Padmini E (2013) Quinic acid as a potent drug candidate for prostate cancer – a comparative pharmacokinetic approach. *Asian Journal of Pharmaceutical and Clinical Research*. **6**(4): 106-112
- Inouye H, Takeda Y, & Nishimura H (1973) On the lignan glucosides of *Symplocos lucida*. *Yakugaku Zasshi*. **93**(1): 44-46
- Inta A, Shengji P, Balslev H, Wangpakapattanawong P, & Trisonthi C (2008) A comparative study on medicinal plants used in Akha's traditional medicine in China and Thailand, cultural coherence or ecological divergence. *Journal of Ethnopharmacology*. **116**(3): 508-517
- Ishida J, Wang HK, Oyama M, Cosentino ML, Hu CQ, & Lee KH (2001) Anti-AIDS agents. 46. Anti-HIV activity of harman, an anti-HIV principle from *Symplocos setchuensis*, and its derivatives. *Journal of Natural Products*. **64**(7): 958-960
- Islam MT, de Alencar MV, da Conceição MK, da Conceição MK, de Carvalho M-CAA, de Sousa DP, & de Freitas RM (2015) Phytol in a pharma-medico-stance. *Chemico-Biological Interactions*. **240**: 60–73
- Ismail M, Hussain J, Khan AU, Khan AL, Ali L, Khan FU, Khan AZ, Niaz U, & Lee IJ (2012) Antibacterial, Antifungal, Cytotoxic, Phytotoxic, Insecticidal, and enzyme inhibitory activities of *Geranium wallichianum*. *Evidence-Based Complementary and Alternative Medicine*. **2012**
- Ismail M, Ibrar M, Iqbal Z, Hussain J, Hussain H, Ahmed M, Ejaz A, & Choudhary MI (2009) Chemical constituents and antioxidant activity of *Geranium wallichianum*. *Records of Natural Products*. **3**(4): 193-197
- Jagetia GC, Baliga MS, Malagi KJ, & Sethukumar KM (2002) The evaluation of the radioprotective effect of Triphala (an ayurvedic rejuvenating drug) in the mice exposed to gamma-radiation. *Phytomedicine*. **9**(2): 99-108
- Jananie RK, Priya V, & Vijayalakshmia K (2011) Determination of bioactive components of *Cynodon dactylon* by GC-MS analysis. *New York Science Journal*. **4**(4): 16-20
- Jang HD, Chang KS, Huang YS, Hsu CL, Lee SH, & Su MS (2007) Principal phenolic phytochemicals and antioxidant activities of three Chinese medicinal plants. *Food Chemistry*. **103**(3): 749–756

- Javale P, & Sabnis S (2010) Antimicrobial properties and phytochemical analysis of *Emblica officinalis*. *Asian Journal of Experimental Biological Sciences*: 91-95
- Jayaprakash GK, Mandadi KK, Poulouse SM, Jadegoud Y, Nagana GG, & Patil BS (2007) Inhibition of colon cancer cell growth and antioxidant activity of bioactive compounds from *Poncirus trifoliata* (L.) Raf. *Bioorganic & Medicinal Chemistry*. **15**(14): 4923-4932
- Jayaraman V, Devarsenapathi K, & Arumugame CT (2011) Phytochemical screening and antimicrobial activity of the stem of *Mallotus philippensis* (Lam.) Muell. Arg. Var. *philippensis* (Euphorbiaceae). *International Journal of Pharmacy and Pharmaceutical Sciences*. **3**(2): 160-163
- Jebastella J, & Reginald AM (2015) Bioactive components of *Cynodon dactylon* using ethanol extract. *World Journal of Pharmaceutical Sciences*. **3**(12): 2388-2391
- Jeena KJ, Joy KL, & Kuttan R (1999) Effect of *Emblica officinalis*, *Phyllanthus amarus* and *Picrorrhiza kurroa* on N-nitrosodiethylamine induced hepatocarcinogenesis. *Cancer Letters*. **136**(1): 11-16
- Jemal A, Bray F, Center M, Ferlay J, Ward E, & Forman D (2011) Global cancer statistics. *CA: A Cancer Journal for Clinicians*. **61**(2): 69–90
- Jeong JB, Hong SC, Jeong HJ, & Koo JS (2011) Anti-inflammatory effect of 2-methoxy-4-vinylphenol *via* the suppression of NF- κ B and MAPK activation, and acetylation of histone H3. *Archives of Pharmacal Research*. **34**(12): 2109-2116
- Jeong JB, & Jeong HJ (2010) 2-Methoxy-4-vinylphenol can induce cell cycle arrest by blocking the hyper-phosphorylation of retinoblastoma protein in benzo[a]pyrene-treated NIH3T3 cells. *Biochemical and Biophysical Research Communications*. **400**(4): 752-757
- Jimoh FO, Adedapo AA, Aliero AA, & Afolayan AJ (2008) Polyphenolic contents and biological activities of *Rumex ecklonianus*. *Pharmaceutical Biology*. **46**(5): 333–340
- Johanning E, Gareis M, Nielsen K, Dietrich R, & Märtlbauer E (2002) *Airborne mycotoxins sampling and screening analysis*, Monterey, CA.
- Johnson IT, Williamson G, & Musk SR (1994) Anticarcinogenic factors in plant foods: a new class of nutrients. *Nutrition Research Reviews*. **7**(1): 175–204
- Johnson LK, Haynes LW, Carlson MA, Fortnum HA, & Gorgas DL (1985) Alarm substances of the stingless bee, *Trigona silvestriana*. *Journal of Chemical Ecology*. **11**(4): 409-416
- Johnson T (1998) *CRC ethnobotany desk reference*. Boca Raton, Florida: CRC Press.
- Joshi AR, & Joshi K (2007) Ethnomedicinal plants used against skin diseases in some villages of Kali Gandaki, Bagmati, and Tadi Likhu watersheds of Nepal. *Ethnobotanical Leaflets*. **11**: 235-246
- Kabir KAA, & Nair VJ (2009) *Flora of Tamil Nadu: Grasses*. Howrah: Botanical Survey of India.
- Kafle L, & Shih CJ (2013) Toxicity and repellency of compounds from clove (*Syzygium aromaticum*) to red imported fire ants *Solenopsis invicta* (Hymenoptera: Formicidae). *Journal of Economic Entomology*. **106**(1): 131-135
- Kagoura MMC, & Morohashi M (1999) Phytol is a novel tumor promoter on ICR mouse skin. *Japanese journal of cancer research*. **90**(4): 377-384
- Kähkönen MP, Hopia AI, Heikki JV, Rauha JP, Pihlaja K, Kujala TS, & Heinonen M (1999) Antioxidant activity of plant extracts containing phenolic compounds. *Journal of Agricultural and Food Chemistry*. **47**(10): 3954–3962

- Kalt FR, & Cock IE (2014) Gas chromatography-mass spectroscopy analysis of bioactive petalostigma extracts: Toxicity, antibacterial and antiviral activities. *Pharmacognosy Magazine*. **10**(Suppl 1): S37-S49
- Kampa M, Nifli AP, Notas G, & Castanas E (2007) Polyphenols and cancer cell growth. *Reviews of Physiology, Biochemistry and Pharmacology*. **159**: 79-113
- Kanjilal UN (1997) *Flora of Assam*. New Delhi: Omsons Publications.
- Kapoor LD (1990) *Handbook of Ayurvedic Medicinal Plants*. Boca Raton, Florida CRC Press.
- Kasabri V, Flatt PR, & Abdel-Wahab YH (2010) *Terminalia bellirica* stimulates the secretion and action of insulin and inhibits starch digestion and protein glycation *in vitro*. *British Journal of Nutrition*. **103**(2): 212-217
- Kashiwada Y, Yamazaki K, Ikeshiro Y, Yamagishi T, Fujioka T, Mihashi K, Mizuki K, Cosentino LM, Fowke K, Morris-Natschke SL, & Lee KH (2001) Isolation of rhododaurichromanic acid B and the anti-HIV principles rhododaurichromanic acid A and rhododaurichromenic acid from *Rhododendron dauricum*. *Tetrahedron*. **57**(8): 1559-1563
- Kavitha S, Packia LM, Mary J, Kala S, & Mohan VR (2014) GC-MS analysis of ethanolic extract of *Nothapodytes nimmoniana* (Graham) Mabb. leaves. *Malaya Journal of Biosciences*. **2**(1): 42-49
- Khan AU, & Gilani AH (2008) Pharmacodynamic evaluation of *Terminalia belerica* for its antihypertensive effect. *Journal of Food and Drug Analysis*. **16**(3): 6-14
- Khan KH (2009) Role of *Embllica officinalis* in medicine - a review. *Botany Research International*. **2**(4): 218-228
- Khan R, Shawl AS, Tantry M, & Alam MS (2008) New coumarin glycosides from *Rhododendron lepidotum*. *Fitoterapia*. **79**(3): 232-233
- Khanahmadi M, Rezazadeh SH, & Taran M (2010) *In vitro* antimicrobial and antioxidant properties of *Smyrniium cordifolium* Boiss. (Umbelliferae) extract. *Asian Journal of Plant Science*. **9**(2): 99-103
- Khanbabaee K, & van Ree T (2001) Tannins: classification and definition. *Natural Product Reports*. **18**(6): 641-649
- Khanna P, & Bansal R (1975) Phyllantidine and phyllantine from *Embllica officinallis* Gaertn. leaves, fruit and *in vitro* tissue cultures. *Indian Journal of Experimental Biology*. **13**: 82-83
- Khatiwora E, Adsul VB, Kulkarni MM, Deshpande NR, & Kashalkar RV (2010) Spectroscopic determination of total phenol and flavonoid contents of *Ipomoea carnea*. *International Journal of ChemTech Research*. **2**(3): 1698-1701
- Kim CW, Lee HJ, Jung JH, Kim YH, Jung DB, Sohn EJ, Lee JH, Woo HJ, Baek NI, Kim YC, & Kim SH (2015) Activation of caspase-9/3 and inhibition of epithelial mesenchymal transition are critically involved in antitumor effect of phytol in hepatocellular carcinoma cells. *Phytotherapy Research*. **29**(7): 1026-1031
- Kim DO, Jeong SW, & Lee CY (2003) Antioxidant capacity of phenolic phytochemicals from various cultivars of plums. *Food Chemistry*. **81**(3): 321-326
- Kim JB, Koo HN, Joeng HJ, Lyu YS, Park SG, Won JH, Kim YK, Hong SH, & Kim HM (2005) Induction of apoptosis by Korean medicine Gagam-whanglyun-haedoktang through

- activation of caspase-3 in human leukemia cell line, HL-60 cells. *Journal of Pharmaceutical Sciences*. **97**(1): 138–145
- King A, & Young G (1999) Characteristics and occurrence of phenolic phytochemicals. *Journal of the American Dietetic Association*. **99**(2): 213-218
- Kinnula VL, & Crapo JD (2004) Superoxide dismutases in malignant cells and human tumors. *Free Radical Biology and Medicine*. **36**(6): 718-744
- Kinsella JE, Frankel E, German B, & Kanner J (1993) Possible mechanisms for the protective role of antioxidants in wine and plant foods. *Food Technol*. **47**: 85-89
- Kintzios SE (2006) Terrestrial plant-derived anticancer agents and plant species used in anticancer research. *Critical Reviews in Plant Sciences*. **25**(2): 79-113
- Kloocke AJ, Mei YH, Shin CF, & Kubo I (1986). *Phytochemistry*. **25**: 1637
- Knobloch K, Pauli A, Iberl B, Weigand H, & Weis N (1989) Antibacterial and antifungal properties of essential oil components. *Journal of Essential Oil Research*. **1**(3): 119-128
- Koch M, Kehop DA, Kinminja B, Sabak M, Wavimbukie G, Barrows KM, Matainaho TK, Barrows LR, & Rai PP (2015) An ethnobotanical survey of medicinal plants used in the East Sepik province of Papua New Guinea. *Journal of Ethnobiology and Ethnomedicine*. **11**(1): 1-26
- Kokate CK (2008) *Practical pharmacognosy*. New Delhi: Vallabh Prakashan.
- Konovalova O, Gergel E, & Herhel V (2013) GC-MS analysis of bioactive components of *Shepherdia argentea* (Pursh.) Nutt. from Ukrainian Flora. *The Pharma Innovation*. **2**(6): 7-12
- Kraus GA, & Frazier K (1980) Synthetic Studies toward Verrucarol. 1. Synthesis of the AB Ring System. *Journal of Organic Chemistry*. **45**(24): 4820-4825
- Krause JA, & Eggleston DS (1991) Structure of 8-(2,6-dideoxy-beta-ribo-hexopyranosyl)-5-hydroxy-2-(4-hydroxyphenyl)-7-methoxy-4H-1-benzopyran-4-one sesquihydrate, aciculatin. *Acta Crystallographica Section C, Crystal Structure Communications*. **47**(Pt 12): 2595–2598
- Krishnaraju AV, Roa-Tayi VN, Sundararaju D, Vanisree M, Tsay HS, & Subbaraju GV (2005) Assessment of bioactivity of Indian medicinal plants using brine shrimp (*Artemia salina*) lethality assay. *International Journal of Applied Science and Engineering*. **3**(2): 125-134
- Kristiansen E, Meyer O, & Thorup I (1996) Refined carbohydrate enhancement of aberrant crypt foci (ACF) in rat colon induced by the food-borne carcinogen 2-amino-3-methylimidazo[4,5-f]quinoline (IQ). *Cancer Letters*. **105**(2): 147–151
- Kulicic T, Radonic A, Katalinic V, & Milos M (2004) Use of different methods for testing antioxidative activity of oregano essential oil. *Food Chemistry*. **85**(4): 633-640
- Kumar A, & Chattopadhyay S (2007) DNA damage protecting activity and antioxidant potential of pudina extract. *Food Chemistry*. **100**(4): 1377-1384
- Kumar GS, Jayaveera KN, Kumar CK, Vrushabendra SBM, Sanjay UP, & Kishore KDV (2007) Antibacterial screening of selected Indian medicinal plants against acne-inducing bacteria. *Pharmacologyonline*. **2**: 34-47

- Kumar GS, Nayaka H, Dharmesh SM, & Slimath PV (2006) Free and bound phenolic antioxidants in amla (*Emblca officinalis*) and turmeric (*Curcuma longa*). *Journal of Food Composition and Analysis*. **19**(5): 446–452
- Kumar N, & Jangwan JS (2012) Phytoconstituents of *Symplocos paniculata* (leaves). *Journal of Current Chemical & Pharmaceutical Sciences*. **2**(1): 76-80
- Kumar NR, Reddy JS, Gopikrishna G, & Solomon KA (2012) GC-MS determination of bioactive constituents of *Cycas beddomei* cones. *International Journal of Pharma and Bio Sciences*. **3**(3): 344 - 350
- Kumar RZA, & Bhaskar A (2012) Phytochemical evaluation by GC-MS and *in vitro* antioxidant activity of *Punica granatum* fruit rind extract. *Journal of Chemical and Pharmaceutical Research*. **4**(6): 2869-2873
- Kumar SR, Sivakumar T, Arulmozhi KT, & Mythili N (2015) Gas chromatography - mass spectroscopy evaluation of bioactive phytochemicals of commercial green teas (*Camellia sinensis*) of India. *Asian Journal of Pharmaceutical and Clinical Research*. **8**(3): 278-282
- Kumaran A, & Karunakaran RJ (2006) Nitric oxide radical scavenging active components from *Phyllanthus emblica* L. *Plant Foods for Human Nutrition*. **61**(1): 1-5
- Kumarappan C, Jaswanth A, & Kumarasunderi K (2011) Antihaemolytic and snake venom neutralizing effect of some Indian medicinal plants. *Asian Pacific Journal of Tropical Medicine*. **4**(9): 743-747
- Kumarasamy Y, Byres M, Cox PJ, Jaspars M, Nahar L, & Sarker SD (2007) Screening seeds of some Scottish plants for free radical scavenging activity. *Phytotherapy Research*. **21**(7): 615-621
- Kumaravel S, Bhabha PK, & Singaravadeivel K (2016) GC-MS and FT-IR analysis of the spice ajwain (*Trachyspermum ammi*). *International Journal of Modern Research and Reviews*. **4**(3): 1114-1120
- Kunwar RM (2006) Non timber forest product of Nepal: a sustainable management approach. Nepal Japan: Center for Biological Conservation, Nepal and International Tropical Timber Organization, Japan.
- Kunwar RM, & Adhikari N (2005) Ethnomedicine of Dolpa district, Nepal: the plants, their vernacular names and uses. *Lyonia*. **8**(1): 43-49
- Kunwar RM, Shrestha KP, & Bussmann RW (2010) Traditional herbal medicine in Far-west Nepal: a pharmacological appraisal. *Journal of Ethnobiology and Ethnomedicine*. **6**(1): 35
- Kuo PC, Abe K, & Schroeder RA (2000) Superoxide enhances interleukin 1beta-mediated transcription of the hepatocyte-inducible nitric oxide synthase gene. *Gastroenterology*. **118**(3): 608-618
- Lage H, Duarte N, Coburger C, Hilgeroth A, & Ferreira MJU (2010) Antitumor activity of terpenoids against classical and atypical multidrug resistant cancer cells. *Phytomedicine*. **17**(6): 441-448
- Lai CY, Tsai AC, Chen MC, Chang LH, Sun HL, Chang YL, Chen CC, Teng CM, & Pan SL (2012) Aciculatin induces p53-dependent apoptosis *via* MDM2 depletion in human cancer cells *in vitro* and *in vivo*. *PLoS ONE*. **7**(8): e42192

- Lamichhane J, Bhattarai K, Shrivastava AK, Shrestha TM, & Jain SC (2014) Chemical constituents of *Rhododendron lepidotum* from Nepal Himalayas. *Chemistry of Natural Compounds*. **50**(4): 767–769
- Laser H (1951) The effect of *cis*-vaccenic acid on respiration and growth of *Bacillus subtilis*. *Biochemical Journal*. **48**(2): 164-170
- Laszlo J, Humphreys SR, & Goldin A (1960) Effects of glucose analogues (2-deoxy-D-glucose, 2-deoxy-D-galactose) on experimental tumors. *Journal of the National Cancer Institute*. **24**(2): 267-281
- Lawal B, Shittu OK, AbdulRasheed-Adeleke T, Ossai PC, & Ibrahim AM (2015) GC-MS determination of bioactive constituents of giant African snail (*Archachatina maginata*) haemolymph. *IOSR Journal of Pharmacy and Biological Sciences*. **10**(2): 59-64
- Lhullier C, Horta PA, & Falkenberg M (2006) Evaluation of macroalgae from Santa Catarina's coast with the brine shrimp assay. *Brazilian Journal of Pharmacognosy*. **16**(2): 158-163
- Li CJ, Wang LQ, Chen SN, & Qin GW (2000) Diterpenoids from the fruits of *Rhododendron molle*. *Journal of Natural Products*. **69**(9): 1214-1217
- Li F, Zheng Y, Lin X, Zhang J, Huang Y, Wu Y, Zheng T, Lin Y, Xie N, & Lin Z (2012) Allelopathy of orchard grass debris extracts on seed germination and seedling growth of *Borreria latifolia*. *Chinese Journal of Tropical Crops*. **33**(2): 290-295
- Li XF (1999) The buttercup medical plant resource of Qian Xi Na Zhou. *Chin Wild Plant Resour*. **18**: 49-50
- Li XH, Shen DD, Li N, & Yu SS (2003) Bioactive triterpenoids from *Symplocos chinensis*. *Journal of Asian Natural Products Research*. **5**(1): 49-56
- Li YH, & Lu XY (2005) Investigation on the origin of 5-HMF in Shengmai Yin decoction by RP-HPLC method. *Journal of Zhejiang University-Science B*. **6**(10): 1015–1021
- Lilhore L, & Pande SS (2016) National status of *in vitro* conservational strategies for *Leptadenia reticulata* (Retz) Wight and Arn. *Journal of Medicinal Plants Studies*. **4**(2): 160-162
- Lim SY, Meyer M, Kjonaas RA, & Ghosh SK (2006) Phytol-based novel adjuvants in vaccine formulation: 1. assessment of safety and efficacy during stimulation of humoral and cell-mediated immune responses. *Journal of Immune Based Therapies and Vaccines*. **4**: 6
- Lim YY, & Quah EPL (2007) Antioxidant properties of different cultivars of *Portulaca oleracea*. *Food Chemistry*. **103**(3): 734-740
- Limbu DK, & Rai BK (2013) Ethnomedicinal practices among the Limbu community in Limbuwan, eastern Nepal. *Global Journal of Human Social Science*. **13**(2)
- Liu X, Zhao M, Wang J, Yang B, & Jiang Y (2008) Antioxidant activity of methanolic extract of emblica fruit (*Phyllanthus emblica* L.) from six regions in China. *Journal of Food Composition and Analysis*. **21**(3): 219-228
- Liu XC, & Liu ZL (2014) Evaluation of larvicidal activity of the essential oil of *Ageratum conyzoides* L. aerial parts and its major constituents against *Aedes albopictus*. *Journal of Entomology and Zoology Studies*. **2**(4): 345-350
- Lohézic-Le DF, Tomasi S, Fontanel D, & Boustie J (2002) Flavonols from *Scurrula ferruginea* Danser (Loranthaceae). *Zeitschrift für Naturforschung C*. **57**(11-12): 1092-1095

- Lopez-Velez M, Martinez-Martinez F, & Del Valle-Ribes C (2003) The study of phenolic compounds as natural antioxidants in wine. *Critical Reviews in Food Science and Nutrition*. **43**(3): 233-244
- Lopez A, Hudson JB, & Towers GH (2001) Antiviral and antimicrobial activities of Colombian medicinal plants. *Journal of Ethnopharmacology*. **77**(2-3): 189-196
- Lopus M, Oroudjev E, Wilson L, Wilhelm S, Widdison W, Chari R, & Jordan MA (2010) Maytansine and cellular metabolites of antibody-maytansinoid conjugates strongly suppress microtubule dynamics by binding to microtubules. *Molecular Cancer Therapeutics*. **9**(10): 2689-2699
- Lopus M, Yenjerla M, & Wilson L (2009) Microtubule dynamics. In Begley TP (Ed.), *Encyclopedia of Chemical Biology* (pp. 153-160). New Jersey: John Wiley & Sons.
- Lu JC, Xu BB, Gao S, Fan L, Zhang HF, Liu RX, & Kodama H (2009) Structure elucidation of two triterpenoid saponins from rhizome of *Anemone raddeana* Regel. *Fitoterapia*. **80**(6): 345-348
- Lu XN, Liu XC, Liu QZ, & Liu ZL (2014) Isolation of insecticidal constituents from the essential oil of *Ageratum houstonianum* Mill. against *Liposcelis bostrychophila* Badonnel. *Journal of Chemistry*. **2014**
- Luceri C, Caderni G, Lancioni L, Aioli S, Dolara P, Mastrandrea V, Scardazza F, & Morozzi G (1996) Effects of repeated boluses of sucrose on proliferation and on AOM-induced aberrant crypt foci in rat colon. *Nutrition and Cancer*. **25**(2): 187-196
- Ma RJ, Liu ZH, Zi CT, Gao W, Dong FW, Yang L, Li J, Zhou J, & Hu JM (2016) Oleanane-type triterpene saponins from *Hydrocotyle nepalensis*. *Fitoterapia*. **110**: 66-71
- Mackeen MM, Ali AM, Lajis NH, Kawazu K, Hassan Z, Amran M, Habsah M, Mooi LY, & Mohamed SM (2000) Antimicrobial, antioxidant, antitumour-promoting and cytotoxic activities of different plant part extracts of *Garcinia atroviridis* Griff. ex T. Anders. *Journal of Ethnopharmacology*. **72**(3): 395-402
- Makihara H, Shimada T, Machida E, Oota M, Nagamine R, Tsubata M, Kinoshita K, Takahashi K, & Aburada M (2012) Preventive effect of *Terminalia bellirica* on obesity and metabolic disorders in spontaneously obese type 2 diabetic model mice. *Journal of Natural Medicines*. **66**(3): 459-467
- Manandhar NP (2002) *Plants and People of Nepal*. Portland, Oregon, USA: Timber Press.
- Mandal M, Paul S, Uddin MR, Mondal MA, Mandal S, & Mandal V (2016) *In vitro* antibacterial potential of *Hydrocotyle javanica* Thunb. *Asian Pacific Journal of Tropical Disease*. **6**(1): 54-62
- Mandelbaum-Shavit F, Wolpert-DeFilippes MK, & Johns DG (1976) Binding of maytansine to rat brain tubulin. *Biochemical and Biophysical Research Communications*. **72**(1): 47-54
- Manjunatha M, Bhalodiya H, Ansari MA, Vada S, & Goli D (2011) Immunomodulatory activity of *Terminalia bellirica* extract in mice. *International Journal of Pharmagenesis*. **2**(1): 103-108
- Manohar VR, Chandrashekar R, & Rao SN (2012) Phytochemical analysis of *Terminalia belerica* fruit pulp extracts. *World journal of Pharmacy and Pharmaceutical Sciences*. **1**(4): 1376-1383

- Manorenjitha MS, Norita AK, Norhisham S, & Asmawi MZ (2013) GC-MS analysis of bioactive components of *Ficus religiosa* (Linn.) stem. *International Journal of Pharma and Bio Sciences*. **4**(2): 99 - 103
- Maruthupandian A, & Mohan VR (2011) GC-MS analysis of some bioactive constituents of *Pterocarpus marsupium* Roxb. *International Journal of ChemTech Research*. **3**(3): 1652-1657
- Mathers C, Boschi-Pinto C, Lopez A, & Murray C (2001) Cancer incidence, mortality and survival by site for 14 regions of the world. Lyon, France: World Health Organization.
- Matsushita A, Sasaki Y, Warashina T, Miyase T, Noguchi H, & Velde DV (2004) Hydrocotylosides I–VII, New Oleanane Saponins from *Hydrocotyle sibthorpioides*. *Journal of Natural Products*. **67**(3): 384-388
- Mayorga P, Pérez KR, Cruz SM, & Cáceres A (2010) Comparison of bioassays using the anostracan crustaceans *Artemia salina* and *Thamnocephalus platyurus* for plant extract toxicity screening. *Revista Brasileira de Farmacognosia*. **20**(6): 897-903
- McLaughlin JL, Chang CJ, & Smith DL (1991) Bench-top bioassays for the discovery of bioactive natural products: an update. In Atta-ur-Rahman (Ed.), *Studies in Natural Products Chemistry* (Vol. 9, pp. 283-409). Amsterdam: Elsevier.
- McLaughlin JL, Rogers LL, & Anderson JE (1998) The use of biological assays to evaluate botanicals. *Drug Information Journal*. **32**: 513-524
- Mehjabeen AM, N J, Mahayrookh, B RA, S M, & Obaidullah (2014) Antidiarrhoeal, Anti-inflammatory and analgesic activities of *Symplocos racemosa* Roxb. bark. *Pakistan Journal of Pharmaceutical Sciences*. **27**(6): 2221-2226
- Mehta R, Murillo G, Naithani R, & Peng X (2010) Cancer chemoprevention by natural products: how far have we come. *Pharmaceutical Research*. **27**: 950-961
- Mei H, Zuo S, Ye L, Wang J, & Ma S (2012) Review of the application of the traditional Chinese medicinal herb, *Ranunculus sceleratus* Linn. *Journal of Medicinal Plants Research*. **6**(10): 1821-1826
- Mendelson R, & Balick MJ (1995) The value of undiscovered pharmaceuticals in tropical forests. *Economic Botany*. **49**(2): 223-228
- Mendes V, Vilaça R, de Freitas V, Ferreira PM, Mateus N, & Costa V (2015) Effect of myricetin, pyrogallol, and phloroglucinol on yeast resistance to oxidative stress. *Oxidative Medicine and Cellular Longevity*. **2015**: 782504
- Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nichols DE, & McLaughlin JL (1982) Brine shrimp: a convenient general bioassay for active plant constituents. *Planta Medica*. **45**(5): 31-34
- Michael AS, Thompson CG, & Abramovitz M (1956) *Artemia salina* as a test organism for bioassay. *Science (NY)*. **123**(3194): 464
- Mickymaray S, Al Aboody MS, Rath PK, Annamalai P, & Nooruddin T (2016) Screening and antibacterial efficacy of selected Indian medicinal plants. *Asian Pacific Journal of Tropical Biomedicine*. **6**(3): 185-191
- Miller NJ, & Ruiz-Larrea MB (2002) Flavonoid and other plant phenols in diet: their significance as antioxidants. *Journal of Nutritional & Environmental Medicine*. **12**(1): 39-51

- Mimaki Y, Watanabe K, Matsuo Y, & Sakagami H (2009) Triterpene glycosides from the tubers of *Anemone coronaria*. *Chemical and Pharmaceutical Bulletin*. **57**(7): 724–729
- Misra SB, & Dixit SN (1978) Antifungal properties of leaf extract of *Ranunculus sceleratus* L. *Experientia*. **34**(11): 1442-1443
- Mitani T, Hoshikawa H, Mori T, Hosokawa T, Tsukamoto I, Yamaguchi F, Kamitori K, Tokuda M, & Mori N (2009) Growth inhibition of head and neck carcinomas by D-allose. *Head & Neck*. **31**(8): 1049-1055
- Molyneux P (2004) The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarinn Journal of Science and Technology*. **26**(2): 211-219
- Moon DO, Lee KJ, Choi YH, & Kim GY (2007) Beta-sitosterol-induced-apoptosis is mediated by the activation of ERK and the downregulation of Akt in MCA-102 murine fibrosarcoma cells. *International Immunopharmacology*. **7**(8): 1044-1053
- Moreno JJ (2003) Effect of olive oil minor components on oxidative stress and arachidonic acid mobilization and metabolism by macrophages RAW 264.7. *Free Radical Biology and Medicine*. **35**(9): 1073-1081
- Moshi MJ, Innocent E, Magadula JJ, Otieno DF, Weisheit A, Mbabazi PK, & Nondo RS (2010) Brine shrimp toxicity of some plants used as traditional medicines in Kagera Region, north western Tanzania. *Tanzania Journal of Health Research*. **12**(1): 63-67
- Mosquera OM, Correa YM, Buitrago DC, & Nino J (2007) Antioxidant activity of twenty five plants from Colombian biodiversity. *Memórias do Instituto Oswaldo Cruz*. **102**(5): 631-634
- Mujeeb F, Bajpai P, & Pathak N (2014) Phytochemical evaluation, antimicrobial activity, and determination of bioactive components from leaves of *Aegle marmelos*. *BioMed Research International*. **2014**
- Mukherjee AK, Basu S, Sarkar N, & Ghosh AC (2001) Advances in cancer therapy with plant based natural products. *Current Medicinal Chemistry*. **8**(12): 1467-1486
- Müller C, Lang R, & Hofmann T (2006) Quantitative precursor studies on di- and trihydroxybenzene formation during coffee roasting using "in bean" model experiments and stable isotope dilution analysis. *Journal of Agricultural and Food Chemistry*. **54**(26): 10086–10091
- Mulligan ME, Murray-Leisure KA, Ribner BS, Standiford HC, John JF, Korvick JA, Kauffman CA, & Yu VL (1993) Methicillin-resistant *Staphylococcus aureus*: a consensus review of the microbiology, pathogenesis, and epidemiology with implications for prevention and management. *American Journal of Medicine*. **94**(3): 313-328
- Murata A, Sekiya K, Watanabe Y, Yamaguchi F, Hatano N, Izumori K, & Tokuda M (2003) A novel inhibitory effect of D-allose on production of reactive oxygen species from neutrophils. *Journal of Bioscience and Bioengineering*. **96**(1): 89-91
- Murwani R (2003) Indonesian tea mistletoe (*Scurrula oortiana*) stem extract increases tumour cell sensitivity to tumour necrosis factor alpha (TNFalpha). *Phytotherapy Research*. **17**(4): 407-409
- Nadkarni KM (1976) *Indian Materia Medica*. Mumbai, India: Popular Prakashan.

- Nagelkerke F, & Postma PW (1978) 2-deoxygalactose, a specific substrate of the *Salmonella typhimurium* galactose permease: its use for the isolation of galP mutants. *Journal of Bacteriology*. **133**(2): 607–613
- Nagendraswamy G, V LR, A BB, T P, Alghorbani M, Zameer F, & Khanum SA (2013) Extraction and evaluation of antimicrobial and antiviral efficacy of *Terminalia bellirica* Fruits. *Indo American Journal of Pharmaceutical Research*. **3**(6): 4262-4268
- Naha N, Lee HY, Jo MJ, Chung BC, Kim SH, & Kim MO (2008) Rare sugar D-allose induces programmed cell death in hormone refractory prostate cancer cells. *Apoptosis*. **13**(9): 1121–1134
- Naidu JR, Ismail R, & Sasidharan S (2014) Acute oral toxicity and brine shrimp lethality of methanol extract of *Mentha spicata* L (Lamiaceae). *Tropical Journal of Pharmaceutical Research*. **13**(1): 101-107
- Nampoothiri SV, Prathapan A, Cherian OL, Raghu KG, Venugopalan VV, & Sundaresan A (2011) *In vitro* antioxidant and inhibitory potential of *Terminalia bellerica* and *Emblica officinalis* fruits against LDL oxidation and key enzymes linked to type 2 diabetes. *Food and Chemical Toxicology*. **49**(1): 125-131
- Nandy AK, Podder G, Sahu NP, & Mahato SB (1989) Triterpenoids and their glucosides from *Terminalia bellerica*. *Phytochemistry*. **28**(10): 2769-2772
- Naqvi SNH, & Parveen F (1991) Toxicity and residual effect of *Nerium indicum* crude extract as compared with Coopex against adults of *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Pakistan Journal of Entomology Karachi*. **6**(1-2): 35-44
- Neda GD, Rabeta MS, & Ong MT (2013) Chemical composition and anti-proliferative properties of flowers of *Clitoria Ternatea*. *International Food Research Journal*. **20**(3): 1229-1234
- Nepal M (1999) Ethnobotany of Rai and Sherpa community in the Makalu Barun conservation area, eastern Nepal. Kathmandu: MSc Thesis submitted to the Central Department of Botany, Tribhuvan University.
- Newman DJ, & Cragg GM (2012) Natural products as sources of new drugs over the 30 years from 1981 to 2010. *Journal of Natural Products*. **75**(3): 311-335
- Newmann DJ, & Cragg GM (2007) Natural products as sources of new drugs over the last 25 years. *Journal of Natural Products*. **70**(3): 461-477
- Newmann DJ, Cragg GM, & Snader KM (2003) Natural products as sources of new drugs over the period 1981-2002. *Journal of Natural Products*. **66**(7): 1022-1037
- Ngamkitidechakul C, Jaijoy K, Hansakul P, Soonthornchareonnon N, & Sireeratawong S (2010) Antitumour effects of *Phyllanthus emblica* L.: induction of cancer cell apoptosis and inhibition of *in vivo* tumour promotion and *in vitro* invasion of human cancer cells. *Phytotherapy Research*. **24**(9): 1405–1413
- Nguyen MTT, Nguyen NT, Nguyen HX, Huynh TNN, & Min BS (2012) Screening of α -glucosidase inhibitory activity of Vietnamese medicinal plants: isolation of active principles from *Oroxylum indicum*. *Natural Product Sciences*. **18**(1): 47-51
- Noltie HJ (2000) *Flora of Bhutan: including a record of plants from Sikkim and Darjeeling* (Vol. 3). Edinburgh: Royal Botanic Garden Edinburgh.
- O'Neil MJ (2006) *The Merck index: an encyclopedia of chemicals, drugs, and biologicals*. Whitehouse Station, NJ: Merck and Co. Inc.

- O'Neill AR, & Rana SK (2016) An ethnobotanical analysis of parasitic plants (Parijibi) in the Nepal Himalaya. *Journal of Ethnobiology and Ethnomedicine*. **12**(1): 14
- Ogugu SE, Kehinde AJ, James BI, & Paul DK (2012) Assessment of cytotoxic effects of methanol extract of *Calliandra portoricensis* using brine shrimp (*Artemia salina*) lethality bioassay. *Global Journal of Bio-Science and BioTechnology*. **1**(2): 257-260
- Ogunnusi TA, & Dosumu OO (2008) Bioactivity of crude extracts of *Euphorbia kamerunica* Pax using brine shrimp (*Artemia salina*) lethality assay. *Journal of medicinal plant research*. **2**(12): 370-373
- Ohashi K, Winarno H, Mukai M, Inoue M, Prana MS, Simanjuntak P, & Shibuya H (2003) Indonesian medicinal plants. XXV. Cancer cell invasion inhibitory effects of chemical constituents in the parasitic plant *Scurrula atropurpurea* (Loranthaceae). *Chemical and Pharmaceutical Bulletin*. **51**(3): 343–345
- Okuda T, Hatano T, & Yazaki K (1982) Dehydrogeraniin, furosinin and furosin, dehydroellagitannins from *Geranium thunbergii*. *Chemical and Pharmaceutical Bulletin*. **30**(3): 1113-1116
- Oskoueian E, Abdullah N, Ahmad S, Saad WZ, Omar AR, & Ho YW (2011) Bioactive compounds and biological activities of *Jatropha curcas* L. kernel meal extract. *International Journal of Molecular Sciences*. **12**(9): 5955-5970
- Ozçelik B, Kartal M, & Orhan I (2011) Cytotoxicity, antiviral and antimicrobial activities of alkaloids, flavonoids, and phenolic acids. *Pharmaceutical Biology*. **49**(4): 396-402
- Ozsoy N, Can A, Yanardag R, & Akev N (2008) Antioxidant activity of *Smilax excelsa* L. leaf extracts. *Food Chemistry*. **110**(3): 571–583
- Palombo EA (2011) Traditional medicinal plant extracts and natural products with activity against oral bacteria: potential application in the prevention and treatment of oral diseases. *Evidence-Based Complementary and Alternative Medicine*. **2011**(1741-427X): 680354
- Pandey G (2002) *Anticancer herbal drugs of India with special reference to Ayurveda*. New Delhi: Sri Satguru Publications.
- Paria ND, & Chattopadhyay SP (2005) *Flora of Hazaribagh District* (Vol. 2). Howrah: Botanical Survey of India.
- Park HJ, Kim MJ, Ha E, & Chung JH (2008) Apoptotic effect of hesperidin through caspase3 activation in human colon cancer cells, SNU-C4. *Phytomedicine*. **15**(1-2): 147-151
- Park KH, & Lee MW (2012) Anti-oxidative, anti-inflammatory and whitening effects of phenolic compounds from *Bambusae Caulis* in Liquamen. *Natural Product Research*. **26**(18): 1687-1691
- Parra LA, Yhebra RS, Sardinas IG, & Buela LI (2001) Comparative study of the assay of *Artemia salina* L. and the estimate of the medium lethal dose (LD50 value) in mice, to determine oral acute toxicity of plant extracts. *Phytomedicine*. **8**(5): 395-400
- Parthipan B, Suky MGT, & Mohan VR (2015) GC-MS analysis of phytochemicals in *Pleiospermium alatum* (Wall. ex Wight & Arn.) Swingle, (Rutaceae). *Journal of Pharmacognosy and Phytochemistry*. **4**(1): 216-222
- Patwardhan B, Vaidhya ADB, & Chorghade M (2004) Ayurveda and natural products drug discovery. *Current Science*. **86**(6): 789-799

- Pejin B, Kojic V, & Bogdanovic G (2014) An insight into the cytotoxic activity of phytol at *in vitro* conditions *Natural Product Research*. **28**(22): 2053–2056
- Pękal A, & Pyrzyńska K (2014) Evaluation of aluminium complexation reaction for flavonoid content assay. *Food Analytical Methods*. **7**(9): 1776-1782
- Pelka M, Danzl C, Distler W, & Petschelt A (2000) A new screening test for toxicity testing of dental materials. *Journal of Dentistry*. **28**(5): 341-345
- Pero RW, & Lund H (2009) *In vivo* treatment of humans with quinic acid enhances DNA repair and reduces the influence of lifestyle factors on risk to disease. *International Journal of Biotechnology and Biochemistry*. **5**(3): 293–305
- Pero RW, Lund H, & Leanderson T (2009) Antioxidant metabolism induced by quinic acid. Increased urinary excretion of tryptophan and nicotinamide. *Phytotherapy Research*. **23**(3): 335-346
- Perry LM (1980) *Medicinal plants of East and Southeast Asia : attributed properties and uses*. Cambridge, Massachusetts: MIT Press.
- Peters GJ, Backus HH, Freemantle S, van Triest B, Codacci-Pisanelli G, van der Wilt CL, Smid K, Lunec J, Calvert AH, Marsh S, McLeod HL, Bloemena E, Meijer S, Jansen G, van Groeningen CJ, & Pinedo HM (2002) Induction of thymidylate synthase as a 5-fluorouracil resistance mechanism. *Biochimica et Biophysica Acta*. **1587**(2-3): 194-205
- Phang CW, Malek SNA, Ibrahim H, & Wahab NA (2011) Antioxidant properties of crude and fractionated extracts of *Alpinia mutica* rhizomes and their total phenolic content. *African journal of pharmacy and pharmacology*. **5**(7): 842-852
- Piddock KJV, & Wise R (1989) Mechanisms of resistance to quinolones and clinical perspective. *Journal of Antimicrobial Chemotherapy*. **23**: 475-483
- Pietta PG (2000) Flavonoids as antioxidants. *Journal of Natural Products*. **63**(7): 1035-1042
- Pinmai K, Chunlaratthanabhorn S, Ngamkitidechakul C, Soonthornchareon N, & Hahnvajanawong C (2008) Synergistic growth inhibitory effects of *Phyllanthus emblica* and *Terminalia bellerica* extracts with conventional cytotoxic agents: doxorubicin and cisplatin against human hepatocellular carcinoma and lung cancer cells. *World Journal of Gastroenterology*. **14**(10): 1491-1497
- Pisutthanan S, Plianbangchang P, Pisutthanan N, Ruanruay S, & Muanrit O (2004) Brine shrimp lethality activity of Thai medicinal plants in the family Meliaceae. *Naresuan University Journal*. **12**(2): 13-18
- Polunin O, & Stainton A (1987) *Flowers of the Himalayas*. New Delhi, India: Oxford University Press.
- Popendorf W, Selim M, & Lewis MQ (1995) Exposure while applying industrial antimicrobial pesticides. *American Industrial Hygiene Association Journal*. **56**(10): 993–1001
- Popescu R, & Kopp B (2013) The genus *Rhododendron*: an ethnopharmacological and toxicological review. *Journal of Ethnopharmacology*. **147**(1): 42-62
- Prabhu VV, & Gopal V (2011) Evaluation of angiogenesis potential of *Terminalia bellirica* Roxb. by chick chorio-allantoic membrane (CAM) assay. *International Journal of Drug Development & Research*. **3**(2): 256-264
- Prabu K, Samyurai P, Subbaiyan B, & Thangapandian V (2013) Phytochemical constituents and gas chromatography - mass spectrometry analysis of *Caralluma diffusa* (Wight) N.

- E. Br. aerial part. *International Journal of Pharmacy & Pharmaceutical Sciences*. **5**(3): 602-605
- Press JR, Shrestha KK, & Sutton DA (2000) *Annotated checklist of the flowering plants of Nepal*. London: Natural History Museum.
- Pyla R, Kim TJ, Silva JL, & Jung YS (2010) Enhanced antimicrobial activity of starch-based film impregnated with thermally processed tannic acid, a strong antioxidant. *International Journal of Food Microbiology*. **137**(2-3): 154-160
- Qadan F, Thewaini AJ, Ali DA, Afifi R, Elkhawad A, & Matalka KZ (2005) The antimicrobial activities of *Psidium guajava* and *Juglans regia* leaf extracts to acne-developing organisms. *The American Journal of Chinese Medicine*. **33**(2): 197-204
- Qiu S, Sun H, Zhang AH, Xu HY, Yan GL, Han Y, & Wang XJ (2014) Natural alkaloids: basic aspects, biological roles, and future perspectives. *Chinese Journal of Natural Medicines*. **12**(6): 401-406
- Quamar MF, & Bera SK (2014) Ethno-medico-botanical studies of plant resources of Hoshangabad district, Madhya Pradesh, India: retrospect and prospects. *Journal of plant science and research*. **1**(1): 1-11
- Quideau S, Deffieux D, Douat-Casassus C, & Pouységu L (2011) Plant polyphenols: chemical properties, biological activities, and synthesis. *Angewandte Chemie International Edition*. **50**(3): 586-621
- Rafat A, Philip K, & Muniandy S (2010) Antioxidant potential and phenolic content of ethanolic extract of selected Malaysian plants. *Research journal of biotechnology*. **5**(1): 16–19
- Raghu HS, & Ravindra P (2010) Antimicrobial activity and phytochemical study of *Phyllanthus emblica* Linn. *International Journal of Pharmaceutical Studies and Research*. **1**(1): 30-33
- Rahman S, Akbor MM, Howlader A, & Jabbar A (2009) Antimicrobial and cytotoxic activity of the alkaloids of Amlaki (*Emblca officinalis*). *Pakistan Journal of Biological Sciences*. **12**(16): 1152–1155
- Rajbhandari KR (2001) *Ethnobotany of Nepal*. Kathmandu: Ethnobotanical Society of Nepal.
- Rajbhandari M, Mentel R, Jha PK, Chaudhary RP, Bhattarai S, Gewali MB, Karmacharya N, Hipper M, & Lindequist U (2009) Antiviral activity of some plants used in Nepalese traditional medicine. *Evidence-Based Complementary and Alternative Medicine*. **6**(4): 517-522
- Rajeswari G, Murugan M, & Mohan VR (2012) GC-MS analysis of bioactive components of *Hugonia mystax* L. (Linaceae). *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. **3**(4): 301-308
- Rajeswari G, Murugan M, & Mohan VR (2013) GC-MS analysis of bioactive components of *Hugonia mystax* L. bark (Linaceae). *Journal of pharmaceutical and biomedical sciences*. **29**(29): 818-824
- Rajeswari J, & Rani S (2014) GC-MS analysis of whole plant of *Leptadenia reticulata*. *International Journal of PharmTech Research*. **6**(7): 2043-2050
- Rajeswari J, & Rani S (2014-15) GC-MS analysis of phytochemical compounds in the ethanolic extract of root of *Lawsonia inermis* Linn. *International Journal of ChemTech Research*. **7**(1): 389-399

- Ramachandran S, Vamsikrishna M, Gowthami KV, Heera B, & Dhanaraju MD (2011) Assessment of cytotoxic activity of *Agave cantula* using brine shrimp (*Artemia salina*) lethality assay. *Asian Journal of Scientific Research*. **4**(1): 90-94
- Ramakrishna BS, Varghese R, Jayakumar S, Mathan M, & Balasubramanian KA (1997) Circulating antioxidants in ulcerative colitis and their relationship to disease severity and activity. *Journal of Gastroenterology and Hepatology*. **12**(7): 490–494
- Ramamoorthy PK, & Awang B (2007) Antioxidant activity, total phenolic and flavonoid content of *Morinda citrifolia* fruit extracts from various extraction processes. *Journal of engineering Science and Technology*. **2**(1): 70-80
- Ramawat KG (2007) Secondary metabolites in nature. In Ramawat KG & Merillon JM (Eds.), *Biotechnology: secondary metabolites* (pp. 21). Enfield, CT: Science Publishers.
- Rameshkumar KB, George V, & Shiburaj S (2007) Chemical constituents and antibacterial activity of the leaf oil of *Cinnamomum chemungianum* Mohan & Henry. *Journal of Essential Oil Research*. **19**(1): 98–100
- Ramos F, Takaishi Y, Kawazoe K, Osorio C, Duque C, Acuña R, Fujimoto Y, Sato M, Okamoto M, Oshikawa T, & Ahmed SU (2006) Immunosuppressive diacetylenes, ceramides and cerebroside from *Hydrocotyle leucocephala*. *Phytochemistry*. **67**(11): 1143-1150
- Rao MRR, & Siddiqui HH (1964) Pharmacological studies on *Emblica officinallis* Gaertn. *Indian Journal of Experimental Biology*. **2**: 29
- Rashid MA, Ahmad VU, Abbasi M, Ali Z, Rasool N, Zubair M, Lodhi MA, Choudhary MI, & Khan IA (2008) α -Chymotrypsin inhibiting benzylated glycoside from *Symplocos racemosa*. *Phytochemistry Letters*. **1**(1): 54-58
- Rastogi RP, & Mehrotra BN (1993) *Compendium of Indian medicinal plants*. New Delhi: Central Drug Research Institute and Publications & Information Directorate.
- Ratnam DV, Ankola DD, Bhardwaj V, Sahana DK, & Kumar MN (2006) Role of antioxidants in prophylaxis and therapy: a pharmaceutical perspective. *Journal of Controlled Release*. **113**(3): 189-207
- Rauha JP, Remes S, Heinonen M, Hopia A, Kähkönen M, Kujala T, Pihlaja K, Vuorela H, & Vuorela P (2000) Antimicrobial effects of finnish plant extracts containing flavonoids and other phenolic compounds. *International Journal of Food Microbiology*. **56**(1): 3-12
- Rawal JR, & Sonawani PR (2016) Determination of bioactive components of *Cynodon dactylon* by GC-MS analysis & its *in vitro* antimicrobial activity. *International Journal of Pharmacy and Life Sciences*. **7**(1): 4880-4885
- Reed JC, & Pellecchia M (2005) Apoptosis-based therapies for hematologic malignancies. *Blood*. **106**(2): 408-418
- Remillard S, Rebhun LI, Howie GA, & Kupchan SM (1975) Antimitotic activity of the potent tumor inhibitor maytansine. *Science*. **189**(4207): 1002–1005
- Revathi P, Parimelazhagan T, & Manian S (2013) Quantification of phenolic compounds, *in vitro* antioxidant analysis and screening of chemical compounds using GC-MS in *Acalypha alnifolia* Klein ex Willd. - a leafy vegetable. *International Journal of Pharma and Bio Sciences*. **4**(2): B973-B986

- Rice-Evans CA, Miller NJ, Bolwell PG, Bramley PM, & Pridham JB (1995) The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radical Research*. **22**(4): 375-383
- Rice-Evans CA, Miller NJ, & Paganga G (1996) Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radical Biology and Medicine*. **20**(7): 933-956
- Richardson MA (2001) Biopharmacologic and herbal therapies for cancer: research update from NCCAM. *Journal of Nutrition*. **131**(11 Suppl): 3037S-3040S
- Rio DD, Rodriguez-Mateos A, Spencer JPE, Tognolini M, Borges G, & Crozier A (2013) Dietary (poly)phenolics in human health: structures, bioavailability, and evidence of protective effects against chronic diseases. *Antioxidants & Redox Signaling*. **18**(14): 1818-1892
- Rocha FF, Almedia CS, dos Santos RT, Santana SA, Costa EA, de Paula JR, & Vanderlinde FA (2011) Anxiolytic-like and sedative effects of *Hydrocotyle umbellata* L., Araliaceae, extract in mice. *Revista Brasileira de Farmacognosia*. **21**: 115-120
- Rockwell S, Mroczkowski Z, & Rupp WD (1982) Evaluation of 2-amino-5-nitrothiazole as a hypoxic cell radiosensitizer. *Radiation Research*. **90**(3): 575-585
- Ryu KR, Choi JY, Chung S, & Kim DH (2011) Anti-scratching behavioral effect of the essential oil and phytol isolated from *Artemisia princeps* Pamp. in mice. *Planta Medica*. **77**(1): 22-26
- Saeed S, & Tariq P (2007) Antibacterial activities of *Emblica officinalis* and *Coriandrum sativum* against gram-negative urinary pathogens. *Pakistan Journal of Pharmaceutical Sciences*. **20**(1): 32-35
- Saha A, Hoque A, Mallick SK, & Panda S (2014) Medicinal uses of grasses by the tribal people in West Bengal - an overview. *International Journal of Basic and Applied Sciences*. **3**(3): 63-70
- Sai Ram M, M N, Dipti P, Vandana M, lavazhagan L, Kumar G, & Selvamurthy W (2003) Cytoprotective activity of Amla (*Emblica officinalis*) against chromium (VI) induced oxidative injury in murine macrophages. *Phytotherapy Research*. **17**(4): 430-433
- Saini A, Sharma S, & Chhibber S (2008) Protective efficacy of *Emblica officinalis* against *Klebsiella pneumoniae* induced pneumonia in mice. *Indian Journal of Medical Research*. **128**(2): 188-193
- Saleem MA, & Wilkins RM (1984) Precocene-I: an anti-juvenile hormone, a potential 4th generation insecticide against a malathion-resistant strain of *Oryzaephilus surinamensis* (L.). *Pakistan Journal of Zoology*. **16**(2): 195-201
- Salem M, Zeidler A, Bohm M, Mohamed M, & Ali H (2015) GC/MS analysis of oil extractives from wood and bark of *Pinus sylvestris*, *Abies alba*, *Picea abies*, and *Larix decidua*. *BioResources*. **10**(4): 7725-7737
- Salunkhe DK, Jadhav SJ, Kadam SS, & Chavan JK (1982) Chemical, biochemical, and biological significance of polyphenols in cereals and legumes. *Critical Reviews in Food Science and Nutrition*. **17**(3): 277-305
- Santos CC, Salvadori MS, Mota VG, Costa LM, de Almeida AA, de Oliveira GA, Costa JP, de Sousa DP, de Freitas RM, & de Almeida RN (2013) Antinociceptive and antioxidant activities of phytol *in vivo* and *in vitro* models. *Neuroscience Journal*. **2013**: 949452

- Saraswathi MN, Karthikeyan M, Kannan M, & Rajasekar S (2012) *Terminalia belerica* Roxb-A phytopharmacological review. *International Journal of Research in Pharmaceutical and Biomedical Sciences*. **3**(1): 96-99
- Saravanan P, Chandramohan G, Mariajancyrani J, & Shanimugasundaram P (2014) GC-MS analysis of phytochemical constituents in ethanolic bark extract of *Ficus religiosa* Linn. *International Journal of Pharmacy and Pharmaceutical Sciences*. **6**(1): 457-460
- Sas K, Robotka H, Toldi J, & Vecsei L (2007) Mitochondria, metabolic disturbances, oxidative stress and the kynurenine system, with focus on neurodegenerative disorders. *Journal of the Neurological Sciences*. **257**(1-2): 221-239
- Sastri BN (1952) *The wealth of India, raw materials* (Vol. X). New Delhi: Council for Scientific and Industrial Research.
- Scalbert A, Manach C, Morand C, Remesy C, & Jimenez L (2005) Dietary polyphenols and the prevention of diseases. *Critical Reviews in Food Science and Nutrition*. **45**(4): 287-306
- Scartezzini P, & Speroni E (2000) Review on some plants of Indian traditional medicine with antioxidant activity. *Journal of Ethnopharmacology*. **71**(1-2): 23-43
- Schippmann U, Leaman D, & Cunningham AB (2006) Plants as source of medicines: new perspectives. In Bogers RJ, Craker LE & Lange D (Eds.), *Medicinal and aromatic plants - agricultural, commercial, ecological, legal, pharmacological and social aspects* (pp. 75). Dordrecht: Springer.
- Sermakkani M, & Thangapandian V (2012) GC-MS analysis of *Cassia italica* leaf methanol extract. *Asian Journal of Pharmaceutical and Clinical Research*. **5**(2): 90-94
- Shahid S, Riaz T, & Asghar MN (2015) Screening of *Ranunculus sceleratus* for enzyme inhibition, antibacterial and antioxidant activities. *Bangladesh Journal of Pharmacology*. **10**: 436-442
- Shahnaj S, Asha U, Mim T, Rumi NSH, Akter S, Ghose SR, Akter S, Islam MT, Das PR, & Rahmatullah M (2015) A survey on the ethnomedicinal practices of a folk medicinal practitioner in Manikganj district, Bangladesh. *Journal of Chemical and Pharmaceutical Research*. **7**(8): 690-696
- Shakeel-U-Rehman, Khan R, Bhat KA, Raja AF, Shawl AS, & Alam MS (2010) Isolation, characterisation and antibacterial activity studies of coumarins from *Rhododendron lepidotum* Wall. ex G. Don, Ericaceae. *Revista Brasileira de Farmacognosia*. **20**(6): 886-890
- Shakeel-U-Rehman, Sofi SN, Khuroo MA, Taneja SC, Bhat KA, & Vishwakarma R (2013) New natural compounds from *Rhododendron lepidotum*. *Natural Product Research*. **27**(21): 2033-2038
- Shao JH, Zhao CC, & Zhao R (2013) A new cerebroside from *Anemone rivularis*. *Chemistry of Natural Compounds*. **49**(4): 694-695
- Sharififar F, Dehghn-Nudeh G, & Mirtajaldini M (2008) Major flavonoids with antioxidant activity from *Teucrium polium* L. *Food Chemistry*. **112**(4): 885-888
- Sharma HK, Chhangte L, & Dolui AK (2001) Traditional medicinal plants in Mizoram, India. *Fitoterapia*. **72**(2): 146-161
- Sharma KK, Kotoky J, Kalita JC, & Barthakur R (2012) Evaluation of antidermatophytic activity of *Ranunculus sceleratus* and *Pongamia pinnata* available in North Eastern Region of India. *Asian Pacific Journal of Tropical Biomedicine*. **2**(2): S808-S811

- Sharma N, Gupta PC, Singh A, & Rao CV (2013) Brine shrimp bioassay of *Pentapetes phoenicea* Linn. and *Ipomoea carnea* Jacq. leaves. *Der Pharmacia Lettre*. **5**(1): 162-167
- Sharma US, Sharma UK, Singh A, Sutar N, & Singh PJ (2010) Screening of *Terminalia bellirica* fruits extracts for its analgesic and antipyretic activities. *Jordan Journal of Biological Sciences*. **3**(3): 121-124
- Shen CC, Cheng JJ, Lay HL, Wu SY, Ni CL, Teng CM, & Chen CC (2012) Cytotoxic apigenin derivatives from *Chrysopogon aciculatis*. *Journal of Natural Products*. **75**(2): 198–201
- Shih KS, Wang JH, Wu YW, Teng CM, Chen CC, & Yang CR (2012) Aciculatin inhibits granulocyte colony-stimulating factor production by human interleukin 1 β -stimulated fibroblast-like synoviocytes. *PLoS ONE*. **7**(7): e42389
- Shimada K, Fujikawa K, Yahara K, & Nakamura T (1992) Antioxidative properties of xanthone on the auto oxidation of soybean in cyclodextrin emulsion. *Journal of Agricultural and Food Chemistry*. **40**: 945–948
- Shimada T (2006) Salivary proteins as a defense against dietary tannins. *Journal of Chemical Ecology*. **32**(6): 1149-1163
- Shinwari ZK, Khan AA, & Nakaike T (2003) *Medicinal and other useful plants of District Swat Pakistan*. Peshawar, Pakistan: WWF-Pakistan.
- Shrestha GL, & Shrestha B (1999) An overview of wild relatives of cultivated plants in Nepal. In Shrestha R & Shrestha B (Eds.), *Wild Relatives of Cultivated Plants in Nepal. Proceedings of National Conference on Wild Relatives of Cultivated Plants in Nepal, June 2–4, 1999* (pp. 19-23). Kathmandu: Green Energy Mission.
- Shrestha KK, Tiwari NN, & Ghimire SK (2002) Medicinal and aromatic plants database of Nepal. In Watanabe T, Takano A, Bista MS & Saiju HK (Eds.), *Proceeding of Nepal-Japan Joint Symposium on Conservation and Utilization of Himalayan Medicinal Plant Resources, Nov 6–11, 2000, Kathmandu, Nepal* (pp. 53-74). Japan: Society for the conservation and Development of Himalayan Medicinal Resources.
- Shrestha P (1988) Contribution to the ethnobotany of the Tamangs of Kathmandu valley. *Contribution to the Nepalese studies*. **15**(2): 247-266
- Shrestha P (1989) Ethnobotanical observation on the Tamangs of Kathmandu valley *Proceedings of National Conference on Science and Technology, 24-29 April 1988* (pp. 353-358). Kathmandu: Royal Nepal Academy of Science and Technology.
- Siddhuraju P, & Becker K (2003) Studies on antioxidant activities of mucuna seed (*Mucuna pruriens* var *utilis*) extract and various non-protein amino/imino acids through *in vitro* models. *Journal of the Science of Food and Agriculture* **83**(14): 1517-1524
- Sikkema J, de Bont JA, & Poolman B (1994) Interactions of cyclic hydrocarbons with biological membranes. *The Journal of Biological Chemistry*. **269**(11): 8022–8028
- Sikkema J, de Bont JA, & Poolman B (1995) Mechanisms of membrane toxicity of hydrocarbons. *Microbiological Reviews*. **59**(2): 201-222
- Silva RO, Sousa FB, Damasceno SR, Carvalho NS, Silva VG, Oliveira FR, Sousa DP, Aragão KS, Barbosa AL, Freitas RM, & Medeiros JV (2014) Phytol, a diterpene alcohol, inhibits the inflammatory response by reducing cytokine production and oxidative stress. *Fundamental & Clinical Pharmacology*. **28**(4): 455-464

- Simoneit BRT, Schauer JJ, Nolte CG, Oros DR, Elias VO, Fraser MP, Rogge WF, & Cass GR (1999) Levoglucosan, a tracer for cellulose in biomass burning and atmospheric particles. *Atmospheric Environment*. **33**(2): 173-182
- Singh AK, Raghubanshi AS, & Singh JS (2002) Medical ethnobotany of the tribals of Sonaghati of Sonbhadra district, Uttar Pradesh, India. *Journal of Ethnopharmacology*. **81**(1): 31-41
- Singh I, Sharma A, Nunia V, & Goyal PK (2005) Radioprotection of Swiss albino mice by *Embllica officinalis*. *Phytotherapy Research*. **19**(5): 444-446
- Singh M, Chaudhry MA, Yadava JN, & Sanyal SC (1992) The spectrum of antibiotic resistance in human and veterinary isolates of *Escherichia coli* collected from 1984-86 in northern India. *Journal of Antimicrobial Chemotherapy*. **29**(2): 159-168
- Singh NP, Khanna KK, Mudgal V, & Dixit RD (2001) *Flora of Madhya Pradesh* (Vol. 3). Howrah: Botanical Survey of India.
- Singh U, & Jialal I (2006) Oxidative stress and atherosclerosis. *Pathophysiology*. **13**(3): 129–142
- Singleton VL, Orthofer R, & Lamuela-Raventos RM (1999) Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*. **299**: 152-178
- Singleton VL, & Rossi JA (1965) Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*. **16**: 144-158
- Sini KR, Haribabu Y, Sajith MS, & Surya SK (2012) *In-vitro* cytotoxic activity of *Orthosiphon thymiflorus* (Roth.) sleensen leaf extract against dalton lymphoma ascites cell line. *Journal of Chemical and Pharmaceutical Research*. **4**(1): 917-921
- Sivakumar T, & Gajalakshmi D (2014) Phytochemical screening and GC-MS analysis of root extract from *Asparagus racemosus* L. *International Journal of Pharmaceutical Science and Research*. **5**(12): 5245-5249
- Slater TF, Sawyer B, & Strauli U (1963) Studies of succinate-tetrazolium reductase systems III. Points of coupling of four different tetrazolium salts. *Biochimica et Biophysica Acta*. **77**: 383-393
- Sleet RB, & Brendel K (1983) Improved methods for harvesting and counting synchronous populations of *Artemia* nauplii for use in developmental toxicology. *Ecotoxicology and Environmental Safety*. **7**(5): 435-446
- Smith MA, Rottkamp CA, Nunomura A, Raina AK, & Perry G (2000) Oxidative stress in Alzheimer's disease. *Biochimica et Biophysica Acta*. **1502**(1): 139–144
- Snider BB, & Amin SG (1978) A synthetic precursor of verrucarin. *Synthetic Communications*. **8**: 117-125
- Soares JR, Dins TC, Cunha AP, & Almeida LM (1997) Antioxidant activities of some extracts of *Thymus zygis*. *Free Radical Research*. **26**(5): 469–478
- Solanki SS, & Selvanayagam M (2013) Phytochemical screening and study of predictive toxicity of certain medicinal plants and extracts using brine shrimp. *Journal of Herbal Science and Technology*. **10**(1): 1-4
- Solis PN, Wright CW, Anderson MM, Gupta MP, & Phillipson JD (1993) A microwell cytotoxicity assay using *Artemia salina* (brine shrimp). *Planta Medica*. **59**(3): 250-252

- Soobrattee MA, Neerghen VS, Luximon-Ramma A, Aruoma OI, & Bahorun T (2005) Phenolics as potential antioxidant therapeutic agents: mechanism and actions. *Mutation Research*. **579**(1-2): 200-213
- Sood K, & Yadav RNS (2014) Phytochemical screening and antimicrobial activity of four members of family Apiaceae. *International Journal of Phytomedicine*. **6**(2): 232-236
- Sreekumar PV, & Nair VJ (1991) *Flora of Kerala - grasses*. Calcutta: Botanical Survey of India.
- Sridevi G, Prema S, Sekar S, & Sembulingam K (2014) Phytochemical analysis of *Pergularia daemia* for its bioactive components through gas chromatographic mass spectrometry (GCMS). *IOSR Journal Of Pharmacy*. **4**(5): 41-46
- Srinivasan D, Nathan S, Suresh T, & Lakshmana PP (2001) Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine. *Journal of Ethnopharmacology*. **74**(3): 217-220
- Sripanidkulchai B, & Junlatat J (2014) Bioactivities of alcohol based extracts of *Phyllanthus emblica* branches: antioxidation, antimelanogenesis and anti-inflammation. *Journal of natural medicines*. **68**(3): 615-622
- Stamp D, Zhang XM, Medline A, Bruce WR, & Archer MC (1993) Sucrose enhancement of the early steps of colon carcinogenesis in mice. *Carcinogenesis*. **14**(4): 777-779
- Stefanello MEA, Cervi AC, Ito IY, Salvador MJ, Wisniewski A, & Simionatto EL (2008) Chemical composition and antimicrobial activity of essential oils of *Eugenia chlorophylla* (Myrtaceae). *Journal of Essential Oil Research*. **20**(1): 75-78
- Stefanello MEA, Salvador MJ, Ito IY, & Macari PAT (2006) Avaliação da atividade antimicrobiana e citotóxica de extratos de *Gochnatia polymorpha* ssp. *floccosa*. *Revista Brasileira de Farmacognosia*. **16**(4): 525-530
- Steffen LM, Jacobs DR, Stevens J, Shahar E, Carithers T, & Folsom AR (2003) Associations of whole-grain, refined-grain, and fruit and vegetable consumption with risks of all-cause mortality and incident coronary artery disease and ischemic stroke: the atherosclerosis risk in communities (ARIC) study. *The American Journal of Clinical Nutrition*. **78**(3): 383-390
- Stevigny C, Bailly C, & Quetin-Leclercq J (2005) Cytotoxic and antitumor potentialities of aporphinoid alkaloids. *Current Medicinal Chemistry - Anti-Cancer Agents*. **5**(2): 173-182
- Strack D, & Wray V (1992) Anthocyanins. In Harborne JB (Ed.), *The Flavonoids: Advances in Research Since 1986* (pp. 1-22). London: Chapman and Hall.
- Subhasree B, Baskar R, Keerthana RL, Susan RL, & Rajasekaran P (2009) Evaluation of antioxidant potential in selected green leafy vegetables. *Food Chemistry*. **115**(4): 1213-1220
- Sudha T, Chidambarampillai S, & Mohan VR (2013) GC-MS analysis of bioactive components of aerial parts of *Fluggea leucopyrus* Willd. (Euphorbiaceae). *Journal of Applied Pharmaceutical Science*. **3**(5): 126-130
- Sui L, Dong Y, Watanabe Y, Yamaguchi F, Hatano N, Izumori K, & Tokuda M (2005a) Growth inhibitory effect of D-allose on human ovarian carcinoma cells *in vitro*. *Anticancer Research*. **25**(4): 2639-2644

- Sui L, Dong Y, Watanabe Y, Yamaguchi F, Hatano N, Tsukamoto I, Izumori K, & Tokuda M (2005b) The inhibitory effect and possible mechanisms of D-allose on cancer cell proliferation. *International Journal of Oncology*. **27**(4): 907-912
- Sumantran VN, Boddul S, Koppikar SJ, Dalvi M, Wele A, Gaire V, & Wagh UV (2007) Differential growth inhibitory effects of *W. somnifera* root and *E. officinalis* fruits on CHO cells. *Phytotherapy Research*. **21**(5): 496–499
- Sun YX, Liu JC, & Liu DY (2011) Phytochemicals and bioactivities of *Anemone raddeana* Regel: a review. *Die Pharmazie*. **66**(11): 813-821
- Sundarraaj S, Thangam R, Sreevani V, Kaveri K, Gunasekaran P, Achiraman S, & Kannan S (2012) γ -Sitosterol from *Acacia nilotica* L. induces G2/M cell cycle arrest and apoptosis through c-Myc suppression in MCF-7 and A549 cells. *Journal of Ethnopharmacology*. **141**(3): 803-809
- Sunil C, Agastian P, Kumarappan C, & Ignacimuthu S (2012) *In vitro* antioxidant, antidiabetic and antilipidemic activities of *Symplocos cochinchinensis* (Lour.) S. Moore bark. *Food and Chemical Toxicology*. **50**(5): 1547-1553
- Takahashi H, Hirata S, Minami H, & Fukuyama Y (2001) Triterpene and flavanone glycoside from *Rhododendron simsii*. *Phytochemistry*. **56**(8): 875-879
- Takhtajan A (1997) *Diversity and classification of flowering plants*. New York: Columbia University Press.
- Tang M, Shen D, Hu YC, Gao S, & Yu S (2004) Cytotoxic triterpenoid saponins from *Symplocos chinensis*. *Journal of Natural Products*. **67**(12): 1969-1974
- Tang W, & Eisenbrand G (1992) *Rhododendron dauricum* L. *Chinese drugs of plant origin: chemistry, pharmacology, and use in traditional and modern medicine* (pp. 877-883). Berlin, Heidelberg: Springer Berlin Heidelberg.
- Tawaha K, Alali FQ, Gharaibeh M, Mohammad M, & El-Elmat T (2007) Antioxidant activity and total phenolic content of selected Jordanian plant species. *Food Chemistry*. **104**(4): 1372-1378
- Temitope AO, Olufemi AG, & Alaba FT (2010) Effect of heat treatment on antioxidant activity of some spices. *Continental Journal of Food Science and Technology*. **4**: 53–59
- Tennyson S, Ravindran KJ, & Arivoli S (2012) Screening of twenty five plant extracts for larvicidal activity against *Culex quinquefasciatus* Say (Diptera: Culicidae). *Asian Pacific Journal of Tropical Biomedicine*. **2**(2): S1130-1134
- Teres S, Barcelo-Coblijn G, Benet M, Alvarez R, Bressani R, Halver JE, & Escriba PV (2008) Oleic acid content is responsible for the reduction in blood pressure induced by olive oil. *Proceedings of the National Academy of Sciences*. **105**(37): 13811-13816
- Tewari K, Pande C, Tewari G, Kharkwal GC, & Punetha D (2015) Volatile constituents of *Geranium wallichianum* D. Don ex Sweet. from northwestern Himalayan region. *Journal of Indian Chemical Society*. **92**: 123-125
- Thakur RS, Puri HS, & Akhtar H (1989) *Major medicinal plants of India*. Lucknow: Central Institute of Medicinal and Aromatic Plants.
- Thanigaivel S, Hindu SV, Vijayakumar S, Mukherjee A, Chandrasekaran N, & Thomas J (2015) Differential solvent extraction of two seaweeds and their efficacy in controlling *Aeromonas salmonicida* infection in *Oreochromis mossambicus*: a novel therapeutic approach. *Aquaculture*. **443**: 56-64

- Thatoi HN, Patra JK, & Das SK (2014) Free radical scavenging and antioxidant potential of mangrove plants: a review. *Acta Physiologiae Plantarum*. **36**(3): 561-579
- Thery JC, Spano JP, Azria D, Raymond E, & Penault LF (2014) Resistance to human epidermal growth factor receptor type 2-targeted therapies. *European Journal of Cancer*. **50**(5): 892-901
- Thresher JS, Podolin DA, Wei Y, Mazzeo RS, & Pagliassotti MJ (2000) Comparison of the effects of sucrose and fructose on insulin action and glucose tolerance. *American Journal of Physiology*. **279**(4): R1334–R1340
- Tiwari KP, & Singh RB (1978) Rivularinin, a new saponin from *Anemone rivularis*. *Phytochemistry*. **17**(11): 1991–1994
- Tiwari NN (1999) Wild relatives of cultivated medicinal and aromatic plants (MAPs) in Nepal. *Proceedings of National Conference on Wild Relatives of Cultivated Plants in Nepal, June 2–4, 1999* (pp. 141-148). Nepal: The Green Energy Mission.
- Tiwari ON, & Chauhan UK (2005) Genus *Rhododendron* status in Sikkim Himalaya: an assessment. *Journal American Rhododendron Society*. **59**: 147–153
- Tuomi T, Reijula K, Johnsson T, Hemminki K, Hintikka EL, Lindroos O, Kalso S, Koukila-Kähkölä P, Mussalo-Rauhamaa H, & Haahtela T (2000) Mycotoxins in crude building materials from water-damaged buildings. *Applied and Environmental Microbiology*. **66**(5): 1899–1904
- Upadhyay N, Tiwari SK, Srivastava A, Seth A, & Maurya SK (2015) Anti-urolithiatic effect of *Terminalia bellirica* Roxb. fruits on ethylene glycol-induced renal calculi in rats. *Indo American Journal of Pharmaceutical Research*. **5**(5): 2031-2040
- Upston JM, Kritharides L, & Stocker R (2003) The role of vitamin E in atherosclerosis. *Progress in Lipid Research*. **42**(5): 405-422
- Vanhaecke P, Persoone G, Claus C, & Sorgeloos P (1981) Proposal for a short-term toxicity test with *Artemia nauplii*. *Ecotoxicology and Environmental Safety*. **5**(3): 382-387
- Vanin AB, Orlando T, Piazza SP, Puton BM, Cansian RL, Oliveira D, & Paroul N (2014) Antimicrobial and antioxidant activities of clove essential oil and eugenyl acetate produced by enzymatic esterification. *Applied Biochemistry and Biotechnology*. **174**(4): 1286-1298
- Velanganni J, Kadamban D, & Ramamoorthy D (2011) Phytochemical screening and antimicrobial activity of the stem of *Mallotus philippensis* (Lam.) Muell. Arg. Var. *philippensis*. *International Journal of Pharmacy and Pharmaceutical Sciences*. **3**(2): 160-163
- Veni T, & Pushpanathan T (2014) Comparison of the *Artemia salina* and *Artemia franciscana* bioassays for toxicity of Indian medicinal plants. *Journal of Coastal Life Medicine*. **2**(6): 453-457
- Venkata RB, Samuel LA, Pardha SM, Narashimha RB, Naga VKA, Sudhakar M, & Radhakrishnan TM (2012) Antibacterial, antioxidant activity and GC-MS analysis of *Eupatorium odoratum*. *Asian Journal of Pharmaceutical and Clinical Research*. **5**(2): 99-106
- Verpoorte R (2000) Pharmacognosy in the new millennium: leadfinding and biotechnology. *Journal of Pharmacy and Pharmacology*. **52**(3): 253-262
- Vijayabaskaran M, Badkhal AK, Babu G, Sivakumar P, Perumal P, Sivakumar T, Sampathkumar R, & Jayakar B (2010) Antitumor activity and antioxidant status of *Symplocos*

- racemosa* Roxb against Ehrlich ascites carcinoma in Swiss albino mice. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. **1**(3): 306-314
- Vinoth PV, Chidambaranathan N, & Gopal V (2012) Evaluation and quantification of angiogenesis activity of *Terminalia bellirica* Roxb. by mice sponge implantation method. *Journal of Young Pharmacists*. **4**(1): 22-27
- Vivancos M, & Moreno JJ (2005) beta-Sitosterol modulates antioxidant enzyme response in RAW 264.7 macrophages. *Free Radical Biology and Medicine*. **39**(1): 91-97
- Wadekar RR, & Patil KS (2008) Effect of *Ziziphus mauritiana* leaf extracts on phagocytosis by human neutrophils. *Journal of Pharmacy Research*. **1**(1): 79-83
- Wan-Ibrahim WI, Sidik K, & Kuppusamy UR (2010) A high antioxidant level in edible plants is associated with genotoxic properties. *Food Chemistry*. **122**(4): 1139-1144
- Wang HX, & Ng TB (1999) Natural products with hypoglycemic, hypotensive, hypocholesterolemic, antiatherosclerotic and antithrombotic activities. *Life Sciences*. **65**(25): 2663-2677
- Wang JL, Liu K, Gong WZ, Wang Q, Xu DT, Liu MF, Bi KL, & Song YF (2012) Anticancer, antioxidant, and antimicrobial activities of anemone (*Anemone cathayensis*). *Food Science and Biotechnology*. **21**(2): 551-557
- Wang JR, Ma JQ, Peng LL, Feng JT, & Ding LS (1998) Chemical constituents and antifeeding activity of *Anemone tomentosa*. *Acta Botanica Boreali-Occidentalia Sinica*. **18**: 643-644
- Wang MK, Wu FE, & Chen YZ (1997) Triterpenoid saponins from *Anemone hupehensis*. *Phytochemistry*. **44**(2): 333-335
- Wang X, Wang M, Xu M, Wang Y, Tang H, & Sun X (2014) Cytotoxic oleanane-type triterpenoid saponins from the Rhizomes of *Anemone rivularis* var. *flore-minore*. *Molecules*. **19**(2): 2121-2134
- Watt G (1972) *Dictionary of the economic products of India*. Delhi: Cosmo Publications.
- Wendakoon C, Calderon P, & Gagnon D (2012) Evaluation of selected medicinal plants extracted in different ethanol concentrations for antibacterial activity against human pathogens. *Journal of Medicinally Active Plants*. **1**(2): 60-68
- WHO (2000) General guidelines for methodologies on research and evaluation of traditional medicine (pp. 41). Geneva: World Health Organization.
- Wilkes HG (1981) *New and potential crops or what to anticipate for the future. Paper presented at the annual meeting of the American Association for the Advancement of Science, January 1981, Toronto, Canada.*
- Wilson TG, Landers MH, & Happ GM (1983) Precocene I and II inhibition of vitellogenic oocyte development in *Drosophila melanogaster*. *Journal of Insect Physiology*. **29**(3): 249-254
- Wu G, Yu D, Tu M, Xia S, Chong Y, & Kang H (1999) Growth and population ecology of *Ranunculus sceleratus* in Donghu Lake, Wuhan. *Acta Hydrobiologica Sinica*. **23**(3): 211-216
- Xiao Y, Fan Y, Chen B, Zhang Q, & Zeng H (2010) Polysaccharides from *Scurrula parasitica* L. inhibit sarcoma S180 growth in mice. *Zhongguo Zhong Yao Za Zhi*. **35**(3): 381-384
- Xu BJ, & Chang SKC (2007) A comparative study on phenolic profiles and antioxidant activities of legumes as affected by extraction solvents. *Journal of Food Science*. **72**(2): 160-161

- Xu Q, Li YH, & Lu XY (2007) Investigation on influencing factors of 5-HMF content in Schisandra. *Journal of Zhejiang University-Science B*. **8**(6): 439–345
- Xu XF, Yang H, & Yang LT (2004) The applying potential of purifying sewage with *Ranunculus sceleratus* L. *Journal of Plant Resources and Environment*. **13**: 17-20
- Yaguchi A, Yoshinari T, Tsuyuki R, Takahashi H, Nakajima T, Sugita-Konishi Y, Nagasawa H, & Sakuda S (2009) Isolation and identification of precocenes and piperitone from essential oils as specific inhibitors of trichothecene production by *Fusarium graminearum*. *Journal of Agricultural and Food Chemistry*. **57**(3): 846-851
- Yannai S (2004) *Dictionary of food compounds with CD-ROM: Additives, flavors, and ingredients*. Boca Raton: Chapman & Hall/CRC.
- Yasuda T, Inaba A, Ohmori M, Endo T, Kubo S, & Ohsawa K (2000) Urinary metabolites of gallic acid in rats and their radical-scavenging effects on 1,1-diphenyl-2-picrylhydrazyl radical. *Journal of Natural Products*. **63**(10): 1444-1446
- Ye WC, Zhang QW, Zhao SX, & Che CT (2001) Four new oleanane saponins from *Anemone anhuiensis*. *Chemical & Pharmaceutical Bulletin*. **49**(5): 632-634
- Yu F, Yu F, McGuire PM, Li R, & Wang R (2007) Effects of *Hydrocotyle sibthorpioides* extract on transplanted tumors and immune function in mice. *Phytomedicine*. **14**(2-3): 166-171
- Yuvaraj G, Sathyanathan V, Shankar S, & Kumar NR (2010) Anti cancer and *in-vitro* activities of *Derris brevipes* var. *brevipes*. *Journal of Chemical and Pharmaceutical Research*. **2**(6): 482-488
- Zablotowicz RM, Hoagland RE, & Wagner SC (1996) Effect of saponins on the growth and activity of rhizosphere bacteria. *Advances in Experimental Medicine and Biology*. **405**: 83-95
- Zeng K, Thompson KE, Yates CR, & Miller DD (2009) Synthesis and biological evaluation of quinic acid derivatives as anti-inflammatory agents. *Bioorganic & Medicinal Chemistry Letters*. **19**(18): 5458–5460
- Zhang LT, Zhang YW, Takaishi Y, & Duan HQ (2008) Antitumor triterpene saponins from *Anemone flaccida*. *Chinese Chemical Letters*. **19**(2): 190-192
- Zhang LZ, Zhao WH, Guo YJ, Tu GZ, Lin S, & Xin LG (2003) Studies on chemical constituents in fruits of Tibetan medicine *Phyllanthus emblica*. *Zhongguo Zhong Yao Za Zhi*. **28**(10): 940–943
- Zhang YJ, Liang RJ, Zhao Q, Hong AH, Wang YF, & Cen YZ (2013) Chemical constituents from the fresh leaves of *Phyllanthus emblica* L. *Lishizhen Medicine and Materia Medica Research*. **24**(6): 1298-1300
- Zhang YJ, Tanaka T, Iwamoto Y, Yang CR, & Kouno I (2000) Phyllaemblic acid, a novel highly oxygenated norbisabolane from the roots of *Phyllanthus emblica*. *Tetrahedron Letters*. **41**(11): 1781-1784
- Zhao C, Li XN, Liang YZ, Fang HZ, Huang LF, & Guo FQ (2006) Comparative analysis of chemical components of essential oils from different samples of *Rhododendron* with the help of chemometrics methods. *Chemometrics and Intelligent Laboratory Systems*. **82**(1-2): 218-228
- Zhao CC, Shao JH, & Fan JD (2012) A new triterpenoid with antimicrobial activity from *Anemone rivularis*. *Chemistry of Natural Compounds*. **48**(5): 803-805

- Zhao YX, Li CS, Luo XD, Liu YQ, & Zhou J (2005) A new sesquilignan from *Tsuga dumosa*. *Acta Botanica Yunnanica*. **27**(2): 217-222
- Zhao YX, Li CS, Luo XD, & Zhou J (2004a) A new tricyclic 6-7-5-ring diterpene from pliocene fossil *Tsuga dumosa*. *Heterocycles*. **63**(4): 861-864
- Zhao YX, Luo XD, & Zhou J (2004b) Lignans from *Tsuga dumosa*. *Acta Botanica Yunnanica*. **26**(2): 229-233
- Zhou X, Zhang Y, Liang H, Huang S, Wang C, & Ping A (2012) Chemical constituents of *Rhododendron lepidotum*. *Zhongguo Zhong Yao Za Zhi*. **37**(4): 483-489

Appendices

List of Reagents and Culture media

1. Preparation of 1 M Na₂CO₃ – 100 ml

10.599 g of Na₂CO₃ (Merk specialities Pvt. Ltd., Mumbai, India) was carefully weighed and then dissolved in distilled water, and the volume was adjusted to 100 ml at the end.

2. Preparation of glacial acetic acid (20%) – 200

40 ml of the commercially supplied glacial acetic acid (Thermo Fisher Scientific, India) was taken and mixed with ethanol. Finally, the volume was adjusted to 200 ml by the addition of ethanol.

3. Preparation of aluminium chloride (10%) – 100 ml

10 g of the commercially supplied aluminium chloride (Merk Specialities Pvt. Ltd., Mumbai, India) was weighed and dissolved in water. Finally, the volume was maintained to 100 ml.

4. Preparation of 1M potassium acetate (CH₃COOK) – 100 ml

Weigh 9.814 g of the potassium acetate Merk Specialities Pvt. Ltd., Mumbai, India) and dissolve on water. Finally maintain the volume to 100 ml by the addition of water.

5. Preparation of 0.2 Mm DPPH solution – 100 ml

1,1-diphenyl-2-picrylhydrazyl (DPPH) has the molecular weight of 394.32 g/mol. Thus, 100 ml of 0.2 mM solution of DPPH is prepared by weighing the 7.886 mg of the DPPH carefully and dissolving it on ethanol and finally maintaining the volume to 100 ml by addition of ethanol.

6. Preparation of the Folin-Ciocalteu pheol reagent (1:10 dilution)

6 ml of the commercially supplied Folin – Ciocalteu phenol reagent (Merk Specialities Pvt. Ltd., Mumbai, India) was taken and mix it with 54 ml of the distilled water to prepare the 60 ml of 1:10 dilution of Folin – Ciocalteu phenol reagent.

7. Composition of nutrient agar media

The composition of nutrient agar media (Hi Media Laboratories Pvt. Ltd., Mumbai, India) is as follows:

Components	gram/L
Peptic digest of animal tissue	5.0
Beef extract	1.5
Yeast extract	1.5
Sodium chloride	5.0
Agar	15.0
Ph	7.4±0.2

8. Composition of Mueller Hinton Agar (MHA)

The composition of Mueller Hinton Agar (MHA) media (Hi Media Laboratories Pvt. Ltd., Mumbai, India) is as follows:

Components	gram/L
Beef infusion form	300
Casein hydrolysate	17.5
Starch	1.56
Agar	17
Final pH	7.3±0.2

9. Composition of Potato Dextrose Agar (PDA)

The composition of PDA broth (Hi Media Laboratories Pvt. Ltd., Mumbai, India) is as follows:

Components	gram/L
Potato	200
Agar	2
Dextrose/Glucose	2

10. Composition of artificial sea water

SN	Salt	Molecular wt (g/mol)	Amount in gram/L
1	NaCl	58.44	23.50
2	Na ₂ SO ₄	142.04	4.00
3	KCl	74.55	0.68
4	H ₃ BO ₃	61.83	0.026
5	MgCl ₂ .6H ₂ O	203.33	10.78
6	CaCl ₂	110.98	1.47
7	NaHCO ₃	84.00	0.196
8	Na ₂ EDTA	372.24	0.0003

pH = 8.0±0.2

11. General Classification of brine shrimp cytotoxicity

LC ₅₀ Values (µg/ml)	Classification
<1	Extremely toxic
1-100	Highly toxic
100-500	medium toxic
500-1000	low toxic
>1000	non-toxic

Photographs

Medicinal Plants



a) *Loxogramme involuta*



b) *Hydrocotyle javanica* Thunb



c) *Phyllanthus emblica* Linnaeus



d) *Chrysopogon aciculatus* (Retzius) Trinius

Plate 1 Some medicinal plants (a) *Loxogramme involuta*, (b) *Hydrocotyle javanica* Thunb, (c) *Phyllanthus emblica* Linnaeus, and (d) *Chrysopogon aciculatus* (Retzius) Trinius



a) *Rhododendron lepidotum* Wallich ex G. Don



b) *Geranium wallichianum* D. Don ex Sweet



c) *Tsuga dumosa* (D. Don) Eichler



d) *Anemone rivularis* Buchanan-Hamilton ex De Candolle

Plate 2 Some medicinal plants (a) *Rhododendron lepidotum* Wallich ex G. Don, (b) *Geranium wallichianum* D. Don ex Sweet, (c) *Tsuga dumosa* (D. Don) Eichler, and (d) *Anemone rivularis* Buchanan-Hamilton ex De Candolle



a) *Scurrula elata* (Edgeworth) Danser



b) *Terminalia bellirica* (Gaertner) Roxburgh



c) *Ranunculus scleratus* Linnaeus



d) *Symplocos lucida* (Thunberg) Siebold & Zuccarini

Plate 3 Some medicinal plants (a) *Scurrula elata* (Edgeworth) Danser, (b) *Terminalia bellirica* (Gaertner) Roxburgh, (c) *Ranunculus scleratus* Linnaeus, and (d) *Symplocos lucida* (Thunberg) Siebold & Zuccarini

Methanol extraction



Methanol extraction



Research team members with principal investigators


The project on methanol extraction for bioprospecting of underutilised medicinal plants of Nepal

GC-MS analysis




GC-MS analysis: the training and the experiment

Brine Shrimp Lethality




Brine shrimp lethality bioassay: death count through stereoscope


MTT Assay



HeLa cell line




MDCK cell line



Cell seeding in 96-well tissue culture plate



Examination of effects of extracts on HeLa cells



Formation of formazan crystals



Dissolution of formazan salts

MTT cell proliferation assay: cell seeding to formazan dissolution